

GLYCERIN - glycerin suppository
Rite Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rite Aid Glycerin Supp.

Active ingredient (in each suppository)

Glycerin 2 g

Purpose

Hyperosmotic laxative

Uses

- for relief of occasional constipation
- this product generally produces a bowel movement in 1/4 to 1 hour

Warnings

For rectal use only

May cause rectal discomfort or burning sensation

Ask a Doctor before using any laxative if you have

- abdominal pain, nausea or vomiting
- a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

Stop use and consult a doctor if you have

- rectal bleeding
- no bowel movement after using this product

These symptoms may indicate a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Single daily dosage

adults and children 6 years and over 1 suppository, or as directed by a doctor

children 2 to under 6 years use Child Suppositories

Insert suppository well up into rectum.

Suppository need not melt completely to produce laxative action.

Other information

- Store container tightly closed.
- Keep away from excessive heat.

Inactive ingredients

purified water, sodium hydroxide, stearic acid

Glycerin Suppositories, 50 count

Compare to the active ingredient in Fleet® Glycerin Suppositories*

LAXATIVE GLYCERIN SUPPOSITORIES

GLYCERIN 2g

often works in minutes for gentle, effective relief of constipation

RITE AID PHARMACY **50 ADULT SIZE SUPPOSITORIES**

| | | |
|--|---|--|
| Drug Facts | Active Ingredient (in each suppository) Glycerin 2 g..... | Purpose Hyperosmotic laxative |
| Uses | <ul style="list-style-type: none"> ■ for relief of occasional constipation ■ this product generally produces bowel movement in 1/4 to 1 hour | |
| Warnings | <p>For rectal use only</p> <p>May cause rectal discomfort or a burning sensation.</p> <p>Ask a doctor before using any laxative if you have</p> <ul style="list-style-type: none"> ■ abdominal pain, nausea or vomiting ■ sudden change in bowel habits lasting more than 2 weeks ■ already used a laxative for more than 1 week <p>Stop use and consult a doctor if you have</p> <ul style="list-style-type: none"> ■ rectal bleeding ■ no bowel movement after using product <p>These symptoms may indicate a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> | |
| Directions | <i>Single daily dosage</i> | adults and children 6 years and over use Child Suppositories |
| | children 2 to under 6 years | use Child Suppositories |
| Insert suppository well up into rectum. Suppository need not melt completely to produce laxative action. | | |
| Other Information | <ul style="list-style-type: none"> ■ Store container tightly closed. ■ Keep away from excessive heat. | |
| Inactive Ingredients | purified water, sodium hydroxide, stearic acid | |

SATISFACTION GUARANTEE
If you're not satisfied, we'll happily refund your money.

DISTRIBUTED BY:
RITE AID
30 HUNTER LANE
CAMP HILL, PA 17011
www.riteaid.com

*This product is not manufactured or distributed by C.B. Fleet Co., owner of the registered trademark Fleet®

0 11822 33215 6

MOUTH OF JAR SEALED FOR SAFETY. IF FOIL IS BROKEN OR MISSING, DO NOT USE.

NO VARNISH AREA

The product package shown above represents a sample of that currently in use. Additional packaging may also be available.

RAGlycerinSupp50ct

Distributed by

Rite Aid

30 Hunter Lane

Camp Hill, PA 17011

GLYCERIN

glycerin suppository

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:11822-3312 |
| Route of Administration | RECTAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX) | GLYCERIN | 2 g |

Inactive Ingredients

| Ingredient Name | Strength |
|-------------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |
| SODIUM HYDROXIDE (UNII: 55X04QC32I) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:11822-3312-9 | 100 in 1 JAR; Type 0: Not a Combination Product | 10/14/2011 | |
| 2 | NDC:11822-3312-5 | 50 in 1 JAR; Type 0: Not a Combination Product | 10/14/2011 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part334 | 10/14/2011 | |

Labeler - Rite Aid Corporation (014578892)**Registrant** - DSC Laboratories Inc. (097807374)**Establishment**

| Name | Address | ID/FEI | Business Operations |
|------------------------|---------|-----------|-------------------------|
| DSC Laboratories, Inc. | | 097807374 | manufacture(11822-3312) |

Revised: 11/2021

Rite Aid Corporation