**RITE AID STOMACH RELIEF ORIGINAL FLAVOR- bismuth subsalicylate tablet, chewable**
**RITE AID CORPORATION**

---

**Rite Aid Pink Bismuth Subsalicylate 262mg 30 Chewable Tablets**

## Active ingredient (in each tablet)

Bismuth subsalicylate 262 mg

## Purpose

Upset stomach reliever and antidiarrheal

## Uses

relieves:

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

***Ask a doctor or pharmacist before use if you are*** *taking any drug for*

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

**When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use**.**

**Keep Out of Reach of Children**

In case of overdose, get medical help or contact a Poison Control Center immediately at 1-800-222-1222

**Directions**

- chew completely or dissolve tablets in mouth. Do not swallow tablets whole.
- **adults and children 12 years and over:**
- 2 tablets (1 dose) every ½ hour or 4 tablets (2 doses) every hour as needed for diarrhea
- 2 tablets (1 dose) every ½ hour as needed for over indulgence (upset stomach, heartburn, indigestion, nausea.
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty clear fluids to help prevent dehydration caused by diarrhea.

**Other information**

- **each tablet contains:** salicylate 102 mg
- store at room temperature.
- avoid excessive heat

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF STRIP UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**Inactive ingredients**

D&C Red No. 27 lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, sorbitol, sucralose.

**Questions or comments?**

1-**866-467-2748**

**Principal Display**

**NDC# 11822-2430-0**

**Compare to the active ingredient in Pepto-Bismol® Chewables***

**Stomach Relief**

**Chewable Tablets**

Bismuth Subsalicylate

Upset Stomach Reliever / Antidiarrheal

**5 Symptoms Digestive Relief:**

- Heartburn
- Indigestion
- Nausea
- Upset Stomach
- Diarrhea

**Original Flavor**

**30 Chewable Tablets**

**DISTRIBUTED BY:**

**RITE AID, 30 HUNTER LANE,**

**CAMP HILL, PA 17011**

**www.riteaid.com**

**SATISFACTION GUARANTEE**

**If you’re not satisfied, we will happily refund your money.**

**IMPORTANT:** Keep this carton for future labeling.

*This product is not manufactured or distributed by Procter & Gamble Inc., the distributor of Pepto-Bismol® Chewables.

# RITE AID STOMACH RELIEF ORIGINAL FLAVOR

bismuth subsalicylate tablet, chewable

| Product Information | | | |
|---|---|---|---|
| **Product Type** | HUMAN OTC DRUG | **Item Code (Source)** | NDC:11822-2430 |
| **Route of Administration** | ORAL | | |

| Active Ingredient/Active Moiety | | |
|---|---|---|
| **Ingredient Name** | **Basis of Strength** | **Strength** |
| **BISMUTH SUBSALICYLATE** (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ) | BISMUTH SUBSALICYLATE | 262 mg |

| Inactive Ingredients | |
|---|---|
| **Ingredient Name** | **Strength** |
| **D&C RED NO. 27** (UNII: 2LRS185U6K) | |
| **MAGNESIUM STEARATE** (UNII: 70097M6I30) | |
| **MALTODEXTRIN** (UNII: 7CVR7L4A2D) | |

| | |
|---|---|
| **MICROCRYSTALLINE CELLULOSE** (UNII: OP1R32D61U) | |
| **SORBITOL** (UNII: 506T60A25R) | |
| **SUCRALOSE** (UNII: 96K6UQ3ZD4) | |

## Product Characteristics

| | | | |
|---|---|---|---|
| **Color** | PINK | **Score** | no score |
| **Shape** | ROUND | **Size** | 17mm |
| **Flavor** | | **Imprint Code** | RP124 |
| **Contains** | | | |

## Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:11822-2430-0 | 5 in 1 CARTON | 03/30/2020 | |
| **1** | | 6 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC Monograph Drug | M008 | 03/30/2020 | |

## Labeler - RITE AID CORPORATION (014578892)

Revised: 10/2024 RITE AID CORPORATION