

**MIGRAINE RELIEF- acetaminophen, aspirin, caffeine tablet, film coated
Walgreen Company**

Walgreen Co. Migraine Relief Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID*)

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever aid

Use

- treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for:
- diabetes
- gout
- arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients

carnauba wax, crospovidone, FD&C blue #1 aluminum lake, hypromellose, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, saccharin sodium, stearic acid, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Walgreens

Compare to the active ingredients in Excedrin[®] Migraine

WALGREENS PHARMACIST RECOMMENDED

Migraine Relief

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFEINE TABLETS

PAIN RELIEVER / PAIN RELIEVER AID

100 COATED CAPLETS**

(**capsule-shaped tablets)

ACTUAL SIZE

W3OR01221-F
REV-0522

Drug Facts

Active ingredients (in each caplet)
 Acetaminophen 250 mg
 Aspirin 250 mg (NSAID)
 Caffeine 65 mg

Purposes
 Pain reliever
 Pain reliever and
 nonsteroidal anti-inflammatory drug

Use ■ treats migraine

Warnings

Have's syndrome. Children and teenagers who have or are recovering from chicken pox or the-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Take aspirin may cause a severe allergic reaction which may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
 Liver damage may occur if you take
 more than 2 caplets in 24 hours, which is the maximum daily amount.

■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Serious allergic warning: This product contains aspirin. It may cause a severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take other drugs containing aspirin or aspirin drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 2 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING



37478 94 C14

Drug Facts (continued)

Medication versus headache warning: Headaches may worsen if this product is used for 10 or more days per month. **Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat.

Do not use
 ■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
 ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if

- you have a fever that migrates, diagnosed by a health professional
 - you have a headache that is different from your usual migraines
 - you have the worst headache of your life
 - you have ever had a stiff neck
 - you have headaches beginning after or caused by head injury, operation, coughing or bending
 - you experience your first headache after the age of 50
 - you have a migraine so severe as to require bed rest
 - you have liver disease
 - you have a history of stomach or kidney disease, such as heartburn
 - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 - you are taking a diuretic
 - you have asthma
 - you have problems or serious side effects from taking pain relievers or fever reducers
 - you have vomiting with your migraine headache
- Ask a doctor or pharmacist before use if you are**
- taking a prescription drug for:
 - diabetes ■ gout ■ arthritis
 - taking any other drug
 - taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Drug Facts (continued)

Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ your migraine is not relieved or worsens after first dose
 ■ new or unexpected symptoms occur
 ■ ringing in the ears or loss of hearing occurs

Important for use: Reading, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-2222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- adults take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age, ask a doctor

Other information

- Store at 20°-25° (68°-77°)
- Use cap tightly after use
- Read all product information before using. Keep this box for important information.

Inactive ingredients: carmellose gum, croscopolone, FD&C color #1 aluminum lake, hydroxytoluene, microcrystalline cellulose, polydioxanone, pregelatinized starch, propylene glycol, saccharin sodium, stearic acid, titanium dioxide

Questions or comments? 1-800-71-9-6280

KEEP BOX FOR REFERENCE

Your pharmacist recommends the Walgreens brand. We invite you to compare to national brands.
 *This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Excedrin® Migraine.
 DISTRIBUTED BY: WALGREENS CO., 200 WILMOT RD., DEERFIELD, IL 60015
 walgreens.com ©2021 Walgreen Co.



NDC 0363-0374-78



Migraine Relief

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFEINE TABLETS
 PAIN RELIEVER / PAIN RELIEVER AID

100 COATED CAPLETS**
 (**capsule-shaped tablets)



ACTUAL SIZE

MIGRAINE RELIEF

acetaminophen, aspirin, caffeine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0374
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	L374
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0374-78	1 in 1 CARTON	05/02/2002	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0363-0374-85	250 in 1 BOTTLE; Type 0: Not a Combination Product	05/02/2002	01/08/2017
3	NDC:0363-0374-62	1 in 1 CARTON	05/02/2002	
3		24 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0363-0374-82	1 in 1 CARTON	05/02/2002	
4		200 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075794	05/02/2002	

Labeler - Walgreen Company (008965063)

Revised: 3/2023

Walgreen Company