

ASPIRIN ENTERIC COATED, REGULAR STRENGTH- aspirin tablet, delayed release

Walgreen Company

Aspirin

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - a cold
 - headache
 - menstrual pain
 - toothache
 - muscle pain
- or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours not to

exceed 12 tablets in 24 hours unless directed by a doctor

- children under 12 years: do not use unless directed by a doctor

Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

black iron oxide, cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, shellac wax, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

VALUE SIZE

NDC 0363-0227-14

Walgreens

Compare to the active ingredient
in Ecotrin® Regular Strength^{††}

WALGREENS PHARMACIST RECOMMENDED[†]

Aspirin 325

325 mg / PAIN RELIEVER (NSAID)

Regular Strength

Enteric-Coated

500

TABLETS

ACTUAL SIZE

HEART

HEALTH**

**Talk to your doctor or other healthcare
provider before using this product for your heart.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

[†]Our pharmacists recommend the Walgreens brand. We invite you to

compare to national brands.

††This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Ecotrin® Regular Strength.

50844 ORG01222714

DISTRIBUTED BY: **WALGREEN CO.**

DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com ©2021 Walgreen Co.

Drug Facts (continued)

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug ■ are age 60 or older
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

††This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Ecotrin® Regular Strength.

50844 ORG01222714 ITEM 907107 W00000-0000-0

DISTRIBUTED BY: WALGREEN CO.
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.

3 11917 17758 8

PEEL HERE FOR MORE DRUG FACTS

REV-0723 W3ORG1221-F

VALUE SIZE NDC 0363-0227-14

Walgreens

Compare to the active ingredient in Ecotrin® Regular Strength†

Aspirin 325

325 mg / PAIN RELIEVER (NSAID)

Regular Strength Enteric-Coated

500 TABLETS

HEART HEALTH

ACTUAL SIZE

**Talk to your doctor or other healthcare provider before using this product for your heart.

Drug Facts **TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Active ingredient (in each tablet) **Purpose**

Aspirin 325 mg (NSAID)*..... Pain reliever

*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:

- minor pain of arthritis ■ a cold
- headache ■ menstrual pain
- toothache ■ muscle pain

■ or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Drug Facts (continued)

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur

These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information ■ use by expiration date on package

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients black iron oxide, cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, shellac wax, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-426-9391

Walgreens 44-227

ASPIRIN ENTERIC COATED, REGULAR STRENGTH

aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0227
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
Inactive Ingredients		
Ingredient Name		Strength
FERROSOFERRIC OXIDE (UNII: XM0M87F357)		
POWDERED CELLULOSE (UNII: SMD1X3XO9M)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
STARCH, CORN (UNII: O8232NY3SJ)		
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
POLYDEXTROSE (UNII: VH2XOU12IE)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
SHELLAC (UNII: 46N107B71O)		
DIMETHICONE (UNII: 92RU3N3Y1O)		
WATER (UNII: 059QF0KO0R)		
SODIUM ALGINATE (UNII: C269C4G2ZQ)		
SODIUM BICARBONATE (UNII: 8MDF5V39QO)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
TALC (UNII: 7SEV7J4R1U)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		
TRIACETIN (UNII: XHX3C3X673)		
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)		

Product Characteristics			
Color	orange	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;227
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0227-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/14/1992	
2	NDC:0363-0227-57	1 in 1 CARTON	12/14/1992	07/20/2020
2		125 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/14/1992	

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	manufacture(0363-0227) , pack(0363-0227)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-0227)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-0227)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(0363-0227)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0363-0227)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-0227)

Revised: 8/2023

Walgreen Company