

**REGULAR STRENGTH ANTACID- aluminum hydroxide, magnesium hydroxide,  
dimethicone liquid  
AMERISOURCEBERGEN DRUG CORPORATION**

-----  
**GNP antacid original**

**Active ingredients (in each 10 mL dose)**

Aluminum hydroxide 400 mg (equivalent to dried gel, USP)  
Magnesium hydroxide 400 mg  
Simethicone 40mg

**Purposes**

Antacid  
Antigas

**Uses**

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

**Warnings**

**Do not take** more than 120 mL (12 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks.

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Antacids may interact with certain prescription drugs.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Directions**

- shake well before use
- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
- do not take more than 24 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

**Other information**

- **each 10 mL dose contains:** magnesium 165 mg, sodium 6 mg
- store at room temperature
- protect from freezing
- keep tightly closed

**Inactive ingredients**

benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

**Questions or comments?**

**1-800-540-3765**

**package Label**



Compare to Mylanta®  
Regular Strength  
active ingredients\*

NDC 46122-433-40

▶ **REGULAR STRENGTH**

# Antacid Liquid

**Alumina, Magnesia, and Simethicone Oral Suspension**

Fast Relief of:

- Acid Indigestion
- Heartburn
- Sour Stomach
- Pressure and Bloating

Original Flavor

Alcohol: 0.15%

**12 FL OZ (355 mL)**

297-05112-0 REV0917

## Drug Facts

**TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.**

### Active ingredients

(in each 10 mL dose)

Active ingredients	Purpose
Aluminum hydroxide 400 mg (equivalent to dried gel, USP)	Antacid
Magnesium hydroxide 400 mg	Antacid
Simethicone 40 mg	Antigas

### Uses

- relieves • heartburn • sour stomach
- acid indigestion • symptoms referred to as gas

### Warnings

Do not take more than 120 mL (12 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks.

Ask a doctor before use if you have

- kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

### Directions

- shake well before use
- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses) between meals, at bedtime or as directed by a doctor
- children under 12 years of age: ask a doctor.

### Other information

- each 10 mL dose contains: magnesium 165 mg, sodium 6 mg
- store at room temperature
- protect from freezing • keep tightly closed

### Inactive ingredients

benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

\*This product is not manufactured or distributed by Johnson & Johnson Merck, distributor of Mylanta®.



Distributed By  
AmerisourceBergen  
1 West First Avenue  
Conshohocken, PA 19428  
Questions or Concerns?  
[www.mygrp.com](http://www.mygrp.com)



## REGULAR STRENGTH ANTACID

aluminum hydroxide, magnesium hydroxide, dimethicone liquid

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-433
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength

<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 10 mL
<b>MAGNESIUM HYDROXIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 10 mL
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 10 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>BUTYLPARABEN</b> (UNII: 3QPI1U3FV8)	
<b>HYDROXYETHYL CELLULOSE, UNSPECIFIED</b> (UNII: T4V6TWG28D)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	LEMON (lemon)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-433-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/31/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	01/01/2000	

**Labeler** - AMERISOURCEBERGEN DRUG CORPORATION (007914906)

**Registrant** - GCP Laboratories (965480861)

### Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(46122-433)