

HEADACHE RELIEF EXTRA STRENGTH- acetaminophen, aspirin, caffeine tablet Safeway, Inc.

Drug Facts

Active ingredients (in each geltab)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever

Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - a cold
 - arthritis
 - muscular aches
 - toothache
 - premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause severe allergic reaction, which may include:

- hives

- facial swelling
- shock
- asthma (wheezing)

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop use and ask doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
 - feel faint
 - vomit blood

- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 geltabs every 6 hours; do not take more than 8 geltabs in 24 hours
- children under 12 years: do not use unless directed by a doctor

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients

black printing ink, corn starch, D&C yellow #10 aluminum lake, edetate disodium, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hydroxypropyl cellulose, hydroxypropyl cellulose-low substituted, hypromellose, iron oxide black, maltitol, povidone, purified water, stearic acid, talc, titanium dioxide

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

COMPARE TO Excedrin® Extra Strength active ingredients†

Extra Strength

Headache Relief

ACETAMINOPHEN 250 mg

ASPIRIN (**NSAID**) 250 mg

CAFFEINE 65 mg

Pain Reliever/Pain Reliever Aid

GELTABS

†This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare Holdings (US) LLC, distributor of Excedrin® Extra Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY

BETTER LIVING BRANDS LLC

P.O. BOX 99

PLEASANTON, CA 94566-0009

Package Label green

This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare Holdings (US) LLC, distributor of Excedrin® Extra Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



how2recycle.info



Scan here for more information

Drug Facts (continued)

hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 2 tablets every 6 hours; do not take more than 8 tablets in 24 hours ■ children under 12 years: do not use unless directed by a doctor

Other information
■ store between 20°-25°C (68°-77°F)

Inactive ingredients black printing ink, corn starch, D&C yellow #10 aluminum lake, croscarellin, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hydroxypropyl cellulose, hydroxypropyl cellulose-low substituted, hypromellose, iron oxide black, mallicol, povidone, purified water, stearic acid, talc, titanium dioxide

Questions or comments?
Call 1-877-753-3955 Monday-Friday 9AM-5PM EST



COMPARE TO
Excedrin® Extra Strength
active ingredients*

NDC 21130-586-20

Extra Strength Headache Relief
ACETAMINOPHEN 250 mg
ASPIRIN (NSAID) 250 mg
CAFFEINE 65 mg
Pain Reliever/Pain Reliever Aid

Actual Size



20 GELTABS

DISTRIBUTED BY
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0099

OUR PROMISE
QUALITY & SATISFACTION
100% GUARANTEED
ON YOUR MONEY BACK.



RD 24030
3 21130 18125 5

PLU-8514C F000877Z
Lot No.: Exp. Date:

Drug Facts

Active ingredients
(in each geltab)

Purposes

Acetaminophen 250 mg.....Pain reliever
Aspirin 250 mg (NSAID)*.....Pain reliever
Caffeine 65 mg.....Pain reliever and nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to
■ headache ■ a cold ■ arthritis ■ muscular aches
■ toothache ■ premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:
■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older

Drug Facts (continued)

■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use ■ if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are
■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away; ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of

SIGNATURE CARE Extra Strength Headache Relief

Package label

This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare Holdings (US) LLC, distributor of Excedrin® Extra Strength.

DANGER: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



Scan here for more information



Drug Facts (continued)

heating occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 2 tablets every 6 hours; do not take more than 8 tablets in 24 hours ■ children under 12 years: do not use unless directed by a doctor

Other information
■ store between 20°-25°C (68°-77°F)

Inactive ingredients black printing ink, corn starch, D&C yellow #10 aluminum lake, edelate disodium, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hydroxypropyl cellulose, hydroxypropyl cellulose-low substituted, hypromellose, iron oxide black, methylol, polydora, purified water, stearic acid, talc, titanium dioxide

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST



COMPARE TO Excedrin® Extra Strength
active ingredients*

NDC 21130-586-20

Extra Strength Headache Relief
ACETAMINOPHEN 250 mg
ASPIRIN (NSAID) 250 mg
CAFFEINE 65 mg
Pain Reliever/Pain Reliever Aid



Actual Size

20 GELTABS

OUR PROMISE
QUALITY ASSURANCE
100% GUARANTEED
OR YOUR MONEY BACK

DISTRIBUTED BY
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0099
1-888-723-3929

RD 204030
3 21130118126 2
PUD-8514C F0008573
Lot No.: Exp. Date:

Drug Facts

Active Ingredients
(in each tablet)

Purposes

Acetaminophen 250 mg.....Pain reliever
Aspirin 250 mg (NSAID)*.....Pain reliever
Caffeine 65 mg.....Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to ■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache ■ premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older

Drug Facts (continued)

■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use ■ if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic disease ■ you have asthma ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away: ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ fringing in the ears or loss of

SIGNATURE CARE Extra Strength Headache Relief

HEADACHE RELIEF EXTRA STRENGTH

acetaminophen, aspirin, caffeine tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:21130-586

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MALTITOL (UNII: D65DG142WK)	
WATER (UNII: 059QF0KO0R)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
HYDROXYPROPYL CELLULOSE, LOW SUBSTITUTED (UNII: 2165RE0K14)	

Product Characteristics

Color	white (green/white)	Score	no score
Shape	ROUND	Size	13mm
Flavor		Imprint Code	P
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-586-20	1 in 1 CARTON	04/18/2022	
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
OTC Monograph Drug	M013	04/18/2022	

Labeler - Safeway, Inc. (009137209)

Revised: 2/2024

Safeway, Inc.