

**AMPICILLIN AND SULBACTAM- ampicillin sodium and sulbactam sodium injection, powder, for solution**  
NorthStar Rx, LLC

-----  
**Ampicillin and Sulbactam for Injection, USP**  
**(For Intramuscular or Intravenous Use)**

---

**PHARMACY BULK PACKAGE - NOT FOR DIRECT INFUSION**

---


**Rx Only**

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Ampicillin and Sulbactam for Injection, USP and other antibacterial drugs, ampicillin and sulbactam for injection should be used only to treat infections that are proven or strongly suspected to be caused by bacteria.


**DESCRIPTION**

Ampicillin and Sulbactam for Injection, USP is an injectable antibacterial combination consisting of the semisynthetic antibacterial ampicillin sodium and the beta-lactamase inhibitor sulbactam sodium for intravenous and intramuscular administration. **THE INTENT OF THIS PHARMACY BULK PACKAGE IS FOR PREPARATION OF SOLUTIONS FOR IV INFUSION ONLY.**

Ampicillin sodium is derived from the penicillin nucleus, 6-aminopenicillanic acid. Chemically, it is monosodium (2S, 5R, 6R)-6-[(R)-2-amino-2-phenylacetamido]-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo[3.2.0]heptane-2-carboxylate and has a molecular weight of 371.39. Its chemical formula is  $C_{16}H_{18}N_3NaO_4S$ . The structural formula is:



Sulbactam sodium is a derivative of the basic penicillin nucleus. Chemically, sulbactam sodium is sodium penicillinate sulfone; sodium (2S, 5R)-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo [3.2.0] heptane-2-carboxylate 4,4-dioxide. Its chemical formula is  $C_8H_{10}NNaO_5S$  with a molecular weight of 255.22. The structural formula is:



Ampicillin and Sulbactam for Injection, USP, ampicillin sodium/sulbactam sodium parenteral combination, is available as a white to off-white dry powder for reconstitution. Ampicillin and Sulbactam for Injection, USP dry powder is freely soluble in aqueous diluents to yield pale yellow to yellow solutions containing ampicillin sodium and sulbactam sodium equivalent to 250 mg ampicillin per mL and 125 mg sulbactam per mL. The pH of the solutions is between 8.0 and 10.0.

Dilute solutions (up to 30 mg ampicillin and 15 mg sulbactam per mL) are essentially colorless to pale yellow. The pH of dilute solutions remains the same.

Each sterile Pharmacy Bulk Package contains 15 grams Ampicillin and Sulbactam for Injection (equivalent to 10 g ampicillin as the sodium salt plus 5 g sulbactam as the sodium salt). The sodium content per vial is 1150 mg (50 mEq) sodium. Ampicillin and Sulbactam for Injection Pharmacy Bulk Package is a vial containing a sterile preparation of ampicillin sodium and sulbactam sodium for parenteral use that contains many single doses. The Pharmacy Bulk Package is for use in a pharmacy admixture setting; it provides many single doses of ampicillin and sulbactam for injection for addition to suitable parenteral fluids in the preparation and administration for intravenous infusion (see **DIRECTIONS FOR PROPER USE OF PHARMACY BULK PACKAGE** section). **FURTHER DILUTION IS REQUIRED BEFORE USE.**

## CLINICAL PHARMACOLOGY

### General

Immediately after completion of a 15-minute intravenous infusion of ampicillin and sulbactam for injection, peak serum concentrations of ampicillin and sulbactam are attained. Ampicillin serum levels are similar to those produced by the administration of equivalent amounts of ampicillin alone. Peak ampicillin serum levels ranging from 109 to 150 mcg/mL are attained after administration of 2000 mg of ampicillin plus 1000 mg sulbactam and 40 to 71 mcg/mL after administration of 1000 mg ampicillin plus 500 mg sulbactam. The corresponding mean peak serum levels for sulbactam range from 48 to 88 mcg/mL and 21 to 40 mcg/mL, respectively. After an intramuscular injection of 1000 mg ampicillin plus 500 mg sulbactam, peak ampicillin serum levels ranging from 8 to 37 mcg/mL and peak sulbactam serum levels ranging from 6 to 24 mcg/mL are attained.

The mean serum half-life of both drugs is approximately 1 hour in healthy volunteers.

Approximately 75 to 85% of both ampicillin and sulbactam are excreted unchanged in the urine during the first 8 hours after administration of ampicillin and sulbactam for injection to individuals with normal renal function. Somewhat higher and more prolonged serum levels of ampicillin and sulbactam can be achieved with the concurrent

administration of probenecid.

In patients with impaired renal function the elimination kinetics of ampicillin and sulbactam are similarly affected, hence the ratio of one to the other will remain constant whatever the renal function. The dose of ampicillin and sulbactam for injection in such patients should be administered less frequently in accordance with the usual practice for ampicillin (see **DOSAGE AND ADMINISTRATION** section).

Ampicillin has been found to be approximately 28% reversibly bound to human serum protein and sulbactam approximately 38% reversibly bound.

The following average levels of ampicillin and sulbactam were measured in the tissues and fluids listed:

**TABLE 1**

| <b>Fluid or Tissue</b>      | <b>Dose<br/>(grams)<br/>Ampicillin/Sulbactam</b> | <b>Concentration<br/>(mcg/mL or mcg/g)<br/>Ampicillin/Sulbactam</b> |
|-----------------------------|--------------------------------------------------|---------------------------------------------------------------------|
| Peritoneal Fluid            | 0.5/0.5 IV                                       | 7/14                                                                |
| Blister Fluid (Cantharides) | 0.5/0.5 IV                                       | 8/20                                                                |
| Tissue Fluid                | 1/0.5 IV                                         | 8/4                                                                 |
| Intestinal Mucosa           | 0.5/0.5 IV                                       | 11/18                                                               |
| Appendix                    | 2/1 IV                                           | 3/40                                                                |

Penetration of both ampicillin and sulbactam into cerebrospinal fluid in the presence of inflamed meninges has been demonstrated after IV administration of ampicillin and sulbactam for injection.

The pharmacokinetics of ampicillin and sulbactam in pediatric patients receiving ampicillin and sulbactam for injection are similar to those observed in adults. Immediately after a 15-minute infusion of 50 to 75 mg ampicillin and sulbactam for injection /kg body weight, peak serum and plasma concentrations of 82 to 446 mcg ampicillin/mL and 44 to 203 mcg sulbactam/mL were obtained. Mean half-life values were approximately 1 hour.

## **MICROBIOLOGY**

Ampicillin is similar to benzyl penicillin in its bactericidal action against susceptible organisms during the stage of active multiplication. It acts through the inhibition of cell wall mucopeptide biosynthesis. Ampicillin has a broad spectrum of bactericidal activity against many gram-positive and gram-negative aerobic and anaerobic bacteria. (Ampicillin is, however, degraded by beta-lactamases and therefore the spectrum of activity does not normally include organisms which produce these enzymes).

A wide range of beta-lactamases found in microorganisms resistant to penicillins and cephalosporins have been shown in biochemical studies with cell free bacterial systems to be irreversibly inhibited by sulbactam. Although sulbactam alone possesses little useful antibacterial activity except against the Neisseriaceae, whole organism studies have shown that sulbactam restores ampicillin activity against beta-lactamase producing

strains. In particular, sulbactam has good inhibitory activity against the clinically important plasmid mediated beta-lactamases most frequently responsible for transferred drug resistance. Sulbactam has no effect on the activity of ampicillin against ampicillin susceptible strains.

The presence of sulbactam in the ampicillin and sulbactam for injection formulation effectively extends the antibacterial spectrum of ampicillin to include many bacteria normally resistant to it and to other beta-lactam antibacterials. Thus, ampicillin and sulbactam for injection possesses the properties of a broad-spectrum antibacterial and a beta-lactamase inhibitor.

While *in vitro* studies have demonstrated the susceptibility of most strains of the following organisms, clinical efficacy for infections other than those included in the **INDICATIONS and USAGE** section has not been documented.

**Gram-Positive Bacteria:** *Staphylococcus aureus* (beta-lactamase and non-beta-lactamase producing), *Staphylococcus epidermidis* (beta-lactamase and non-beta-lactamase producing), *Staphylococcus saprophyticus* (beta-lactamase and non-beta-lactamase producing), *Streptococcus faecalis*† (Enterococcus), *Streptococcus pneumoniae*† (formerly *D. pneumoniae*), *Streptococcus pyogenes*†, *Streptococcus viridans*†.

**Gram-Negative Bacteria:** *Hemophilus influenzae* (beta-lactamase and non-beta-lactamase producing), *Moraxella* (*Branhamella*) *catarrhalis* (beta-lactamase and non-beta-lactamase producing), *Escherichia coli* (beta-lactamase and non-beta-lactamase producing), *Klebsiella* species (all known strains are beta-lactamase producing), *Proteus mirabilis* (beta-lactamase and non-beta-lactamase producing), *Proteus vulgaris*, *Providencia rettgeri*, *Providencia stuartii*, *Morganella morganii*, and *Neisseria gonorrhoeae* (beta-lactamase and non-beta-lactamase producing).

**Anaerobes:** *Clostridium* species,† *Peptococcus* species,† *Peptostreptococcus* species, *Bacteroides* species, including *B. fragilis*.

† These are not beta-lactamase producing strains and, therefore, are susceptible to ampicillin alone.

## Susceptibility Testing

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

## INDICATIONS AND USAGE

Ampicillin and sulbactam for injection is indicated for the treatment of infections due to susceptible strains of the designated microorganisms in the conditions listed below.

**Skin and Skin Structure Infections** caused by beta-lactamase producing strains of *Staphylococcus aureus*, *Escherichia coli*,\* *Klebsiella* spp.\* (including *K. pneumoniae*\*), *Proteus mirabilis*,\* *Bacteroides fragilis*,\* *Enterobacter* spp.,\* and *Acinetobacter calcoaceticus*.\*

NOTE: For information on use in pediatric patients (see **PRECAUTIONS - Pediatric Use** and **CLINICAL STUDIES** sections).

**Intra-Abdominal Infections** caused by beta-lactamase producing strains of *Escherichia coli*, *Klebsiella* spp. (including *K. pneumoniae*\*), *Bacteroides* spp. (including *B. fragilis*), and *Enterobacter* spp.\*

**Gynecological Infections** caused by beta-lactamase producing strains of *Escherichia coli*,\* and *Bacteroides* spp.\* (including *B. fragilis*\*).

\* Efficacy for this organism in this organ system was studied in fewer than 10 infections.

While ampicillin and sulbactam for injection is indicated only for the conditions listed above, infections caused by ampicillin-susceptible organisms are also amenable to treatment with ampicillin and sulbactam for injection due to its ampicillin content. Therefore, mixed infections caused by ampicillin-susceptible organisms and beta-lactamase producing organisms susceptible to ampicillin and sulbactam for injection should not require the addition of another antibacterial.

Appropriate culture and susceptibility tests should be performed before treatment in order to isolate and identify the organisms causing infection and to determine their susceptibility to ampicillin and sulbactam for injection.

Therapy may be instituted prior to obtaining the results from bacteriological and susceptibility studies when there is reason to believe the infection may involve any of the beta-lactamase producing organisms listed above in the indicated organ systems. Once the results are known, therapy should be adjusted if appropriate.

To reduce the development of drug-resistant bacteria and maintain effectiveness of ampicillin and sulbactam for injection and other antibacterial drugs, ampicillin and sulbactam for injection should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

## **CONTRAINDICATIONS**

The use of ampicillin and sulbactam for injection is contraindicated in individuals with a history of serious hypersensitivity reactions (e.g., anaphylaxis or Stevens-Johnson syndrome) to ampicillin, sulbactam or to other beta-lactam antibacterial drugs (e.g., penicillins and cephalosporins).

Ampicillin and sulbactam for injection is contraindicated in patients with a previous history of cholestatic jaundice/hepatic dysfunction associated with ampicillin and sulbactam for injection.

## **WARNINGS**

### **Hypersensitivity**

Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients on penicillin therapy. These reactions are more apt to occur in individuals with a history of penicillin hypersensitivity and/or hypersensitivity reactions to

multiple allergens. There have been reports of individuals with a history of penicillin hypersensitivity who have experienced severe reactions when treated with cephalosporins. Before therapy with a penicillin, careful inquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins, and other allergens. If an allergic reaction occurs, ampicillin and sulbactam for injection should be discontinued and the appropriate therapy instituted.

### **Hepatotoxicity**

Hepatic dysfunction, including hepatitis and cholestatic jaundice has been associated with the use of ampicillin and sulbactam for injection. Hepatic toxicity is usually reversible; however, deaths have been reported. Hepatic function should be monitored at regular intervals in patients with hepatic impairment.

### **Severe Cutaneous Adverse Reactions**

Ampicillin and sulbactam for injection may cause severe skin reactions, such as toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome (SJS), dermatitis exfoliative, erythema multiforme, and Acute generalized exanthematous pustulosis (AGEP). If patients develop a skin rash they should be monitored closely and ampicillin and sulbactam for injection discontinued if lesions progress (see **CONTRAINDICATIONS** and **ADVERSE REACTIONS** sections).

### ***Clostridium difficile*-Associated Diarrhea**

*Clostridium difficile* associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including ampicillin and sulbactam for injection, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

*C. difficile* produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibacterial drug use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibacterial drug use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibacterial treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

## **PRECAUTIONS**

### **General**

A high percentage of patients with mononucleosis who receive ampicillin develop a skin rash. Thus, ampicillin class antibacterial should not be administered to patients with mononucleosis. In patients treated with ampicillin and sulbactam for injection the possibility of superinfections with mycotic or bacterial pathogens should be kept in mind during therapy. If superinfections occur (usually involving *Pseudomonas* or *Candida*), the drug should be discontinued and/or appropriate therapy instituted.

Prescribing ampicillin and sulbactam for injection in the absence of proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

### **Information for Patients**

Patients should be counseled that antibacterial drugs including ampicillin and sulbactam for injection should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When ampicillin and sulbactam for injection is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by ampicillin and sulbactam for injection or other antibacterial drugs in the future.

Diarrhea is a common problem caused by antibacterial which usually ends when the antibacterial is discontinued. Sometimes after starting treatment with antibacterial, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibacterial. If this occurs, patients should contact their physician as soon as possible.

### **Drug Interactions**

Probenecid decreases the renal tubular secretion of ampicillin and sulbactam. Concurrent use of probenecid with ampicillin and sulbactam for injection may result in increased and prolonged blood levels of ampicillin and sulbactam. The concurrent administration of allopurinol and ampicillin increases substantially the incidence of rashes in patients receiving both drugs as compared to patients receiving ampicillin alone. It is not known whether this potentiation of ampicillin rashes is due to allopurinol or the hyperuricemia present in these patients. There are no data with ampicillin and sulbactam for injection and allopurinol administered concurrently. Ampicillin and sulbactam for injection and aminoglycosides should not be reconstituted together due to the *in vitro* inactivation of aminoglycosides by the ampicillin component of ampicillin and sulbactam for injection.

### **Drug/Laboratory Test Interactions**

Administration of ampicillin and sulbactam for injection will result in high urine concentration of ampicillin. High urine concentrations of ampicillin may result in false positive reactions when testing for the presence of glucose in urine using Clinitest™, Benedict's Solution or Fehling's Solution. It is recommended that glucose tests based on enzymatic glucose oxidase reactions (such as Clinistix™ or Testape™) be used. Following administration of ampicillin to pregnant women, a transient decrease in plasma concentration of total conjugated estriol, estriol-glucuronide, conjugated estrone and estradiol has been noted. This effect may also occur with ampicillin and sulbactam for injection.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term studies in animals have not been performed to evaluate carcinogenic or mutagenic potential.

## **Pregnancy**

Reproduction studies have been performed in mice, rats, and rabbits at doses up to ten (10) times the human dose and have revealed no evidence of impaired fertility or harm to the fetus due to ampicillin and sulbactam for injection. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed. (see **PRECAUTIONS-Drug/Laboratory Test Interactions** section).

## **Labor and Delivery**

Studies in guinea pigs have shown that intravenous administration of ampicillin decreased the uterine tone, frequency of contractions, height of contractions, and duration of contractions. However, it is not known whether the use of ampicillin and sulbactam for injection in humans during labor or delivery has immediate or delayed adverse effects on the fetus, prolongs the duration of labor, or increases the likelihood that forceps delivery or other obstetrical intervention or resuscitation of the newborn will be necessary.

## **Nursing Mothers**

Low concentrations of ampicillin and sulbactam are excreted in the milk; therefore, caution should be exercised when ampicillin and sulbactam for injection is administered to a nursing woman.

## **Pediatric Use**

The safety and effectiveness of ampicillin and sulbactam for injection have been established for pediatric patients one year of age and older for skin and skin structure infections as approved in adults. Use of ampicillin and sulbactam for injection in pediatric patients is supported by evidence from adequate and well-controlled studies in adults with additional data from pediatric pharmacokinetic studies, a controlled clinical trial conducted in pediatric patients and post-marketing adverse events surveillance. (see **CLINICAL PHARMACOLOGY, INDICATIONS AND USAGE, ADVERSE REACTIONS, DOSAGE AND ADMINISTRATION**, and **CLINICAL STUDIES** sections).

The safety and effectiveness of ampicillin and sulbactam for injection have not been established for pediatric patients for intra-abdominal infections.

## **ADVERSE REACTIONS**

**Adult Patients:** Ampicillin and sulbactam for injection is generally well tolerated. The following adverse reactions have been reported in clinical trials.

### **Local Adverse Reactions**

Pain at IM injection site – 16%

Pain at IV injection site – 3% Thrombophlebitis – 3% Phlebitis – 1.2%

### **Systemic Adverse Reactions**

The most frequently reported adverse reactions were diarrhea in 3% of the patients and



rash in less than 2% of the patients.

Additional systemic reactions reported in less than 1% of the patients were: itching, nausea, vomiting, candidiasis, fatigue, malaise, headache, chest pain, flatulence, abdominal distension, glossitis, urine retention, dysuria, edema, facial swelling, erythema, chills, tightness in throat, substernal pain, epistaxis and mucosal bleeding.

**Pediatric Patients:** Available safety data for pediatric patients treated with ampicillin and sulbactam for injection demonstrate a similar adverse events profile to those observed in adult patients. Additionally, atypical lymphocytosis has been observed in one pediatric patient receiving ampicillin and sulbactam for injection.

### **Adverse Laboratory Changes**

Adverse laboratory changes without regard to drug relationship that were reported during clinical trials were:

*Hepatic:* Increased AST (SGOT), ALT (SGPT), alkaline phosphatase, and LDH.

*Hematologic:* Decreased hemoglobin, hematocrit, RBC, WBC, neutrophils, lymphocytes, platelets and increased lymphocytes, monocytes, basophils, eosinophils, and platelets.

*Blood Chemistry:* Decreased serum albumin and total proteins.

*Renal:* Increased BUN and creatinine.

*Urinalysis:* Presence of RBC's and hyaline casts in urine.

### **Postmarketing Experience**

In addition to adverse reactions reported from clinical trials, the following have been identified during post-marketing use of ampicillin sodium/sulbactam sodium or other products containing ampicillin. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. These events have been chosen for inclusion due to a combination of their seriousness, frequency, or potential causal connection to ampicillin sodium/sulbactam sodium.

**Blood and Lymphatic System Disorders:** Hemolytic anemia, thrombocytopenic purpura, and agranulocytosis have been reported. These reactions are usually reversible on discontinuation of therapy and are believed to be hypersensitivity phenomena. Some individuals have developed positive direct Coombs Tests during treatment with ampicillin and sulbactam for injection, as with other beta-lactam antibacterials.

**Gastrointestinal Disorders:** Abdominal pain, cholestatic hepatitis, cholestasis, hyperbilirubinemia, jaundice, abnormal hepatic function, melena, gastritis, stomatitis, dyspepsia, black "hairy" tongue, and *Clostridium difficile* associated diarrhea (see **CONTRAINDICATIONS** and **WARNINGS** sections).

**General Disorders and Administration Site Conditions:** Injection site reaction

**Immune System Disorders:** Serious and fatal hypersensitivity (anaphylactic) reactions (see **WARNINGS** section)

**Nervous System Disorders:** Convulsion and dizziness

**Renal and Urinary Disorders:** Tubulointerstitial nephritis

**Respiratory, Thoracic and Mediastinal Disorders:** Dyspnea

**Skin and Subcutaneous Tissue Disorders:** Toxic epidermal necrolysis, Stevens-Johnson syndrome, angioedema, Acute generalized exanthematous pustulosis (AGEP), erythema multiforme, exfoliative dermatitis, and urticaria (see **CONTRAINDICATIONS** and **WARNINGS** sections), and linear IgA bullous dermatosis.

## OVERDOSAGE

Neurological adverse reactions, including convulsions, may occur with the attainment of high CSF levels of beta-lactams. Ampicillin may be removed from circulation by hemodialysis. The molecular weight, degree of protein binding and pharmacokinetics profile of sulbactam suggest that this compound may also be removed by hemodialysis.

## CLINICAL STUDIES

### Skin and Skin Structure Infections in Pediatric Patients

Data from a controlled clinical trial conducted in pediatric patients provided evidence supporting the safety and efficacy of ampicillin and sulbactam for injection for the treatment of skin and skin structure infections. Of 99 pediatric patients evaluable for clinical efficacy, 60 patients received a regimen containing intravenous ampicillin and sulbactam for injection, and 39 patients received a regimen containing intravenous cefuroxime. This trial demonstrated similar outcomes (assessed at an appropriate interval after discontinuation of all antimicrobial therapy) for ampicillin and sulbactam for injection- and cefuroxime-treated patients:

**TABLE 2**

| <b><u>Therapeutic Regimen</u></b>      | <b><u>Clinical Success</u></b> | <b><u>Clinical Failure</u></b> |
|----------------------------------------|--------------------------------|--------------------------------|
| Ampicillin and Sulbactam for Injection | 51/60 (85%)                    | 9/60 (15%)                     |
| Cefuroxime                             | 34/39 (87%)                    | 5/39 (13%)                     |

Most patients received a course of oral antimicrobials following initial treatment with intravenous administration of parenteral antimicrobials. The study protocol required that the following three criteria be met prior to transition from intravenous to oral antimicrobial therapy: (1) receipt of a minimum of 72 hours of intravenous therapy; (2) no documented fever for prior 24 hours; and (3) improvement or resolution of the signs and symptoms of infection.

The choice of oral antimicrobial agent used in this trial was determined by susceptibility testing of the original pathogen, if isolated, to oral agents available. The course of oral antimicrobial therapy should not routinely exceed 14 days.

## DOSAGE AND ADMINISTRATION

Ampicillin and sulbactam for injection may be used for parenteral administration (**following dilution**). **THE INTENT OF THIS PHARMACY BULK PACKAGE IS FOR PREPARATION OF SOLUTIONS FOR IV INFUSION ONLY.**

For IV administration, the dose can be given by slow intravenous injection over at least 10–15 minutes or can also be delivered in greater dilutions with 50–100 mL of a compatible diluent as an intravenous infusion over 15–30 minutes.

The recommended adult dosage of ampicillin and sulbactam for injection is 1.5 g (1 g ampicillin as the sodium salt plus 0.5 g sulbactam as the sodium salt) to 3 g (2 g ampicillin as the sodium salt plus 1 g sulbactam as the sodium salt) every six hours. This 1.5 to 3 g range represents the total of ampicillin content plus the sulbactam content of ampicillin and sulbactam for injection, and corresponds to a range of 1 g ampicillin/0.5 g sulbactam to 2 g ampicillin/1 g sulbactam. The total dose of sulbactam should not exceed 4 grams per day.

### **Pediatric Patients 1 Year of Age or Older**

The recommended daily dose of ampicillin and sulbactam for injection in pediatric patients is 300 mg per kg of body weight administered via intravenous infusion in equally divided doses every 6 hours. This 300 mg/kg/day dosage represents the total ampicillin content plus the sulbactam content of ampicillin and sulbactam for injection, and corresponds to 200 mg ampicillin/100 mg sulbactam per kg per day. The safety and efficacy of ampicillin and sulbactam for injection administered via intramuscular injection in pediatric patients have not been established. Pediatric patients weighing 40 kg or more should be dosed according to adult recommendations, and the total dose of sulbactam should not exceed 4 grams per day. The course of intravenous therapy should not routinely exceed 14 days. In clinical trials, most children received a course of oral antimicrobials following initial treatment with intravenous ampicillin and sulbactam for injection. (see **CLINICAL STUDIES** section).

### **Impaired Renal Function**

In patients with impairment of renal function the elimination kinetics of ampicillin and sulbactam are similarly affected, hence the ratio of one to the other will remain constant whatever the renal function. The dose of ampicillin and sulbactam for injection in such patients should be administered less frequently in accordance with the usual practice for ampicillin and according to the following recommendations:

**TABLE 3**

**Ampicillin and Sulbactam for Injection, USP  
Dosage Guide for Patients with Renal Impairment**

| <b>Creatinine Clearance<br/>(mL/min/1.73m<sup>2</sup>)</b> | <b>Ampicillin/Sulbactam<br/>Half-Life (Hours)</b> | <b>Recommended<br/>Ampicillin and Sulbactam<br/>for Injection, USP Dosage</b> |
|------------------------------------------------------------|---------------------------------------------------|-------------------------------------------------------------------------------|
| ≥30                                                        | 1                                                 | 1.5 g - 3 g q 6h - q 8h                                                       |
| 15 - 29                                                    | 5                                                 | 1.5 g - 3 g q 12h                                                             |
| 5 - 14                                                     | 9                                                 | 1.5 g - 3 g q 24h                                                             |

When only serum creatinine is available, the following formula (based on sex, weight, and age of the patient) may be used to convert this value into creatinine clearance. The serum creatinine should represent a steady state of renal function.

Males  $\frac{\text{weight (kg)} \times (140 - \text{age})}{72 \times \text{serum creatinine}}$

Females 0.85 x above value

## **COMPATIBILITY, RECONSTITUTION AND STABILITY**

Ampicillin and sulbactam for injection sterile powder is to be stored at or below 30°C (86°F) prior to reconstitution.

When concomitant therapy with aminoglycosides is indicated, ampicillin and sulbactam for injection and aminoglycosides should be reconstituted and administered separately, due to the in vitro inactivation of aminoglycosides by any of the aminopenicillins.

## **DIRECTIONS FOR USE**

### **Intravenous Administration**

#### Directions for Proper Use of Pharmacy Bulk Package

Ampicillin and sulbactam for injection sterile powder for intravenous use may be reconstituted with any of the compatible diluents described in this insert. Solutions should be allowed to stand after dissolution to allow any foaming to dissipate in order to permit visual inspection for complete solubilization.

Ampicillin and Sulbactam for Injection, USP concentrations between 3 and 45 mg (2 to 30 mg ampicillin/1 to 15 mg sulbactam/mL) are recommended for intravenous use.

The 15 gram may be reconstituted with either 92 mL Sterile Water for Injection or 0.9% Sodium Chloride Injection. The diluent should be added in two separate aliquots in a suitable work area, such as a laminar flow hood. Add 50 mL of solution, shake to dissolve. Then add an additional 42 mL and shake. The solution should be allowed to stand after dissolution to allow any foaming to dissipate in order to permit visual inspection for complete solubilization. The resultant solution should have a final concentration of approximately 100 mg/mL ampicillin and 50 mg/mL sulbactam. The closure may be penetrated only one time after reconstitution, if needed, using a suitable sterile transfer device or dispensing set which allows measured dispensing of the contents.

After reconstitution, use within two hours if stored at room temperature, or within four hours if stored under refrigeration.

---

### **Reconstituted Bulk Solution Should Not Be Used for Direct Infusion**

---

If the reconstituted bulk solution is stored for less than one hour at room temperature (20°C/68°F) prior to further dilution, the use periods indicated in Table 4 apply for the diluted solutions.

If the bulk solution is stored for one to two hours at room temperature (20°C/68°F) and then diluted with Sterile Water for Injection or 0.9% Sodium Chloride Injection to the following concentrations, the use periods indicated in Table 5 apply.

Any unused portions of solution that remain after the indicated time periods should be discarded.

**TABLE 4**

| <b>Diluent</b>                 | <b>Maximum Concentration (mg/mL)<br/>(Ampicillin and Sulbactam for<br/>Injection)</b> | <b>Use Periods</b> |
|--------------------------------|---------------------------------------------------------------------------------------|--------------------|
| Sterile Water for Injection    | 45 (30/15)                                                                            | 8 hrs at 25°C      |
|                                | 45 (30/15)                                                                            | 48 hrs at 4°C      |
|                                | 30 (20/10)                                                                            | 72 hrs at 4°C      |
| 0.9% Sodium Chloride Injection | 45 (30/15)                                                                            | 8 hrs at 25°C      |
|                                | 45 (30/15)                                                                            | 48 hrs at 4°C      |
|                                | 30 (20/10)                                                                            | 72 hrs at 4°C      |
| 5% Dextrose Injection          | 30 (20/10)                                                                            | 2 hrs at 25°C      |
|                                | 30 (20/10)                                                                            | 4 hrs at 4°C       |
|                                | 3 (2/1)                                                                               | 4 hrs at 25°C      |
| Lactated Ringer's Injection    | 45 (30/15)                                                                            | 8 hrs at 25°C      |
|                                | 45 (30/15)                                                                            | 24 hrs at 4°C      |
| M/6 Sodium Lactate Injection   | 45 (30/15)                                                                            | 8 hrs at 25°C      |
|                                | 45 (30/15)                                                                            | 8 hrs at 4°C       |
| 5% Dextrose in 0.45% Saline    | 3 (2/1)                                                                               | 4 hrs at 25°C      |
|                                | 15 (10/5)                                                                             | 4 hrs at 4°C       |
| 10% Invert Sugar               | 3 (2/1)                                                                               | 4 hrs at 25°C      |
|                                | 30 (20/10)                                                                            | 3 hrs at 4°C       |

**TABLE 5**

| <b>IV Solution</b>                   | <b>Maximum Concentration (mg/mL)<br/>Ampicillin and Sulbactam for<br/>Injection<br/>(Ampicillin/Sulbactam)</b> | <b>Use Period</b> |
|--------------------------------------|----------------------------------------------------------------------------------------------------------------|-------------------|
| Sterile Water for Injection, USP     | 45 (30/15)                                                                                                     | 4 hrs at 21°C     |
|                                      | 45 (30/15)                                                                                                     | 24 hrs at 4°C     |
| 0.9% Sodium Chloride Injection (USP) | 45 (30/15)                                                                                                     | 4 hrs at 21°C     |
|                                      | 45 (30/15)                                                                                                     | 24 hrs at 4°C     |

### **Animal Pharmacology**

While reversible glycogenosis was observed in laboratory animals, this phenomenon was dose- and time-dependent and is not expected to develop at the therapeutic doses and corresponding plasma levels attained during the relatively short periods of combined ampicillin/sulbactam therapy in man.

### **HOW SUPPLIED**

Ampicillin and Sulbactam for Injection, USP (ampicillin sodium/sulbactam sodium) Pharmacy Bulk Package is supplied as follows:

| <b>NDC</b>            | <b>Ampicillin and Sulbactam for Injection, USP</b>                                                                                                 | <b>Package Factor</b> |
|-----------------------|----------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| 72603- <b>444</b> -10 | 15 gram Pharmacy Bulk Package<br>Bottle containing<br>ampicillin sodium and sulbactam sodium<br>equivalent to<br>10 g ampicillin and 5 g sulbactam | 10 vials per tray     |

Ampicillin and Sulbactam for Injection, USP powder should be stored at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature], prior to reconstitution.

**Sterile, Nonpyrogenic, Preservative-free.**

**The container closure is not made with natural rubber latex.**

**To report SUSPECTED ADVERSE REACTIONS, contact Northstar Rx, LLC at 1-800-206-7821 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

Brands listed are the trademarks of their respective owners.

Manufactured for:  
Northstar Rx LLC,  
Memphis, TN 38141.  
Toll-free: 1-800-206-7821

Manufactured by:  
Istituto Biochimico Italiano Giovanni Lorenzini S.p.A.  
Via Fossignano 2, 04011 Aprilia (LT), Italy  
Issued June 2024

860693

**Principal Display Panel - Ampicillin and Sulbactam for Injection, USP 15 gram Bottle Label**

NDC 72603-**444**-01  
Rx only

**Ampicillin and Sulbactam for Injection, USP**

**15 grams\***

**per Pharmacy Bulk Package**

**PHARMACY BULK PACKAGE -**

**NOT FOR DIRECT INFUSION**

**For Intravenous Use**

**NorthStar**

**3** **8**  
**72603 44401**  
 ©2005 Northstar Healthcare Holdings Ltd.  
 Lot: \_\_\_\_\_  
 Exp: \_\_\_\_\_

NDC 72603-444-01 **Rx only**

**Ampicillin and Sulbactam for Injection, USP**  
**15 grams\***  
 per Pharmacy Bulk Package

**PHARMACY BULK PACKAGE— NOT FOR DIRECT INFUSION**

For Intravenous Use

**NORTHSTAR**

**Reconstituted bulk solution should not be used for direct infusion.**

**PHARMACY BULK PACKAGE**  
 Add Sterile Water for Injection or 0.9% Sodium Chloride Injection-stroke well

| Amount of Diluent | Concentration of Solution                  |
|-------------------|--------------------------------------------|
| 92 mL             | ampicillin 100 mg/mL<br>sulbactam 50 mg/mL |

Reconstitution by adding 92 mL of Sterile Water for Injection or 0.9% Sodium Chloride Injection in two separate aliquots. Add 50 mL, shake to dissolve and add 42 mL, shake for final solution. The resulting solution will contain ampicillin 100 mg/mL and sulbactam 50 mg/mL. Transfer individual doses to appropriate intravenous infusion solutions. Discard unused solution from bulk package within 2 hours after initial entry. After reconstitution, no significant loss of potency occurs for 2 hours at room temperature and for 4 hours if refrigerated. Discard unused transferred solutions held longer than these time periods.

A single entry through the vial closure should be made with a sterile dispensing set or transfer device. Transfer individual doses to appropriate intravenous infusion solutions. Use of a syringe with needle is not recommended. The above process should be carried out under a laminar flow hood using aseptic technique.

**RECOMMENDED STORAGE IN DRY FORM.**  
 Ampicillin and Sulbactam for Injection, USP powder should be stored at 20° to 25°C (68° to 77°F) [see USP], prior to reconstitution.  
**DOSAGE AND USE:** See accompanying prescribing information.  
 \*Each bottle contains 15 g ampicillin sodium/sulbactam sodium equivalent to 10 g ampicillin plus 5 g sulbactam.  
 Sterile, Nonpyrogenic, Preservative-free.  
 This container closure is not made with natural rubber latex.

Dealer/Time/Concentration \_\_\_\_\_  
 Mfg. for: Northstar Rx LLC, Memphis, TN 38141 - Iss. 11/2023  
 Mfg. by: Istituto Biotechimico Italiano Giovanni Lorenzini S.p.A. - Italy

3 8  
 Ampicillin and Sulbactam for Injection, USP  
 15 grams\* per Pharmacy Bulk Package

## AMPICILLIN AND SULBACTAM

ampicillin sodium and sulbactam sodium injection, powder, for solution

### Product Information

|                                |                            |                           |               |
|--------------------------------|----------------------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN PRESCRIPTION DRUG    | <b>Item Code (Source)</b> | NDC:72603-444 |
| <b>Route of Administration</b> | INTRAVENOUS, INTRAMUSCULAR |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name                                                            | Basis of Strength | Strength |
|----------------------------------------------------------------------------|-------------------|----------|
| <b>ampicillin sodium</b> (UNII: JFN36L5S8K) (ampicillin - UNII:7C782967RD) | ampicillin        | 10 g     |
| <b>sulbactam sodium</b> (UNII: DKQ4T82YE6) (sulbactam - UNII:S4TF6I2330)   | sulbactam         | 5 g      |

### Packaging

| # | Item Code        | Package Description                              | Marketing Start Date | Marketing End Date |
|---|------------------|--------------------------------------------------|----------------------|--------------------|
| 1 | NDC:72603-444-10 | 10 in 1 CARTON                                   | 03/01/2024           |                    |
| 1 |                  | 1 in 1 BOTTLE; Type 0: Not a Combination Product |                      |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| ANDA               | ANDA065314                               | 03/01/2024           |                    |

**Labeler** - NorthStar Rx, LLC (830546433)

**Registrant** - NorthStar Rx, LLC (830546433)

**Establishment**

| Name                                            | Address | ID/FEI    | Business Operations    |
|-------------------------------------------------|---------|-----------|------------------------|
| Istituto Biochimico Italiano Giovanni Lorenzini |         | 432432581 | manufacture(72603-444) |

Revised: 6/2024

NorthStar Rx, LLC