

**NAPROXEN SODIUM - naproxen sodium tablet, coated**  
**AAA Pharmaceutical, Inc.**

-----  
**Naproxen Sodium Tablets**

***Drug Facts***

**Active ingredient (in each caplet)**

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID<sup>1</sup>)

---

<sup>1</sup> nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
  - the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert**

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

**Stop use and ask a doctor if**

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- you develop heartburn
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- drink a full glass of water with each dose

adults and children 12 years and older	<ul style="list-style-type: none"> <li>▪ take 1 caplet every 8 to 12 hours while symptoms last</li> <li>▪ for the first dose you may take 2 caplets within the first hour</li> <li>▪ do not exceed 2 caplets in any 8- to 12-hour period</li> <li>▪ do not exceed 3 caplets in a 24-hour period</li> </ul>
children under 12 years	<ul style="list-style-type: none"> <li>▪ ask a doctor</li> </ul>

### Other information

- **each caplet contains:** sodium 20 mg
- store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- retain carton for complete product information

### Inactive ingredients

croscarmellose sodium, FD&C blue #2, macrogol, magnesium stearate, polyvinyl alcohol, povidone, pregelatinized starch, talc, titanium dioxide

### Distributed by:

AAA Pharmaceutical, Inc.  
681 Main Street  
Lumberton, NJ 08048

### PRINCIPAL DISPLAY PANEL - 24 Caplet Bottle Carton

#### **RESTORE u**

NDC 57344-145-01

†COMPARE TO THE ACTIVE  
INGREDIENT IN ALEVE®

#### ***Naproxen***

Naproxen Sodium Tablets, 220 mg

***Pain Reliever / Fever Reducer***  
***(NSAID)***

**24 COATED CAPLETS\*\* (\*\*capsule shaped tablets)**

COATING FREE AREA

**Drug Facts (continued)**

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 ■ right before or after heart surgery

**Ask a doctor before use if**  
 ■ you have a history of stomach problems, such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
 ■ you are taking a diuretic  
 ■ you have asthma  
 ■ you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**  
 ■ under a doctor's care for any serious condition  
 ■ taking any other drug

**When using this product**  
 ■ take with food or milk if stomach upset occurs  
 ■ long term continuous use may increase the risk of heart attack or stroke

**Stop use and ask a doctor if**  
 ■ side effects occur. You may report side effects to FDA at 1-800-FDA-1088.  
 ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood  
 ■ have bloody or black stools  
 ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ you have difficulty swallowing  
 ■ it feels like the pill is stuck in your throat  
 ■ you develop heartburn  
 ■ redness or swelling is present in the painful area  
 ■ any new symptoms appear

**Drug Facts**

**Active ingredient (in each caplet)** Purpose  
 Naproxen sodium 220 mg Pain reliever/  
 (naproxen 200 mg) (NSAID)\*,.....fever reducer  
 \*nonsteroidal anti-inflammatory drug

**USES**  
 ■ temporarily relieves minor aches and pains due to:  
 ■ minor pain of arthritis ■ muscular aches  
 ■ headache ■ menstrual cramps  
 ■ toothache  
 ■ the common cold

**Warnings**  
**A very alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing)  
 ■ shock ■ skin redening ■ rash ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinner (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription (or, naproxen, or others)  
 ■ have 3 or more alcoholic drinks every day while using this product  
 ■ take more or for a longer time than directed

INK AND COATING FREE FOR LOT AND EXPIRATION STAMPING

060106189566  
 7

DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

This product is not manufactured or distributed by Bayer Consumer Healthcare, LLC, distributor of Advair®

**RESTORE U** NDC 57344-145-01

COMPARE TO THE ACTIVE INGREDIENT IN ALEVE®

**Naproxen**  
 Naproxen Sodium Tablets, 220 mg  
 Pain Reliever / Fever Reducer (NSAID)

**24 COATED CAPLETS\*\*** (\*\*capsule shaped tablets)

**Drug Facts (continued)**

**Inactive ingredients** croscarmellose sodium, FD&C blue #2, macrogol, magnesium stearate, polyvinyl alcohol, povidone, pregelatinized starch, talc, titanium dioxide

Distributed by:  
 AAA Pharmaceuticals, Inc.  
 681 Main Street  
 Lumberton, NJ 08048

F114501RES\_RT

**Drug Facts (continued)**

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not take more than directed  
 ■ the smallest effective dose should be used  
 ■ do not take longer than 10 days, unless directed by a doctor (see Warnings)  
 ■ drink a full glass of water with each dose

adults and children 12 years and older  
 ■ take 1 caplet every 8 to 12 hours while symptoms last  
 ■ for the first dose you may take 2 caplets within the first hour  
 ■ do not exceed 2 caplets in any 8- to 12-hour period  
 ■ do not exceed 3 caplets in a 24-hour period

children under 12 years ■ ask a doctor

**Other information**  
 ■ each caplet contains: sodium 20 mg  
 ■ store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F).  
 ■ retain carton for complete product information

**NAPROXEN SODIUM**  
 naproxen sodium tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57344-145
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

**Inactive Ingredients**

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONES (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

**Product Characteristics**

Color	BLUE	Score	no score
Shape	OVAL (Biconvex)	Size	12mm
Flavor		Imprint Code	I7
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57344-145-01	1 in 1 CARTON		
1		24 in 1 BOTTLE, PLASTIC		
2	NDC:57344-145-02	1 in 1 CARTON		
2		50 in 1 BOTTLE, PLASTIC		
3	NDC:57344-145-03	1 in 1 CARTON		
3		100 in 1 BOTTLE, PLASTIC		
4	NDC:57344-145-04	1 in 1 CARTON		
4		150 in 1 BOTTLE, PLASTIC		
5	NDC:57344-145-05	1 in 1 CARTON		
5		200 in 1 BOTTLE, PLASTIC		
6	NDC:57344-145-06	1 in 1 CARTON		
6		300 in 1 BOTTLE, PLASTIC		
7	NDC:57344-145-07	500 in 1 BOTTLE, PLASTIC		
8	NDC:57344-145-08	200 in 1 BOTTLE, PLASTIC		
9	NDC:57344-145-10	400 in 1 BOTTLE, PLASTIC		
10	NDC:57344-145-11	600 in 1 BOTTLE, PLASTIC		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079096	12/15/2012	

**Labeler** - AAA Pharmaceutical, Inc. (181192162)

Revised: 1/2013

AAA Pharmaceutical, Inc.