

## **STOMACH RELIEF REGULAR STRENGTH- bismuth subsalicylate liquid Best Choice (Valu Merchandisers Company)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Drug Facts**

#### **Active ingredient (in each 30 mL dose)**

Bismuth subsalicylate 525 mg

#### **Purpose**

Upset stomach reliever/Antidiarrheal

#### **Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - gas
  - indigestion
  - nausea
  - fullness
  - belching

#### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

#### **Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

#### **Do not use**

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are**

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

**When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. do not use any other dosing device
- keep dosing cup with product
- adults and children 12 years and over:
  - 30 mL (1 dose) every 1/2 or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
  - 30 mL (1 dose) every 1/2 as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
  - do not exceed 8 doses (240 mL) in 24 hours
  - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**

- **each 30 mL contains:** potassium 8 mg, sodium 9 mg
- **each 30 mL contains:** salicylate 206 mg
- low sodium

- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

### **Inactive ingredients**

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

### **Principal Display Panel**

Compare to the active ingredient in Pepto-Bismol®\*

REGULAR STRENGTH

### **Stomach Relief**

**BISMUTH SUBSALICYLATE 525 mg**

**UPSET STOMACH RELIEVER**

**/ANTIDIARRHEAL**

For relief of:

- Heartburn
- Indigestion
- Nausea
- Upset stomach
- Diarrhea

Alcohol free

Sugar free

FL OZ (mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

PROUDLY DISTRIBUTED BY:

ASSOCIATED WHOLESALE GROCERS, INC.

KANSAS CITY, KANSAS 66106

### **Package Label**

**Drug Facts (continued)**

**Directions** ■ mL = milliliter ■ shake well before using  
 ■ measure only with dosing cup provided. Do not use any other dosing device.  
 ■ keep dosing cup with product  
 ■ adults and children 12 years and over  
 ■ 30 mL (1 dose) every 1/2 hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea  
 ■ 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)  
 ■ do not exceed 8 doses (240 mL) in 24 hours  
 ■ use until diarrhea stops but not more than 2 days  
 ■ children under 12 years of age: ask a doctor  
 ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL contains: potassium 8 mg, sodium 9 mg  
 ■ each 30 mL contains: salicylate 206 mg ■ low sodium ■ keep tightly closed  
 ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)

**Inactive ingredients** benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

**Questions or comments?** Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company. PLD-C482B LB009086

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

PROUDLY DISTRIBUTED BY:  
 ASSOCIATED WHOLESALE GROCERS, INC.  
 KANSAS CITY, KANSAS 66106  
 BEST CHOICE 20148-0723-PLD

Scan here for more product information. Call 1-844-252-1112 for more product information.

**Best Choice, 100% Guaranteed**  
[www.bestchoicebrand.com](http://www.bestchoicebrand.com)

**Best Choice®**

Compare to the active ingredient in  
**Pepto-Bismol®**

**REGULAR STRENGTH**  
**Stomach Relief**  
**BISMUTH SUBSALICYLATE 525mg**  
**UPSET STOMACH RELIEVER**  
**/ANTIDIARRHEAL**

**For relief of:**

- Heartburn
- Indigestion
- Nausea
- Upset stomach
- Diarrhea

**Alcohol free**  
**Sugar free**

**12 fl oz (355 mL)**

**Drug Facts**

**Active ingredient (in each 30 mL dose)** Purpose  
 Bismuth subsalicylate 525 mg.....Upset stomach reliever/  
 Antidiarrheal

**Uses** relieves ■ diarrhea ■ travelers' diarrhea  
 ■ upset stomach due to overindulgence in food and drink, including  
 ■ heartburn ■ gas ■ indigestion ■ nausea  
 ■ fullness ■ belching

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert: Contains salicylate. Do not take if you are:**  
 ■ allergic to salicylates (including aspirin)  
 ■ taking other salicylate products

**Do not use if you have**  
 ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have**  
 ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for**  
 ■ diabetes ■ gout ■ arthritis  
 ■ anticoagulation (thinning the blood)

**When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.**

**Stop use and ask a doctor if**  
 ■ symptoms get worse or last more than 2 days  
 ■ ringing in the ears or loss of hearing occurs  
 ■ diarrhea lasts more than 2 days

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.**

### BEST CHOICE Regular Strength Stomach Relief

## STOMACH RELIEF REGULAR STRENGTH

bismuth subsalicylate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63941-724
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 7671P0Y5NH)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Packaging

Marketing Start      Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-724-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/29/2019	
2	NDC:63941-724-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/29/2019	
3	NDC:63941-724-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/29/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	11/29/2019	

**Labeler** - Best Choice (Valu Merchandisers Company) (868703513)

Revised: 8/2023

Best Choice (Valu Merchandisers Company)