ITHAPPENZ MORNING AFTER - levonorgestrel tablet Aurohealth LLC

ITHAPPENZTM MORNING AFTER

(levonorgestrel tablet 1.5 mg)

Drug Facts

Active ingredient (in each tablet)

Levonorgestrel USP 1.5 mg

Purpose

Emergency contraceptive

Use

for women to reduce chance of pregnancy after unprotected sex (if a contraceptive failed or if you did not use birth control)

Warnings

Allergy alert: Do not use if you have ever had an allergic reaction to levonorgestrel

Sexually transmitted diseases (STDs) alert: This product does **not** protect against HIV/AIDS or other STDs

Do not use

- if you are already pregnant (because it will not work)
- for regular birth control

Ask a doctor or pharmacist before use if you are

taking efavirenz (HIV medication) or rifampin (tuberculosis treatment) or medication for seizures (epilepsy). These medications may reduce the effectiveness of levonorgestrel.

When using this product you may have

menstrual changes

- nausea
- lower stomach (abdominal) pain
- tiredness
- headache
- dizziness
- breast pain
- vomiting

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- take as soon as possible within 72 hours (3 days) after unprotected sex. The sooner you take it the better it will work.
- if you vomit within 2 hours after taking the medication, call a healthcare professional to find out if you should repeat the dose.

Other information

- read the instructions, warnings and enclosed product leaflet before use
- this product works mainly by preventing ovulation (egg release). It may also prevent fertilization of a released egg (joining of sperm and egg) or attachment of a fertilized egg to the uterus (implantation).
- do not use if carton is open or tear strip is removed or the blister seal is broken or missing
- store at 20° to 25°C (68° to 77°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, lactose monohydrate, magnesium stearate, potato starch and talc.

Questions?

Call **1-855-274-4122**

Distributed by:

AUROHEALTH LLC

279 Princeton-Hightstown Road East Windsor, NJ 08520

Made in India

Code: TS/DRUGS/22/2009

Patient Information

ITHAPPENZTM MORNING AFTER
(levonorgestrel tablet 1.5 mg)
Emergency Contraceptive
One Tablet. One Step.
What You Need to Know
What is ITHAPPENZTM MORNING AFTER?

ITHAPPENZTM MORNING AFTER is emergency contraception that helps prevent pregnancy after birth control failure or unprotected sex. It is a **backup** method of preventing pregnancy and should not be used as regular birth control.

What ITHAPPENZTM MORNING AFTER is not.

ITHAPPENZTM MORNING AFTER will not work if you are already pregnant and will not affect an existing pregnancy. ITHAPPENZTM MORNING AFTER will not protect you from HIV infection (the virus that causes AIDS) and other sexually transmitted diseases (STDs).

When should I use ITHAPPENZTM MORNING AFTER?

The sooner you take emergency contraception, the better it works. You should use ITHAPPENZTM MORNING AFTER within 72 hours (3 days) **after you have had unprotected sex**.

ITHAPPENZTM MORNING AFTER is a backup or emergency method of birth control you can use when:

- your regular birth control was used incorrectly or failed
- you did not use any birth control method

When not to use ITHAPPENZTM MORNING AFTER.

ITHAPPENZTM MORNING AFTER should not be used:

- as a regular birth control method, because it's not as effective as regular birth control.
- if you are already pregnant, because it will not work.
- if you are allergic to levonorgestrel or any other ingredients in ITHAPPENZTM MORNING AFTER.

When should I talk to a doctor or pharmacist?

Ask a doctor or pharmacist before use if you are taking efavirenz (HIV medication) or rifampin (tuberculosis treatment) or medication for seizures (epilepsy). These medications may reduce the effectiveness of ITHAPPENZTM MORNING AFTER and increase your chance of becoming pregnant. Your doctor may prescribe another form of emergency contraception that may not be affected by these medications.

How does ITHAPPENZTM MORNING AFTER work?

ITHAPPENZTM MORNING AFTER is one tablet with levonorgestrel, a hormone that has been used in many birth control pills for several decades. ITHAPPENZTM MORNING AFTER contains a higher dose of levonorgestrel than birth control pills, but works in a similar way to prevent pregnancy. It works mainly by stopping the release of an egg from the ovary. It is possible that ITHAPPENZTM MORNING AFTER may also work by preventing fertilization of an egg (the uniting of sperm with the egg) or by preventing attachment (implantation) to the uterus (womb).

How can I get the best results from ITHAPPENZTM MORNING AFTER?

You have 72 hours (3 days) to try to prevent pregnancy after birth control failure or unprotected sex. The sooner you take ITHAPPENZTM MORNING AFTER, the better it works.

How effective is ITHAPPENZTM MORNING AFTER?

If ITHAPPENZTM MORNING AFTER is taken as directed, it can significantly decrease the chance that you will get pregnant. About 7 out of every 8 women who would have gotten pregnant will not become pregnant.

How will I know ITHAPPENZTM MORNING AFTER worked?

You will know ITHAPPENZTM MORNING AFTER has been effective when you get your next period, which should come at the expected time, or within a week of the expected time. If your period is delayed beyond 1 week, it is possible you may be pregnant. You should get a pregnancy test and follow up with your healthcare professional.

Will I experience any side effects?

- some women may have changes in their period, such as a period that is heavier or lighter or a period that is early or late. If your period is more than a week late, you may be pregnant.
- if you have severe abdominal pain, you may have an ectopic pregnancy, and should get immediate medical attention.
- when used as directed, ITHAPPENZTM MORNING AFTER is safe and effective. Side effects may include changes in your period, nausea, lower stomach (abdominal) pain, tiredness, headache, dizziness, and breast tenderness.
- if you vomit within 2 hours of taking the medication, call a healthcare professional to find out if you should repeat the dose.

What if I still have questions about ITHAPPENZTM MORNING AFTER?

If you have questions or need more information, call our toll-free number AUROHEALTH LLC at 1-855-274-4122.

Other Information

Keep out of reach of children:

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Do not use if the blister seal is opened.

Store at room temperature 20° to 25°C (68° to 77°F).

Active ingredient: levonorgestrel 1.5 mg

Inactive ingredients: colloidal silicon dioxide, corn starch, lactose monohydrate, magnesium stearate, potato starch, and talc.

If you are sexually active, you should see a healthcare provider for routine checkups. Your healthcare provider will talk to you about and, if necessary, test you for sexually transmitted diseases, teach you about effective methods of routine birth control, and answer any other questions you may have.

Distributed by:

AUROHEALTH LLC

279 Princeton-Hightstown Road East Windsor, NJ 08520

Manufactured by:

Aurobindo Pharma Limited

Hyderabad-500 032, India

Issued: 11/2022

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL -1.5 mg (1 Tablet Carton Label)

[†]Compare to the NDC 58602-892-57 active ingredient of **Plan B One-Step**®

ITHAPPENZTM

MORNING AFTER

levonorgestrel tablet 1.5 mg emergency contraceptive

Reduces chance of pregnancy after unprotected sex.

Not for regular birth control.

1 Tablet
One Tablet. One Stop.

Levonorgestrel 1.5 mg



ITHAPPENZ MORNING AFTER

levonorgestrel tablet

_			
Pro	duct	Inform	ation

Product Type HUMAN OTC DRUG Item Code (Source) NDC:58602-892

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

LEVONORGESTREL (UNII: 5W7SIA7YZW) (LEVONORGESTREL - UNII:5W7SIA7YZW) LEVONORGESTREL 1.5 mg

Inactive Ingredients

mactive mg. calends				
Ingredient Name	Strength			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
STARCH, CORN (UNII: O8232NY3SJ)				
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
STARCH, POTATO (UNII: 81089SAH3T)				
TALC (UNII: 7SEV7J4R1U)				

Product Characteristics

Color	WHITE (White to off-white)	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	S;11
Contains			

Packaging

Ш	1			
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 NDC:58602-892-	1 in 1 CARTON	11/14/2022	
	1	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph Citation	Marketing Start	Marketing End
Category		Date	Date
ANDA	ANDA206867	11/14/2022	

Labeler - Aurohealth LLC (078728447)

Establishment				
Name	Address	ID/FEI	Business Operations	
Aurobindo Pharma Limited		650381903	ANALYSIS(58602-892), MANUFACTURE(58602-892)	

Revised: 3/2023 Aurohealth LLC