# CHILDRENS MULTI SYMPTOM FEVER AND COLD- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl suspension Kroger Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

# Kroger Co. Children's Multi-Symptom Fever & Cold Drug Facts

# Active ingredients (in each 5 mL)

Acetaminophen 160 mg Chlorpheniramine maleate 1 mg Dextromethorphan HBr 5 mg Phenylephrine HCl 2.5 mg

## **Purposes**

Pain reliever/fever reducer Antihistamine Cough suppressant Nasal decongestant

#### Uses

- temporarily relieves
- minor aches and pains
- headache
- minor sore throat pain
- runny nose
- itchy nose or throat
- sneezing
- itchy, watery eyes due to hay fever
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily reduces fever

# **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- in a child under 4 years of age
- if your child has ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- to make a child sleepy

# Ask a doctor before use if your child has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

# Ask a doctor or pharmacist before use your child is taking

- sedatives or tranquilizers
- the blood thinning drug warfarin

# When using this product

- do not exceed recommended dosage
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness
- excitability may occur, especially in children

## Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

## Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical even if you do not notice any signs or symptoms.

#### **Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- shake well before using
- remove the child protective cap and squeeze your child's dose into the dosing cup
- may be given every 4 hours, while symptoms persist. Do not give more than 5 doses in 24 hours unless directed by a doctor.
- use enclosed dosing cup only. Keep for use with this product. Do not use any other dosing device.

Age	Dose	
children under 4	do not us e	
years of age		
children 4 to under 6	do not use unless directed by a doctor	
years of age		
children 6 to under 12 years of age	10 mL	

### Other information

- each 5 mL contains: sodium 3 mg
- contains no aspirin
- store at 20-25°C (68-77°F). Protect from light.
- close cap tightly

## **Inactive ingredients**

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, tribasic sodium phosphate, xanthan gum

#### Questions or comments?

1-800-632-6900

#### **Principal Display Panel**

COMPARE TO the active ingredients of TRIAMINIC® MULTI-SYMPTOM FEVER & COLD

See side panel

OUR PHARMACIST RECOMMENDED

Ages 6 TO 11 Years

children's Multi-Symptom Fever & Cold

Acetaminophen Pain Reliever/Fever Reducer

Chlorpheniramine Maleate Antihistamine

Dextromethorphan HBr Cough Suppressant

Phenylephrine HCl Nasal Decongestant

**ORAL SUSPENSION** 

Cold & Flu Symptom Relief

Fever, Runny & Stuffy Nose, Aches & Pains, Sore Throat & Cough

Grape Flavor

4 FL OZ (118 mL)



#### CHILDRENS MULTI SYMPTOM FEVER AND COLD

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl suspension

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-023
Route of Administration	ORAL		

## **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	160 mg in 5 mL
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg in 5 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients		
Ingredient Name	Strength	
CARRAGEENAN (UNII: 5C69 YCD2YJ)		
D&C RED NO. 33 (UNII: 9DBA0SBB0L)		
EDETATE DISO DIUM (UNII: 7FLD9 1C86K)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6A3C0OX)		
POLYSORBATE 80 (UNII: 6OZP39ZG8H)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SORBITOL (UNII: 506T60A25R)		
SUCROSE (UNII: C151H8M554)		
XANTHAN GUM (UNII: TTV12P4NEE)		
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)		
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0 DDB505)		
SO DIUM PHO SPHATE, TRIBASIC (UNII: A752Q30 A6 X)		

Product Characteristics			
Color	PURPLE (opaque)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

	Packaging			
:	# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
	1 NDC:30142-023-26	1 in 1 CARTON	06/23/2015	
	1	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	06/23/2015	

Revised: 5/2020 Kroger Company