

**THERAFLU NIGHTTIME SEVERE COLD AND COUGH- acetaminophen,
diphenhydramine hcl, phenylephrine hcl powder, for solution
GlaxoSmithKline Consumer Healthcare Holdings (US) LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each packet)

Acetaminophen 650 mg
Diphenhydramine HCl 25 mg
Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer
Antihistamine/cough suppressant
Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - runny nose
 - sneezing
 - itchy nose or throat
 - itchy, watery eyes due to hay fever
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may

include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not exceed recommended dosage**
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- **each packet contains:** potassium 10 mg, sodium 23 mg
- **phenylketonurics:** contains phenylalanine 13 mg per packet
- store at controlled room temperature 20°-25°C (68°-77°F). Protect product from heat and moisture.

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, flavors, maltodextrin, silicon dioxide, sodium citrate, sucrose,

tribasic calcium phosphate

Questions or comments?

call **1-855-328-5259**

Additional Information

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.

KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

TAMPER EVIDENT INNER UNIT

DO NOT USE IF SEALED THERAFLU PACKET IS TORN OR BROKEN.

1-855-328-5259

Distributed by: **GSK Consumer Healthcare**

Warren, NJ 07059

©2019 GSK group of companies or its licensor.

Trademarks are owned by or licensed to the GSK group of companies.

SEVERE COLD & COUGH

COUGH

NASAL CONGESTION

SORE THROAT PAIN

HEADACHE

BODY ACHES

FEVER

RUNNY NOSE

SNEEZING

Principal Display Panel

NDC 0067-7918-06

THERAFLU

SEVERE COLD & COUGH

NIGHTTIME

Acetaminophen

Pain Reliever/Fever Reducer

Diphenhydramine HCl

Antihistamine/Cough Suppressant

Phenylephrine HCl

Nasal Decongestant

- ***Cough***
- ***Nasal Congestion***
- ***Sore Throat Pain***
- ***Headache***
- ***Body Ache***
- ***Fever***
- ***Runny Nose***
- ***Sneezing***

6 PACKETS

HONEY LEMON INFUSED WITH CHAMOMILE & WHITE TEA FLAVORS

gsk

62000000033903

NDC 0067-7918-06

gsk

THERAFLU

SEVERE COLD & COUGH

NIGHTTIME

Acetaminophen
Pain Reliever/Fever Reducer

Diphenhydramine HCl
Antihistamine/Cough Suppressant

Phenylephrine HCl
Nasal Decongestant

- ▶ Cough
- ▶ Nasal Congestion
- ▶ Sore Throat Pain
- ▶ Headache
- ▶ Body Ache
- ▶ Fever
- ▶ Runny Nose
- ▶ Sneezing

6 PACKETS

HONEY LEMON
INFUSED WITH
CHAMOMILE &
WHITE TEA FLAVORS

THERAFLU NIGHTTIME SEVERE COLD AND COUGH

acetaminophen, diphenhydramine hcl, phenylephrine hcl powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0067-7918
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 237 mL
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40)	DIPHENHYDRAMINE	25 mg

(DIPHENHYDRAMINE - UNII:8GTS82S83M)	HYDROCHLORIDE	in 237 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 237 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
ASPARTAME (UNII: Z0H242BBR1)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SUCROSE (UNII: C151H8M554)	
TRIBASIC CALCIUM PHOSPHATE (UNII: 91D9GV0Z28)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	HONEY, LEMON (HONEY LEMON INFUSED WITH CHAMOMILE & WHITE TEA FLAVORS)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0067-7918-06	6 in 1 CARTON	07/01/2014	
1	NDC:0067-7918-01	237 mL in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/01/2014	

Labeler - GlaxoSmithKline Consumer Healthcare Holdings (US) LLC (079944263)