

**MELoxicAM, meloxicam tablet**  
Lundbeck Pharmaceuticals (USA), Inc.

**REGIMENTS OF PRESCRIBING INFORMATION**  
These highlights do not include all of the information needed for the MELoxicAM TABLETS USP, safety and effectiveness. See full prescribing information for MELoxicAM TABLETS USP.  
MELoxicAM TABLETS USP, for oral use  
INDICATIONS, Approved 2008

**WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS**  
See full prescribing information for complete boxed warning.

- **Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in treatment and may increase with duration of use (2.1).**
- **Meloxicam tablets are contraindicated in the setting of coronary artery bypass graft (CABG) surgery (2.2).**
- **NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious GI events (2.2).**

**RECENT MAJOR CHANGES**

boxed warning (2018)  
Indications and Usage: Juvenile Rheumatoid Arthritis (JRA) (Pauciarticular and Polyarticular Course) (1.3) (6/2016)  
Dosage and Administration: General Dosing Instructions (2.1) (2/2016)  
Dosage and Administration: Juvenile Rheumatoid Arthritis (JRA) (Pauciarticular and Polyarticular Course) (2.4) (6/2016)  
Warnings and Precautions: Cardiovascular: Thrombotic Events (2.1) (2/2016)  
Warnings and Precautions: Heart Failure and Edema (5.3) (2/2016)

**INDICATIONS AND USAGE**  
Meloxicam tablets are a nonsteroidal anti-inflammatory drug indicated for:

- Osteoarthritis (OA) (1.1)
- Rheumatoid Arthritis (RA) (1.2)
- Juvenile Rheumatoid Arthritis (JRA) in patients who weigh ≥60 kg (1.3)

**DOSE AND ADMINISTRATION**  
Use the boxed warning design for the chosen duration consistent with individual patient treatment goals (2.1)

- **OA (2.1) (see 2.1.1)**  
Starting dose: 7.5 mg once daily  
Dose may be increased to 15 mg once daily
- **RA (2.2)**  
7.5 mg once daily in adults and ≥60 kg
- **Juvenile RA (2.4)** or use in combination with approved formulations of oral meloxicam even if the total meloxicam strength is less than 15 mg

**DOSE FORMS AND STRENGTHS**  
• Meloxicam Tablets USP 7.5 mg and 15 mg (3)

**CONTRAINDICATIONS**

- Known hypersensitivity to any component of the drug product (6)
- History of asthma, urticaria, or other allergic-type reaction after taking aspirin or other NSAIDs (4)
- In the setting of CABG surgery (2)

**WARNINGS AND PRECAUTIONS**

- **Cardiovascular**: Serious patients of warning signs and symptoms of hypersensitivity (Dyspepsia or discomfort of abdominal area, possible nausea or flatulent signs and symptoms of liver disease develop) (5.1)
- **Cardiovascular**: Please advise your prescriber if you have had any heart-related symptoms (5.1)
- **Cardiovascular**: Monitor blood pressure (5.1)
- **GI Bleeding and Ulcers**: Avoid use of Meloxicam in patients with severe heart failure unless benefits are expected to outweigh the risk of worsening heart failure (5.2)
- **GI Bleeding and Ulcers**: Monitor renal function in patients with renal or hepatic impairment, heart failure, dehydration, or hypotension. Avoid use of Meloxicam in patients with advanced renal disease unless benefits are expected to outweigh risk of worsening renal function (5.2)
- **Anaphylactoid Reactions**: Avoid emergency help if an anaphylactoid reaction occurs (5.3)
- **Aspirin-Related Adverse Effects and Other NSAIDs**: Meloxicam is contraindicated in patients with aspirin-sensitive asthma. Those patients who have used aspirin with caution should be advised (5.3)
- **Aspirin-Related Adverse Effects and Other NSAIDs**: Avoid use of Meloxicam in patients with severe heart failure unless benefits are expected to outweigh the risk of worsening heart failure (5.3)
- **Aspirin-Related Adverse Effects and Other NSAIDs**: Avoid use in pregnant women starting at 30 weeks gestation (5.1, 8.1)
- **Aspirin-Related Adverse Effects and Other NSAIDs**: Monitor benefits or harms in patients with signs or symptoms of events (5.1, 7)

**ADVERSE REACTIONS**  
Most common (≥5% and greater than placebo) adverse events in adults are diarrhea, upper respiratory tract infection, dyspepsia, sinusitis, and headache (5.1)

Most common adverse events in pediatric patients were similar to those in the adult rheumatoid arthritis study (5.1)

To report SUSPECTED ADVERSE REACTIONS, contact Lundbeck Pharmaceuticals (USA), Inc. at 1-866-562-4136 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

**DRUG INTERACTIONS**

- **Aspirin**: Avoid use of Meloxicam in patients with aspirin-sensitive asthma. Those patients who have used aspirin with caution should be advised (5.3)
- **Aspirin**: Avoid use of Meloxicam in patients with severe heart failure unless benefits are expected to outweigh the risk of worsening heart failure (5.3)
- **Aspirin**: Avoid use in pregnant women starting at 30 weeks gestation (5.1, 8.1)
- **Aspirin**: Monitor benefits or harms in patients with signs or symptoms of events (5.1, 7)

**USE IN SPECIFIC POPULATIONS**

- **Pediatric**: Use of NSAIDs during the third trimester of pregnancy increases the risk of premature closure of the fetal ductus arteriosus. Avoid use of NSAIDs in pregnant women starting at 30 weeks gestation (5.1, 8.1)
- **Elderly**: NSAIDs are associated with reversible interstitial nephritis. Consider withdrawal of Meloxicam in patients who have interstitial nephritis (5.1)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide. Revised 2/2018

**FULL PRESCRIBING INFORMATION: CONTENTS\***

- 1 INDICATIONS AND USAGE
  - 1.1 Osteoarthritis (OA)
  - 1.2 Rheumatoid Arthritis (RA)
  - 1.3 Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course
- 2 DOSE AND ADMINISTRATION
  - 2.1 General Dosing Instructions
  - 2.2 Osteoarthritis
  - 2.3 Rheumatoid Arthritis
  - 2.4 Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course
  - 2.5 Renal Impairment
  - 2.6 Non-Interchangeability with Other Formulations of Meloxicam
- 3 DOSE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
  - 5.1 Cardiovascular: Thrombotic Events
  - 5.2 Cardiovascular: Bleeding, Ulceration, and Perforation
  - 5.3 Hypotension
  - 5.4 Hypertension
  - 5.5 Heart Failure and Edema
  - 5.6 Renal Toxicity and Hyperkalemia
  - 5.7 Anaphylactoid Reactions
  - 5.8 Exacerbation of Asthma Related to Aspirin Sensitivity
  - 5.9 Stomach Acid Reflux
  - 5.10 Premature Closure of Fetal Ductus Arteriosus
  - 5.11 Hemorrhagic Tendency
  - 5.12 Masking of Inflammation and Fever
  - 5.13 Laboratory Monitoring
- 6 ADVERSE REACTIONS
  - 6.1 Clinical Trial Experience
  - 6.2 Post-Marketing Experience
- 7 DRUG INTERACTIONS
- 8 USE IN SPECIFIC POPULATIONS
  - 8.1 Pregnancy
  - 8.2 Lactation
  - 8.3 Fertility and Modes of Reproductive Potential
  - 8.4 Pediatric Use
  - 8.5 Geriatric Use
  - 8.6 Hepatic Impairment
  - 8.7 Renal Impairment
- 9 OVERDOSAGE
- 10 DESCRIPTION
- 11 CLINICAL PHARMACOLOGY
  - 11.1 Mechanism of Action
  - 11.2 Pharmacokinetics
- 12 NONCLINICAL TOXICOLOGY
  - 12.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13 CLINICAL STUDIES
  - 13.1 Osteoarthritis and Rheumatoid Arthritis
  - 13.2 Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course
- 14 HOW SUPPLIED/STORAGE AND HANDLING
- 15 PATIENT COUNSELING INFORMATION

**FULL PRESCRIBING INFORMATION**

**WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS**

**Cardiovascular Thrombotic Events**

- **Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in treatment and may increase with duration of use (see Warnings and Precautions (5.1)).**
- **Meloxicam tablets are contraindicated in the setting of coronary artery bypass graft (CABG) surgery (see Contraindications (2) and Warnings and Precautions (5.1)).**

**Gastrointestinal Bleeding, Ulceration, and Perforation**

- **NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious GI events (see Warnings and Precautions (5.2)).**

**1 INDICATIONS AND USAGE**

**1.1 Osteoarthritis (OA)**  
Meloxicam tablets are indicated for relief of the signs and symptoms of osteoarthritis (see Clinical Studies (14.1)).

**1.2 Rheumatoid Arthritis (RA)**  
Meloxicam tablets are indicated for relief of the signs and symptoms of rheumatoid arthritis (see Clinical Studies (14.1)).

**1.3 Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course**  
Meloxicam tablets are indicated for relief of the signs and symptoms of pauciarticular or polyarticular course juvenile Rheumatoid Arthritis in patients who weigh ≥60 kg (see Dosage and Administration (2.4) and Clinical Studies (14.2)).

**2 DOSE AND ADMINISTRATION**

**2.1 General Dosing Instructions**  
Carefully consider the potential benefits and risks of Meloxicam tablets and other treatment options before deciding to use Meloxicam tablets. Use the lowest effective dosage for the chosen duration consistent with individual patient treatment goals (see Warnings and Precautions (5)).  
After observing the response to initial therapy with Meloxicam tablets, adjust the dose to suit an individual patient's needs.  
In adults, the maximum recommended daily oral dose of Meloxicam tablets is 15 mg regardless of formulation. In patients with renal failure, a maximum daily dosage of 7.5 mg is recommended (see Use in Specific Populations (8.7) and Clinical Pharmacology (12.3)).  
Meloxicam tablets may be taken without regard to timing of meals.

**2.2 Osteoarthritis**  
For the relief of the signs and symptoms of osteoarthritis the recommended starting and maintenance oral dose of Meloxicam tablets is 7.5 mg once daily. Some patients may receive additional benefit by increasing the dose to 15 mg once daily.

**2.3 Rheumatoid Arthritis**  
For the relief of the signs and symptoms of rheumatoid arthritis, the recommended starting and maintenance oral dose of Meloxicam tablets is 7.5 mg once daily. Some patients may receive additional benefit by increasing the dose to 15 mg once daily.

**2.4 Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course**  
For the treatment of juvenile rheumatoid arthritis, the recommended dose of Meloxicam tablets is 7.5 mg once daily in children who weigh ≥60 kg. There was no additional benefit demonstrated by increasing the dose above 7.5 mg in clinical trials.  
Meloxicam tablets should not be used in children who weigh <60 kg.

**2.5 Renal Impairment**  
The use of Meloxicam tablets in subjects with severe renal impairment is not recommended.

In patients on hemodialysis, the maximum dosage of Meloxicam tablets is 7.5 mg per day (see Clinical Pharmacology (12.3)).

**2.6 Non-Interchangeability with Other Formulations of Meloxicam**  
Meloxicam tablets have not shown equivalent systemic exposure to other approved formulations of oral meloxicam. Therefore, Meloxicam tablets are not interchangeable with other formulations of oral meloxicam products even if the total meloxicam strength is the same. Do not substitute similar dose strengths of Meloxicam tablets with other formulations of oral meloxicam product.

**3 DOSE FORMS AND STRENGTHS**  
Meloxicam Tablets USP:

- 7.5 mg: Light yellow, round flat beveled edge, tablet with U & L debossed on one side and 7.5 debossed centrally on the other side
- 15 mg: Light yellow, capsule shaped, biconvex, tablet with U & L debossed on one side and 15 debossed centrally on the other side

**4 CONTRAINDICATIONS**

Meloxicam tablets are contraindicated in the following patients:

- Known hypersensitivity (e.g., angioedematous reaction and serious skin reaction) to meloxicam or any component of the drug product *See Warnings and Precautions (5.7, 5.9)*
- History of asthma, urticaria, or other allergic-type reactions after taking aspirin or other NSAIDs. Severe, sometimes fatal, anaphylactic reactions to NSAIDs have been reported in such patients *See Warnings and Precautions (5.7, 5.9)*
- In the setting of coronary artery bypass graft (CABG) surgery *See Warnings and Precautions (5.1)*

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Cardiovascular Thrombotic Events

Clinical trials of several COX-2 selective and nonselective NSAIDs of up to three years duration have shown an increased risk of serious cardiovascular (CV) thrombotic events, including myocardial infarction (MI) and stroke, which can be fatal. Based on available data, it is unclear that the risk for CV thrombotic events is similar for all NSAIDs. The relative increase in serious CV thrombotic events over baseline conferred by NSAID use appears to be similar in patients with and without known CV disease or risk factors for CV disease. However, patients with known CV disease or risk factors had a higher absolute incidence of excess serious CV thrombotic events, due to their increased baseline rate. Some observational studies found that this increased risk of serious CV thrombotic events began as early as the first weeks of treatment. The increase in CV thrombotic risk has been observed most consistently at higher doses.

To minimize the potential risk for an adverse CV event in NSAID-treated patients, use the lowest effective dose for the shortest duration possible. Physicians and patients should remain alert for the development of such events, throughout the entire treatment course, even in the absence of previous CV symptoms. Patients should be informed about symptoms of serious CV events and the steps to take if they occur.

There is no conclusive evidence that concurrent use of aspirin mitigates the increased risk of serious CV thrombotic events associated with NSAID use. The concurrent use of aspirin and an NSAID, such as meloxicam, increases the risk of serious gastrointestinal (GI) events *See Warnings and Precautions (5.2)*.

### 5.2 Serious Postoperative Bleeding: Risks of CABG Surgery

Two large, controlled clinical trials of a COX-2 selective NSAID for the treatment of pain in the first 10-14 days following CABG surgery found an increased incidence of myocardial infarction and stroke. NSAIDs are contraindicated in the setting of CABG *See Contraindications (4)*.

### 5.3 MI and Stroke

Observational studies conducted in the Danish National Registry have demonstrated that patients treated with NSAIDs in the post-acute period were at increased risk of myocardial infarction, CV-related death, and all-cause mortality beginning in the first week of treatment. In this same cohort, the incidence of death in the first year post-acute MI was 20 per 100 person-years in NSAID-treated patients compared to 12 per 100 person-years in non-NSAID exposed patients. Although the absolute rate of death declined somewhat after the first year post-MI, the increased relative risk of death in NSAID users persisted over at least the first four year of follow-up.

Avoid the use of Meloxicam in patients with a recent MI unless the benefits are expected to outweigh the risk of recurrent CV thrombotic events. If Meloxicam is used in patients with a recent MI, monitor patients for signs of cardiac ischemia.

### 5.2 Gastrointestinal Bleeding, Ulceration, and Perforation

NSAIDs, including meloxicam, can cause serious gastrointestinal (GI) adverse events including inflammation, bleeding, ulceration, and perforation of the esophagus, stomach, small intestine, or large intestine, which can be fatal. These serious adverse events can occur at any time, with or without warning symptoms, in patients treated with NSAIDs. Only one of four patients who develop a serious upper GI adverse event on NSAID therapy is symptomatic. Upper GI ulcers, gross bleeding, or perforation caused by NSAIDs occurred in approximately 1% of patients treated for 3-6 months, and in about 2-4% of patients treated for one year. However, even short-term NSAID therapy is not without risk.

### 5.3 Risk Factors for GI Bleeding, Ulceration, and Perforation

Patients with a prior history of peptic ulcer disease and/or GI bleeding who used NSAIDs had a greater than 10-fold increased risk for developing a GI bleed compared to patients without these risk factors. Other factors that increase the risk of GI bleeding in patients treated with NSAIDs include longer duration of NSAID therapy; concomitant use of oral corticosteroids, aspirin, anticoagulants, or selective serotonin reuptake inhibitors (SSRIs); smoking; use of alcohol; older age; and poor general health status. Most postmarketing reports of fatal GI events occurred in elderly or debilitated patients. Additionally, patients with advanced liver disease and/or coagulopathy are at increased risk for GI bleeding.

### 5.4 Strategies to Minimize the GI Risks in NSAID-treated Patients

- Use the lowest effective dosage for the shortest possible duration.
- Avoid administration of more than one NSAID at a time.
- Avoid use in patients at higher risk unless benefits are expected to outweigh the increased risk of bleeding. For such patients, as well as those with active GI bleeding, consider alternate therapies other than NSAIDs.
- Remains alert for signs and symptoms of GI ulceration and bleeding during NSAID therapy.
- If a serious GI adverse event is suspected, promptly initiate evaluation and treatment, and discontinue Meloxicam until a serious GI adverse event is ruled out.
- In the setting of concomitant use of low-dose aspirin for cardiac prophylaxis, monitor patients more closely for evidence of GI bleeding *See Drug Interactions (7)*.

### 5.3 Hepatotoxicity

Elevations of ALT or AST (three or more times the upper limit of normal [ULN]) have been reported in approximately 1% of NSAID-treated patients in clinical trials. In addition, rare, sometimes fatal, cases of severe hepatic injury, including fulminant hepatitis, liver necrosis, and hepatic failure have been reported.

Elevations of ALT or AST (less than three times ULN) may occur in up to 15% of patients treated with NSAIDs including meloxicam.

Inform patients of the warning signs and symptoms of hepatotoxicity (e.g., nausea, fatigue, lethargy, diarrhea, pruritus, jaundice, right upper quadrant tenderness, and “flu-like” symptoms). If clinical signs and symptoms consistent with liver disease develop, or if systemic manifestations occur (e.g., eosinophilia, rash, etc.), discontinue Meloxicam immediately and perform a clinical evaluation of the patient *See Use in Specific Populations (8.6) and Clinical Pharmacology (12.3)*.

### 5.4 Hypertension

NSAIDs, including Meloxicam, can lead to new onset or worsening of preexisting hypertension, either of which may contribute to the increased incidence of CV events. Patients taking antihypertensive converting enzyme (ACE) inhibitors, thiazide diuretics, or loop diuretics may have impaired response to these therapies when taking NSAIDs *See Drug Interactions (7)*.

Monitor blood pressure (BP) during the initiation of NSAID treatment and throughout the course of therapy.

### 5.5 Heart Failure and Edema

The Cox-2 and traditional NSAID Trialists' Collaboration meta-analysis of randomized controlled trials demonstrated an approximately two-fold increase in hospitalization for heart failure in COX-2 selective-treated patients and nonselective NSAID-treated patients compared to placebo-treated patients. In Danish National Registry study of patients with heart failure, NSAID use increased the risk of MI, hospitalization for heart failure, and death.

Additionally, fluid retention and edema have been observed in some patients treated with NSAIDs. Use of meloxicam may blunt the CV effects of several therapeutic agents used to treat these medical conditions (e.g., diuretics, ACE inhibitors, or angiotensin receptor blockers [ARBs]) *See Drug Interactions (7)*.

Avoid the use of Meloxicam in patients with severe heart failure unless the benefits are expected to outweigh the risk of worsening heart failure. If Meloxicam is used in patients with severe heart failure, monitor patients for signs of worsening heart failure.

### 5.6 Renal Toxicity and Hypokalemia

**Renal Toxicity**  
Long-term administration of NSAIDs, including Meloxicam, has resulted in renal papillary necrosis, renal insufficiency, acute renal failure, and other renal injury.

Renal toxicity has also been seen in patients in whom renal prostaglandins have a compensatory role in the maintenance of renal perfusion. In these patients, administration of an NSAID may cause a dose-dependent reduction in prostaglandin formation and, secondarily, in renal blood flow, which may precipitate overt renal decompensation. Patients at greatest risk of this reaction are those with impaired renal function, dehydration, hypovolemia, heart failure, liver dysfunction, those taking diuretics and ACE inhibitors or ARBs, and the elderly. Discontinuation of NSAID therapy is usually followed by recovery to the pretreatment state.

The renal effects of Meloxicam may hasten the progression of renal dysfunction in patients with preexisting renal disease. Because some Meloxicam metabolites are excreted by the kidney, monitor patients for signs of worsening renal function.

Correct volume status in dehydrated or hypovolemic patients prior to initiating Meloxicam. Monitor renal function in patients with renal or hepatic impairment, heart failure, aortic stenosis, or hypertension during use of Meloxicam *See Drug Interactions (7)*.

No information is available from controlled clinical studies regarding the use of Meloxicam in patients with advanced renal disease. Avoid the use of Meloxicam in patients with advanced renal disease unless the benefits are expected to outweigh the risk of worsening renal function. If Meloxicam is used in patients with advanced renal disease, monitor patients for signs of worsening renal function *See Clinical Pharmacology (12.3)*.

### 5.7 Hypokalemia

Increases in serum potassium concentration, including hyperkalemia, have been reported with use of NSAIDs, even in some patients without renal impairment. Patients with renal insufficiency, these effects have been attributed to a hyporeninemic-hypaldosteronism state.

### 5.7 Anaphylactic Reactions

Meloxicam has been associated with anaphylactic reactions in patients with and without known hypersensitivity to meloxicam and in patients with aspirin-sensitive asthma *See Contraindications (4) and Warnings and Precautions (5.9)*.

### 5.8 Exacerbation of Asthma Related to Aspirin Sensitivity

A subgroup of patients with asthma may have aspirin-sensitive asthma which may include chronic rhinitis complicated by nasal polyps, and/or potentially fatal bronchospasm, and/or intolerance to aspirin and other NSAIDs. Because cross-reactivity between aspirin and NSAIDs has been reported in such aspirin-sensitive patients, Meloxicam is contraindicated in patients with this form of aspirin sensitivity *See Contraindications (4)*. When Meloxicam is used in patients with preexisting asthma (whether known aspirin sensitivity), monitor patients for changes in the signs and symptoms of asthma.

### 5.9 Serious Skin Reactions

NSAIDs, including meloxicam, can cause serious skin adverse reactions such as exfoliative dermatitis, Stevens-Johnson Syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. These serious events may occur without warning. Inform patients about the signs and symptoms of serious skin reactions, and to discontinue the use of Meloxicam at the first appearance of skin rash or any other sign of hypersensitivity. Meloxicam is contraindicated in patients with previous serious skin reactions to NSAIDs *See Contraindications (4)*.

### 5.10 Premature Closure of Fetal Ductus Arteriosus

Meloxicam may cause premature closure of the fetal ductus arteriosus. Avoid use of NSAIDs, including Meloxicam, in pregnant women starting at 30 weeks of gestation (third trimester) *See Use in Specific Populations (8.1)*.

### 5.11 Hematologic Toxicity

Anemia has occurred in NSAID-treated patients. This may be due to occult or gross blood loss, fluid retention, or an idiosyncratic decreased effect on erythropoiesis. If a patient treated with Meloxicam has any sign or symptoms of anemia, monitor hemoglobin or hematocrit.

NSAIDs, including Meloxicam, may increase the risk of bleeding events. Co-medical conditions such as coagulation disorders or concomitant use of warfarin, other anticoagulants, antiplatelet agents (e.g., aspirin), serotonin reuptake inhibitors (SSRIs), and serotonin norepinephrine reuptake inhibitors (NRNIs) may increase this risk. Monitor these patients for changes in the signs and symptoms of anemia.

### 5.12 Masking of Inflammation and Fever

The pharmacological activity of Meloxicam in reducing inflammation, and possibly fever, may diminish the utility of diagnostic signs in detecting infections.

### 5.13 Laboratory Monitoring

Because serious GI bleeding, hepatotoxicity, and renal injury can occur without warning symptoms or signs, consider monitoring patients on long-term (NSAID) treatment with a CBC, and a chemistry profile periodically *See Warnings and Precautions (5.2, 5.3, 5.9)*.

## 6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- Cardiovascular Thrombotic Events *See Blood Warnings and Precautions (5.1)*
- GI Bleeding, Ulceration, and Perforation *See Blood Warnings and Precautions (5.2)*
- Hypertension *See Warnings and Precautions (5.4)*
- Heart Failure and Edema *See Warnings and Precautions (5.5)*
- Renal Toxicity and Hypokalemia *See Warnings and Precautions (5.6)*
- Anaphylactic Reactions *See Warnings and Precautions (5.7)*
- Serious Skin Reactions *See Warnings and Precautions (5.9)*
- Hematologic Toxicity *See Warnings and Precautions (5.11)*

### 6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

### 6.2.1

#### Disorders and Rheumatoid Arthritis

The Meloxicam Phase 2b clinical trial database includes 10,122 OA patients and 1012 RA patients treated with Meloxicam 7.5 mg/day, 1505 OA patients and 1311 RA patients treated with Meloxicam 15 mg/day. Meloxicam in these doses was administered to 661 patients for at least 6 months and to 312 patients for at least one year. Approximately 10,200 of these were treated in placebo- and/or active-controlled observational trials and 2863 of these were treated in placebo- and/or active-controlled treatment arthritis trials. Gastrointestinal (GI) adverse events were the most frequently reported adverse events in all treatment groups across Meloxicam trials.

A 12-week multicenter, double-blind, randomized trial was conducted in patients with osteoarthritis of



## Labor or Delivery

There are no studies on the effects of Meloxicam during labor or delivery. In animal studies, NSAIDs, including meloxicam, inhibit prostaglandin synthesis, cause delayed parturition, and increase the incidence of stillbirth.

## Data

### Animal Data

Meloxicam was not teratogenic when administered to pregnant rats during fetal organogenesis at oral doses up to 4 mg/kg/day (2.6-fold greater than the MRHD of 15 mg of Meloxicam based on BSA comparison). Administration of meloxicam to pregnant rabbits throughout embryogenesis produced an increased incidence of spinal defects of the heart at an oral dose of 60 mg/kg/day (78-fold greater than the MRHD based on BSA comparison). The no effect level was 20 mg/kg/day (26-fold greater than the MRHD based on BSA comparison). In rats and rabbits, embryofetotoxicity occurred at oral meloxicam doses of 1 mg/kg/day and 2 mg/kg/day, respectively (1.6-fold greater, respectively, than the MRHD based on BSA comparison) when administered throughout organogenesis.

Oral administration of meloxicam to pregnant rats during gestation through lactation increased the incidence of dystocia, delayed parturition, and decreased offspring survival at meloxicam doses of 0.125 mg/kg/day or greater (0.08-times MRHD based on BSA comparison).

### Human Data

There are no human data available on whether meloxicam is present in human milk, or on the effects on breastfed infants, or on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Meloxicam and any potential adverse effects on the breastfed infant from the Meloxicam from the underlying maternal condition.

## Data

### Animal Data

Meloxicam was present in the milk of lactating rats at concentrations higher than those in plasma.

### 8.3 Female and Male of Reproductive Potential

#### Infertility

##### Females

Based on the mechanism of action, the use of prostaglandin-mediated NSAIDs, including Meloxicam may delay or prevent rupture of ovarian follicles, which has been associated with reversible infertility in some women. Published animal studies have shown that administration of prostaglandin synthesis inhibitors has the potential to disrupt prostaglandin-mediated follicular rupture required for ovulation. Small studies in women treated with NSAIDs have also shown a reversible delay in ovulation. Consider withdrawal of NSAIDs, including Meloxicam, in women who have difficulties conceiving or who are undergoing investigations of infertility.

#### 8.4 Pediatric Use

The safety and effectiveness of meloxicam in pediatric JRA patients from 2 to 17 years of age has been evaluated in three clinical trials (see Dosage and Administration (2.3), Adverse Reactions (6.1) and Clinical Studies (14.2)).

#### 8.5 Geriatric Use

Elderly patients, compared to younger patients, are at greater risk for NSAID-associated serious cardiovascular, gastrointestinal, and/or renal adverse reactions. If the anticipated benefit for the elderly patient outweighs these potential risks, start dosing at the low end of the dosing range, and monitor patients for adverse effects (see Warnings and Precautions (6.1, 6.2, 6.5, 6.6, 6.7)).

#### 8.6 Hepatic Impairment

No dose adjustment is necessary in patients with mild to moderate hepatic impairment. Patients with severe hepatic impairment have not been adequately studied. Since meloxicam is significantly metabolized in the liver and hepatotoxicity may occur, use meloxicam with caution in patients with hepatic impairment (see Warnings and Precautions (5.3) and Clinical Pharmacology (12.3)).

#### 8.7 Renal Impairment

No dose adjustment is necessary in patients with mild to moderate renal impairment. Patients with severe renal impairment have not been studied. The use of Meloxicam in subjects with severe renal impairment is not recommended. In patients on hemodialysis, meloxicam should not exceed 7.5 mg per day. Meloxicam is not dialyzable (see Dosage and Administration (2.1) and Clinical Pharmacology (12.3)).

## 10 OVERDOSAGE

Symptoms following acute NSAID overdosage have been typically limited to headache, dizziness, nausea, vomiting, and epigastric pain, which have been generally reversible with supportive care. Gastrointestinal bleeding has occurred. Hypertension, acute renal failure, secondary depression, and coma have occurred, but were rare (see Warnings and Precautions (6.1, 5.2, 5.4, 5.6)).

Monitor patients with symptoms and supportive care following an NSAID overdosage. There are no specific antidotes. Consider emesis and/or activated charcoal (60 to 100 grams in adults, 1 to 2 gram per kg of body weight in pediatric patients) within 4 hours, unless a symptomatic patient vomits within four hours of ingestion or in patients with a large overdosage (5 to 10 times the recommended dosage). Forced diuresis, alkalization of urine, hemodialysis, or hemoperfusion may not be useful due to high protein binding.

There is limited experience with meloxicam overdosage. Cholestyramine is known to accelerate the clearance of meloxicam. Accelerated removal of meloxicam by 4 g oral doses of cholestyramine given three times a day was demonstrated in a clinical trial. Administration of cholestyramine may be useful following an overdosage.

For additional information about overdosage treatment, call a poison control center (1-800-222-1222).

## 11 DESCRIPTION

Meloxicam Tablets USP are a nonsteroidal anti-inflammatory drug (NSAID). Each tablet contains 7.5 mg of 5 mg meloxicam for oral administration. Meloxicam is chemically designated as 4-chlorobenzoic-2-methyl-3-(5-methyl-2-thienyl)-2H-1,2-benzothiazine-3-carboxamide-1,1-dioxide. The molecular weight is 321.4. Its empirical formula is C<sub>17</sub>H<sub>15</sub>N<sub>2</sub>O<sub>4</sub>S<sub>2</sub> and it is in the following structural formula:



### Chemical Structure

Meloxicam is a pale yellow solid, practically insoluble in water, with higher solubility observed in strong acids and bases. It is very slightly soluble in methanol. Meloxicam has an apparent partition coefficient (log P) of 1.1 in octanol-buffered pH 7.4. Meloxicam has pKa values of 1.1 and 4.2. Meloxicam is available as a tablet for oral administration containing 7.5 mg or 15 mg meloxicam.

The inactive ingredients in Meloxicam tablets USP include colloidal silicon dioxide, croscarmellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyvinylpyrrolidone and xanthan gum dyeblack.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Meloxicam has analgesic, anti-inflammatory, and antipyretic properties.

The mechanism of action of Meloxicam, like that of other NSAIDs, is not completely understood but involves inhibition of cyclooxygenase (COX-1 and COX-2).

Meloxicam is a potent inhibitor of prostaglandin synthesis *in vitro*. Meloxicam concentration reached during therapy here produced *in vivo* effects. Prostaglandins sensitize afferent nerves and potentiate the action of bradykinin in inducing pain animal models. Prostaglandins are mediators of inflammation. The case meloxicam is an inhibitor of prostaglandin synthesis, its mode of action may be due to a decrease of prostaglandin in peripheral tissues.

### 12.3 Pharmacokinetics

#### Absorption

The absolute bioavailability of meloxicam capsules was 89% following a single oral dose of 30 mg compared with 30 mg IV bolus injection. Following single intravenous doses, dose-proportional pharmacokinetics were shown in the range of 5 mg to 60 mg. After multiple oral doses the pharmacokinetics of meloxicam capsules were dose-proportional over the range of 7.5 mg to 15 mg. Mean C<sub>max</sub> was achieved within four to five hours after a 7.5 mg meloxicam tablet was taken under fast conditions, indicating a prompt drug absorption. Following multiple dosing, steady-state concentrations were reached by Day 5. A second meloxicam concentration peak occurs around 12 to 14 hours post-dose suggesting biliary recycling. Meloxicam capsules have been shown to be bioequivalent to Meloxicam tablets.

Table 4 Single Dose and Steady-State Pharmacokinetic Parameters for Oral 7.5 mg and 15 mg Meloxicam (Mean and % CV)<sup>a</sup>

Pharmacokinetic Parameters (% CV)	Steady State			
	Healthy male adults (7.5 mg capsules)	Elderly males (7.5 mg capsules)	Elderly females (7.5 mg capsules)	Renal failure (15 mg capsules)
N	12	5	12	12
C <sub>max</sub> (ng/mL)	1.05 (20)	1.3 (26)	0.75 (26)	0.75 (26)
t <sub>max</sub> (h)	4.9 (8)	6 (27)	4.0 (5)	10 (87)
t <sub>1/2</sub> (h)	21 (40)	24 (54)	16 (40)	16 (29)
C <sub>12h</sub> (ng/mL)	0.8 (20)	0.9 (26)	0.7 (22)	0.9 (43)
AUC <sub>0-12h</sub> (ng·h/mL)	14.7 (25)	15 (42)	10 (30)	20 (44)

<sup>a</sup> The parameters values in the table are from various studies.

<sup>b</sup> *n* = under high fat conditions

<sup>c</sup> Meloxicam tablets

<sup>d</sup> V<sub>d</sub>1 = dose/(AUC×k<sub>e</sub>)

#### Food and Anacid Effects

Administration of meloxicam capsules following a high fat breakfast (75 g of fat) resulted in mean peak drug levels (i.e., C<sub>max</sub>) being increased by approximately 22%, while the extent of absorption (AUC) was unchanged. The time to maximum concentration (T<sub>max</sub>) was achieved between 5 and 6 hours. No pharmacokinetic interaction was detected with concurrent administration of atidazole. Based on these results, Meloxicam can be administered without regard to timing of meals or concurrent administration of atidazole.

#### Distribution

The mean volume of distribution (V<sub>d</sub>) of meloxicam is approximately 10 L. Meloxicam is ~99.4% bound to human plasma proteins (primarily albumin) within the therapeutic dose range. The fraction of protein binding is independent of drug concentration, over the clinically relevant concentration range, but decreases to ~95% in patients with renal disease. Meloxicam penetration into human red blood cells, after oral dosing, is less than 10%. Following a radiolabeled dose, over 90% of the radioactivity derived in the plasma was present as unchanged meloxicam.

Meloxicam concentrations in synovial fluid, after a single oral dose, range from 40% to 50% of those in plasma. The free fraction in synovial fluid is 2.5 times higher than in plasma, due to the lower albumin content in synovial fluid as compared to plasma. The significance of this penetration is unknown.

#### Elimination

Meloxicam is extensively metabolized in the liver. Meloxicam metabolites include 5-carboxy meloxicam (60% of dose), from P-450 mediated metabolism by oxidation of an imine-carboxamide metabolite 5-hydroxymethyl meloxicam which is also excreted in a lesser extent (9% of dose). *In vitro* studies indicate that CYP2C3 (cytochrome P450 metabolizing enzyme) plays an important role in this metabolic pathway with a minor contribution of the CYP3A4 isoenzyme. P-glycoprotein activity is probably responsible for the dose-to-metabolites which account for 10% and 4% of the administered dose, respectively. All the four metabolites are not known to have any pharmacological activity.

#### Excretion

Meloxicam excretion is predominantly in the form of metabolites, and occurs to equal extent in the urine and feces. Only traces of the unchanged parent compound are excreted in the urine (0.2%) and feces (1.6%). The extent of the urinary excretion was confirmed for radiolabeled multiple 7.5 mg doses: 0.5%, 0%, and 13% of the dose were found in urine in the form of meloxicam, and the 5-hydroxymethyl and 5-carboxy metabolites, respectively. There is significant biliary and/or renal excretion of the drug. This was demonstrated when oral administration of cholestyramine following a single IV dose of meloxicam decreased the AUC of meloxicam by 20%.

The mean elimination half-life (t<sub>1/2</sub>) ranges from 15 hours to 20 hours. The elimination half-life is constant across dose levels indicating linear metabolism within the therapeutic dose range. Plasma clearance ranges from 7 to 9 mL/min.

#### Specific Population

##### Pediatric

After single (0.25 mg/kg) dose administration and after achieving steady state (0.375 mg/kg/day), there was a general trend of approximately 20% lower exposure in younger patients (2 to 6 years old) compared to the older patients. The older patients had meloxicam exposure similar (single dose) or slightly reduced (steady state) to those in the adult patients, when using AUC values normalized to a dose of 0.25 mg/kg (see Dosage and Administration (2.4)). The meloxicam mean (SD) elimination half-life was 15.2 (5.1) and 13.0 (5.0) for the 2 to 6 year old patients, and 7 to 16 year old patients, respectively.

In a covariate analysis, utilizing population pharmacokinetics, body-weight, but not age, was the single predictive covariate for differences in the meloxicam apparent oral plasma clearance. The body-weight normalized apparent oral clearance values were adequate predictors of meloxicam exposure in pediatric patients.

The pharmacokinetics of Meloxicam in pediatric patients under 2 years of age have not been investigated.

##### Geriatric

Elderly males (65 years of age) exhibited meloxicam plasma concentrations and steady-state pharmacokinetics similar to young males. Elderly females (65 years of age) had a 47% higher AUCs and 23% higher C<sub>max</sub> as compared to younger females (55 years of age) after both body weight normalization. Despite the increased oral concentrations in the elderly females, the adverse event profile was comparable for both elderly patient populations. A smaller free fraction was found in elderly female patients in comparison to elderly male patients.

##### Sex

Young females exhibited slightly lower plasma concentrations relative to young males. After single doses of 7.5 mg Meloxicam, the mean elimination half-life was 19.5 hours for the female group as compared to 23.4 hours for the male group. At steady state, the data were similar (173 hours vs 174

hours). This pharmacokinetic difference due to gender is likely to be of little clinical importance. There was linearity of pharmacokinetics and no appreciable difference in the C<sub>max</sub> or T<sub>max</sub> across genders.

#### Hepatic Impairment

Following a single 15 mg dose of meloxicam there was no marked difference in plasma concentration in patients with mild (Child-Pugh Class I) or moderate (Child-Pugh Class II) hepatic impairment compared to healthy volunteers. Protein binding of meloxicam was not affected by hepatic impairment. No dosage adjustment is necessary in patients with mild to moderate hepatic impairment. Patients with severe hepatic impairment (Child-Pugh Class III) have not been adequately studied [see Warnings and Precautions (5.3) and Use in Specific Populations (8.6)].

#### Renal Impairment

Meloxicam pharmacokinetics have been investigated in subjects with mild and moderate renal impairment. Total drug plasma concentrations of meloxicam decreased and renal clearance of meloxicam increased with the degree of renal impairment while free AUC<sub>0-24</sub> values were similar in all groups. The higher meloxicam clearance in subjects with renal impairment may be due to increased fractional renal excretion of meloxicam which is available for hepatic metabolism and subsequent excretion. No dosage adjustment is necessary in patients with mild to moderate renal impairment. Patients with severe renal impairment have not been adequately studied. The use of Meloxicam in subjects with severe renal impairment is not recommended [see Dosage and Administration (2.5), Warnings and Precautions (5.4) and Use in Specific Populations (8.7)].

#### Hemodialysis

Following a single dose of meloxicam, the free C<sub>max</sub> plasma concentrations were higher in patients with renal failure on chronic hemodialysis (1% free fraction) in comparison to healthy volunteers (0.2% free fraction). Hemodialysis did not lower the total drug concentration in plasma; therefore, additional doses are not necessary after hemodialysis. Meloxicam is not dialyzable [see Dosage and Administration (2.1) and Use in Specific Populations (8.7)].

#### Drug Interaction Studies

**Aspirin:** When NSAIDs were administered with aspirin, the protein binding of NSAIDs were reduced, although the clearance of free NSAID was not altered. When Meloxicam was administered with aspirin (1000 mg three times daily) to healthy volunteers, it tended to increase the AUC<sub>0-24</sub> (10%) and C<sub>max</sub> (24%) of meloxicam. The clinical significance of this interaction is not known. See Table 2 for clinically significant drug interactions of NSAIDs with aspirin [see Drug Interactions (7)].

**Cholelithiasis:** Pretreatment for four days with cholestyramine significantly increased the clearance of meloxicam by 50%. This resulted in a decrease in t<sub>1/2</sub> from 19.2 hours to 12.5 hours, and a 37% reduction in AUC<sub>0-24</sub>. This suggests the existence of a recirculating pathway for meloxicam in the gastrointestinal tract. The clinical relevance of this interaction has not been established.

**Cimetidine:** Concurrent administration of 200 mg cimetidine four times daily did not alter the single-dose pharmacokinetics of 30 mg meloxicam.

**Digoxin:** Meloxicam 15 mg once daily for 7 days did not alter the plasma concentration profile of digoxin after a single administration for 7 days at clinical doses. In vivo testing found no protein binding drug interaction between digoxin and meloxicam.

**Lithium:** In a study conducted in healthy subjects, mean pre-dose lithium concentration and AUC<sub>0-24</sub> were increased by 27% in subjects receiving lithium doses ranging from 16 to 1072 mg twice daily with meloxicam 15 mg QD every day as compared to subjects receiving lithium alone [see Drug Interactions (7)].

**Methotrexate:** A study in 13 rheumatoid arthritis (RA) patients evaluated the effects of multiple doses of meloxicam on the pharmacokinetics of methotrexate tolerance weeks. Meloxicam did not have significant effect on the pharmacokinetics of single doses of methotrexate. In vivo, methotrexate did not displace meloxicam from its human serum binding sites [see Drug Interactions (7)].

**Warfarin:** The effect of meloxicam on the anticoagulant effect of warfarin was studied in a group of healthy subjects receiving daily doses of warfarin (produced an INR (International Normalized Ratio) between 1.2 and 1.8). In these subjects, meloxicam did not alter warfarin pharmacokinetics and the average anticoagulant effect of warfarin as determined by prothrombin time. However, one subject showed an increase in INR from 1.5 to 2.1. Caution should be used when administering Meloxicam with warfarin in patients on warfarin therapy who are hospitalized in INR and an increased risk of bleeding complications when a new medication is introduced [see Drug Interactions (7)].

## 11 NONCLINICAL TOXICOLOGY

### 11.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

#### Carcinogenesis

There was no increase in tumor incidence in long-term carcinogenicity studies in rats (104 weeks) and mice (87 weeks) administered meloxicam oral doses up to 0.8 mg/kg/day in rats and up to 8.0 mg/kg/day in mice (up to 0.5- and 2.6-times, respectively, the maximum recommended human dose (MRHD) of 15 mg/day Meloxicam based on body surface area (BSA) comparison).

#### Mutagenesis

Meloxicam was not mutagenic in Ames assay, or clastogenic in a chromosome aberration assay with human lymphocytes and in vivo micronucleus test in mouse bone marrow.

#### Impairment of Fertility

Meloxicam did not impair male and female fertility in rats at oral doses up to 8 mg/kg/day in males and 5 mg/kg/day in females (up to 5.8- and 3.2-times doses, respectively, than the MRHD based on BSA comparison).

## 14 CLINICAL STUDIES

### 14.1 Osteoarthritis and Rheumatoid Arthritis

The use of Meloxicam in the treatment of the signs and symptoms of osteoarthritis of the knee and hip was evaluated in a 12-week, double-blind, controlled trial. Meloxicam (7.5 mg, 7.5 mg, and 15 mg daily) was compared to placebo. The four primary endpoints were investigator's global assessment, patient global assessment, patient pain assessment, and total WOMAC score (a self-administered questionnaire addressing pain, function, and stiffness). Patients on Meloxicam 7.5 mg daily and Meloxicam 15 mg daily showed significant improvement in each of these endpoints compared with placebo.

The use of Meloxicam for the management of the signs and symptoms of osteoarthritis was evaluated in six double-blind, active-controlled trials outside the U.S. ranging from 6 weeks to 6 months' duration. In these trials, the efficacy of Meloxicam, in doses of 7.5 mg/day and 15 mg/day, was comparable to placebo or 20 mg/day and diclofenac SR 100 mg/day and consistent with the efficacy seen in the U.S. trial.

The use of Meloxicam for the treatment of the signs and symptoms of rheumatoid arthritis was evaluated in a 12-week, double-blind, controlled medication trial. Meloxicam (7.5 mg, 15 mg, and 22.5 mg daily) was compared to placebo. The primary endpoint in this study was the ACR20 response rate, a composite measure of clinical, laboratory, and functional measures of RA response. Patients receiving Meloxicam 7.5 mg and 15 mg daily showed significant improvement in the primary endpoint compared with placebo. No incremental benefit was observed with the 22.5 mg dose compared to the 15 mg dose.

### 14.2 Juvenile Rheumatoid Arthritis (JRA) Pauciarticular and Polyarticular Course

The use of Meloxicam for the treatment of the signs and symptoms of pauciarticular or polyarticular course Juvenile Rheumatoid Arthritis in patients 2 years of age and older was evaluated in two 12-week, double-blind, parallel-arm, active-controlled trials.

Both studies included three arms: naproxen and two doses of meloxicam. In both studies, meloxicam dosing began at 0.25 mg/kg/day (7.5 mg maximum) or 0.25 mg/kg/day (15 mg maximum), and naproxen dosing began at 10 mg/kg/day. The study used these doses throughout the 12-week dosing period, while the other two groups started after 4 weeks at doses of 0.25 mg/kg/day and 0.375 mg/kg/day (22.5 mg maximum of meloxicam and 15 mg/kg/day of naproxen).

The efficacy analysis used the ACR Pediatric 30 responder definition, a composite of parent and investigator assessments, count of active joints and joints with limited range of motion, and erythrocyte sedimentation rate. The proportion of responders were similar in all three groups in both studies, and no difference was observed between the meloxicam dose groups.

### 14 HOW SUPPLIED/STORAGE AND HANDLING

Meloxicam tablets USP are available as a light yellow, round, film-coated, uncoated tablet containing meloxicam 7.5 mg or as a light yellow, oblong, film-coated, uncoated tablet containing meloxicam 15 mg. The 7.5 mg tablet is imprinted with letter 'L' and 15 on one side and tablet code '75' on the other side. The 15 mg tablet is imprinted with letter 'L' and 15 on one side and tablet code '15' on the other side.

Meloxicam Tablets USP 7.5 mg are available as follows:

NDC 29300-124-13, Bottles of 30

NDC 29300-124-01, Bottles of 100

NDC 29300-124-10, Bottles of 1,000

NDC 29300-124-50, Bottles of 5,000

Meloxicam Tablets USP 15 mg are available as follows:

NDC 29300-125-13, Bottles of 30

NDC 29300-125-01, Bottles of 100

NDC 29300-125-10, Bottles of 1,000

NDC 29300-125-50, Bottles of 5,000

**Storage:** Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Keep

Meloxicam Tablets USP in a dry place.

Dispense tablets in a tight container.

Keep this and all medications out of the reach of children.

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide) that accompanies each prescription dispensed.

**Additional Medication Guides can be obtained by calling Unichem at 1-866-562-4616.**

Inform patients, families or their caregivers of the following information before initiating therapy with an NSAID and periodically during the course of ongoing therapy.

#### Cardiovascular Thrombotic Events

Advise patients to be alert for the symptoms of cardiovascular thrombotic events, including chest pain, shortness of breath, weakness, or slurring of speech, and to report any of these symptoms to their healthcare provider immediately [see Warnings and Precautions (5.1)].

#### Concomitant Bleeding, Ulceration, and Perforation

Advise patients to report symptoms of ulceration and bleeding, including epigastric pain, dyspepsia, melena, and hematemesis to their healthcare provider. In the setting of concomitant use of low-dose aspirin for cardiac prophylaxis, inform patients of the increased risk for the signs and symptoms of GI bleeding [see Warnings and Precautions (5.2)].

#### Hepatotoxicity

Inform patients of the warning signs and symptoms of hepatotoxicity (e.g., nausea, fatigue, lethargy, diarrhea, pruritus, jaundice, right upper quadrant tenderness, and "flu-like" symptoms). If these occur, instruct patients to stop Meloxicam tablets and seek immediate medical therapy [see Warnings and Precautions (5.3)].

#### Heart Failure and Edema

Advise patients to be alert for the symptoms of congestive heart failure including shortness of breath, unexplained weight gain, or edema and to contact their healthcare provider if such symptoms occur [see Warnings and Precautions (5.5)].

#### Anaphylactic Reactions

Inform patients of the signs of anaphylactic reaction (e.g., difficulty breathing, swelling of the face or throat). Instruct patients to seek immediate emergency help if these occur [see Contraindications (4) and Warnings and Precautions (5.7)].

#### Serious Skin Reactions

Advise patients to stop Meloxicam tablets immediately if they develop any type of rash and to contact their healthcare provider as soon as possible [see Warnings and Precautions (5.9)].

#### Female Fertility

Advise females of reproductive potential who desire pregnancy that NSAIDs, including Meloxicam tablets, may be associated with a reversible delay in ovulation [see Use in Specific Populations (8.3)].

#### Fetal Toxicity

Inform pregnant women to avoid use of Meloxicam tablets and other NSAIDs starting at 30 weeks gestation because of the risk of premature closure of the fetal ductus arteriosus [see Warnings and Precautions (8.4) and Use in Specific Populations (8.5)].

#### Avoid Concomitant Use of NSAIDs

Inform patients that the concomitant use of Meloxicam tablets with other NSAIDs or salicylates (e.g., difflural, salicylate) is not recommended due to the increased risk of gastrointestinal toxicity, and little or no increase in efficacy [see Warnings and Precautions (5.2) and Drug Interactions (7)]. Advise patients that NSAIDs may be present in "over the counter" medications for treatment of colds, fever, or headache.

#### Use of NSAIDs and Low-Dose Aspirin

Inform patients not to use low-dose aspirin concomitantly with Meloxicam tablets until they talk to their healthcare provider [see Drug Interactions (7)].

For current prescribing information, call Unichem at 1-866-562-4616.

#### Manufactured by

UNICHEM LABORATORIES LTD.

Pleasant Hill, Estate,

Pleasant Hill, Giza 40511, India

Manufactured for:

UNICHEM  
PHARMACEUTICALS (USA) INC.

Unichem Logo

East Brunswick, NJ 08816

08-8-062019

13011603

NPL MEDGUIDE

**Medication Guide for Nonsteroidal Anti-inflammatory Drugs (NSAIDs)**

**What is the most important information I should know about medicines called Nonsteroidal Anti-inflammatory Drugs (NSAIDs)?**  
 NSAIDs can cause serious side effects, including:  
 • Increased risk of a heart attack or stroke that can lead to death. This risk may happen early in treatment and may increase:  
 ◦ with increasing doses of NSAIDs  
 ◦ with longer use of NSAIDs.  
 • Do not take NSAIDs right before or after a heart surgery called a "coronary artery bypass graft (CABG)."  
 • Avoid taking NSAIDs after a recent heart attack, unless your healthcare provider tells you to. You may have an increased risk of another heart attack if you take NSAIDs after a recent heart attack.  
 • Increased risk of bleeding, ulcers, and tears (perforation) of the esophagus (tube leading from the mouth to the stomach), stomach and intestines:  
 ◦ anytime during use  
 ◦ without warning symptoms  
 ◦ that may cause death  
 The risk of getting an ulcer or bleeding increases with:  
 ◦ past history of stomach ulcers, or stomach or intestinal bleeding with use of NSAIDs  
 ◦ taking medicines called "corticosteroids," "anti-coagulants," "SSRIs," or "SNRIs"  
 ◦ increasing doses of NSAIDs  
 ◦ longer use of NSAIDs  
 ◦ smoking  
 ◦ drinking alcohol  
 ◦ older age  
 ◦ poor health  
 ◦ advanced liver disease  
 ◦ bleeding problems  
 NSAIDs should only be used:  
 ◦ exactly as prescribed  
 ◦ at the lowest dose possible for your treatment  
 ◦ for the shortest time needed

**What are NSAIDs?**  
 NSAIDs are used to treat pain and redness, swelling, and heat (inflammation) from medical conditions such as different types of arthritis, menstrual cramps, and other types of rheumatic pain.

**When should not take NSAIDs?**  
 Do not take NSAIDs:  
 • if you have had an asthma attack, hives, or other allergic reactions with aspirin or any other NSAID, right before or after heart bypass surgery.  
**Before taking NSAIDs, tell your healthcare provider about all of your medical conditions, including if you:**  
 • have liver or kidney problems  
 • have high blood pressure  
 • have asthma  
 • are pregnant or plan to become pregnant. Talk to your healthcare provider if you are considering taking NSAIDs during pregnancy. You should not take NSAIDs after 29 weeks of pregnancy.  
 • are breastfeeding or plan to breast feed.  
**Tell your healthcare provider about all of the medicines you take, including prescription or over-the-counter medicines, vitamins or herbal supplements.** NSAIDs and some other medicines can interact with each other and cause serious side effects. Do not start taking any new medicine without talking to your healthcare provider first.

**What are the possible side effects of NSAIDs?**  
 NSAIDs can cause serious side effects, including:  
**See "What is the most important information I should know about medicines called Nonsteroidal Anti-inflammatory Drugs (NSAIDs)?"**  
 • new or worse high blood pressure  
 • heart failure  
 • liver problems including liver failure  
 • kidney problems including kidney failure  
 • low red blood cells (anemia)  
 • life-threatening skin rashes  
 • life-threatening allergic reaction  
 • Other side effects of NSAIDs include stomach pain, constipation, diarrhea, gas, heartburn, nausea, vomiting, and dizziness.  
**Get emergency help right away if you get any of the following symptoms:**  
 • chest pain or trouble breathing  
 • dizziness  
 • weakness in one part or side of your body  
 • slurred speech  
 • swelling of the face or throat

**Stop taking your NSAID and call your healthcare provider right away if you get any of the following symptoms:**  
 • Nausea  
 • new fluid or weaker than usual  
 • diarrhea  
 • itching  
 • your skin or eyes look yellow  
 • indigestion or stomach pain  
 • flu-like symptoms  
 • small blood  
 • there is blood in your bowel movement or it is black and sticky like tar  
 • unusual weight gain  
 • skin rash or blisters with fever  
 • swelling of the arms, legs, hands and feet

**If you take too much of your NSAID, call your healthcare provider or get medical help right away.**  
 There are not all of the possible side effects of NSAIDs. For more information, ask your healthcare provider or pharmacist about NSAIDs.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**Other information about NSAIDs:**  
 Aspirin is an NSAID but it does not increase the chance of a heart attack. Aspirin can cause bleeding in the brain, stomach, and intestines. Aspirin can also cause ulcers in the stomach and intestines. Some NSAIDs are sold in lower doses without a prescription (over-the-counter). Talk to your healthcare provider before using over-the-counter NSAIDs for more than 10 days.

**General information about the safe and effective use of NSAIDs:**  
 Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use NSAIDs for a condition for which it was not prescribed. Do not give NSAIDs to other people, even if they have the same symptoms that you have. It may harm them.  
 You would like the more information about NSAIDs, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about NSAIDs that is written for health professionals.

**Additional Medication Guides can be obtained by calling Unichem at 1-866-562-4616.**  
 The other trademarks referenced are owned by third parties not affiliated with Unichem Laboratories Limited.

Manufactured by:  
**UNICHEM LABORATORIES LTD.**  
 Plot No. 16, Estate,  
 Plot No. 16, Estate, Gurgaon, Haryana, India  
 Manufactured for:  
  
 East Rutherford, NJ 08816  
 875-662-0219  
 30115623

This Medication Guide has been approved by the U.S. Food and Drug Administration.  
 Revised: June 2019

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL**



**NDC 20300-124-15**  
 30 TABLETS  
 Meloxicam Tablets USP 7.5 mg  
 Manufactured by UNICHEM LABORATORIES LTD.  
 Plot No. 16, Estate, Gurgaon, Haryana, India  
 Manufactured for UNICHEM PHARMACEUTICALS (USA), INC.  
 East Rutherford, NJ 08816  
 875-662-0219  
 30115623

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL**



**NDC 20300-125-15**  
 30 TABLETS  
 Meloxicam Tablets USP 15 mg  
 Manufactured by UNICHEM LABORATORIES LTD.  
 Plot No. 16, Estate, Gurgaon, Haryana, India  
 Manufactured for UNICHEM PHARMACEUTICALS (USA), INC.  
 East Rutherford, NJ 08816  
 875-662-0219  
 30115623

MELOXICAM			
meloxicam tablet			
<b>Product Information</b>			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC 20300-01
Route of Administration	ORAL		
<b>Active Ingredient/Active Moiety</b>			
Ingredient Name	Ratio of Strength	Strength	
MELoxicAM (UNE VEGICORIC) (MELoxicAM - UNICHEM (US))		MELoxicAM	7.5 mg
<b>Inactive Ingredients</b>			
Ingredient Name	Strength		
CELLULOSE MICROCRYSTALLINE (UNE OPERICUS)			
CRUONIPYRONE (UNE 25V 20116)			
LACTOSE MONOHYDRATE (UNE TAYGUSIA)			
MAGNESIUM HYDROXIDE (UNE TAYGUSIA)			
POLYBUTYLENE (UNE 11720 3812)			
SILICON DIOXIDE (UNE 11720 3812)			
TITANIUM DIOXIDE (UNE 11720 3812)			
<b>Product Characteristics</b>			
Color	YELLOW	Shape	16 3039
Shape	OVAL	Marking	None
Flavor		Registration Code	02,75
Contains			
<b>Packaging</b>			
# (US Code)	Package Description	Marketing Start Date	Marketing End Date
1, NDC 20300-01-01 (30 in 1 BOTTLE, Type 1, Not a Combination Product)		11/07/2017	
2, NDC 20300-01-01 (30 in 1 BOTTLE, Type 1, Not a Combination Product)		11/07/2017	
3, NDC 20300-01-01 (30 in 1 BOTTLE, Type 1, Not a Combination Product)		11/07/2017	
4, NDC 20300-01-01 (30 in 1 BOTTLE, Type 1, Not a Combination Product)		11/07/2017	
<b>Marketing Information</b>			
Marketing Category	Application Number and Monograph Classification	Marketing Start Date	Marketing End Date
ANDA	ANDA07927	11/07/2017	

MELOXICAM			
meloxicam tablet			
<b>Product Information</b>			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC 20300-01
Route of Administration	ORAL		
<b>Active Ingredient/Active Moiety</b>			
Ingredient Name	Ratio of Strength	Strength	
MELoxicAM (UNE VEGICORIC) (MELoxicAM - UNICHEM (US))		MELoxicAM	15 mg
<b>Inactive Ingredients</b>			
Ingredient Name	Strength		
CELLULOSE MICROCRYSTALLINE (UNE OPERICUS)			
CRUONIPYRONE (UNE 25V 20116)			
LACTOSE MONOHYDRATE (UNE TAYGUSIA)			
MAGNESIUM HYDROXIDE (UNE TAYGUSIA)			
POLYBUTYLENE (UNE 11720 3812)			
SILICON DIOXIDE (UNE 11720 3812)			
TITANIUM DIOXIDE (UNE 11720 3812)			
<b>Product Characteristics</b>			
Color	YELLOW	Shape	16 3039
Shape	OVAL	Marking	None
Flavor		Registration Code	02,15

Combiner				
<b>Packaging</b>				
#	Item Code	Packaging Description	Marketing Start Date	Marketing End Date
1	NDC-28393-025-01	30 in 1 BOTTLE, Type 0, Not a Combination Product	11/01/2017	
2	NDC-28393-025-02	60 in 1 BOTTLE, Type 2, Not a Combination Product	11/01/2017	
3	NDC-28393-025-03	120 in 1 BOTTLE, Type 0, Not a Combination Product	11/01/2017	
4	NDC-28393-025-04	240 in 1 BOTTLE, Type 0, Not a Combination Product	11/01/2017	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA17927	11/01/2017		

Labeler - Unichem Pharmaceuticals (USA), Inc. (1832013)

Revised: 2/2020

Unichem Pharmaceuticals (USA), Inc.