

**REGULAR STRENGTH ANTACID AND GAS RELIEF- aluminum hydroxide,
magnesium hydroxide, dimethicone suspension
RITE AID PHARMACY**

riteaid ANTACID original

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 200 mg (equivalent to dried gel, USP)
Magnesium hydroxide 200 mg
Simethicone 20mg

Purposes

Antacid
Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Do not take more than 24 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
- children under 12 years: ask a doctor

Other information

- each 5 mL teaspoonful contains: **magnesium 85 mg, sodium 3 mg**
- store at room temperature
- protect from freezing
- keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments?

1-800-540-3765

package Label

Compare to the active ingredient in
Regular Strength Mylanta®*

NDC 11822-3324-7

REGULAR STRENGTH ANTACID & GAS RELIEF

ALUMINUM HYDROXIDE 200 mg
MAGNESIUM HYDROXIDE 200 mg
SIMETHICONE 20 mg

FAST ACTING
Soothing relief of:
heartburn • acid indigestion
sour stomach • pressure
& bloating

218-05112-1 REV 0223

33247



**ORIGINAL
FLAVOR**
CONTAINS LESS
THAN 0.15% ALCOHOL

12 FL OZ (355 mL)

Drug Facts

TAMPER-EVIDENT: Do not use if breakaway band on bottle cap is missing or broken.

Active ingredients

(in each 5 mL teaspoonful)

Active ingredients	Purpose
Aluminum hydroxide (equivalent to dried gel, USP) 200 mg	Antacid
Magnesium hydroxide 200 mg	Antacid
Simethicone 20 mg	Antigas

Uses relieves • heartburn • sour stomach • acid indigestion
• the symptoms referred to as gas

Warnings

Do not take more than 24 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks.

Ask a doctor before use if you have

• kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Directions

• shake well before use
• adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
• children under 12 years: ask a doctor

Other information

• each 5 mL teaspoonful contains: magnesium 85 mg, sodium 3 mg
• store at room temperature • protect from freezing
• keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

*RITE AID is not affiliated with the owner of the registered trademark MYLANTA®
918-05112-1 REV GC0223

DISTRIBUTED BY: RITE AID,
200 NEWBERRY COMMONS
LETTERS, PA 17319
www.riteaid.com



SATISFACTION GUARANTEE:
If you're not satisfied, we'll
happily refund your money.

REGULAR STRENGTH ANTACID AND GAS RELIEF

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-3324
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE -	ALUMINUM	200 mg

UNII:5QB0T2IUN0)	HYDROXIDE	in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	200 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	20 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-3324-7	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2014	
2	NDC:11822-3324-8	710 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2014	11/01/2018

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	04/01/2014	

Labeler - RITE AID PHARMACY (014578892)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(11822-3324) , pack(11822-3324)

Revised: 11/2023

RITE AID PHARMACY