

**COLD AND FLU DAY TIME MULTI-SYMPTOM RELIEF- acetaminophen,
dextromethorphan hbr, phenylephrine hcl liquid
Pharmacy Value Alliance, LLC**

Cold & Flu Day Time Multi-Symptom Relief

Drug Facts

Active Ingredients

(in each 15mL, 1tablespoon)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5mg

Purpose

Acetaminophen Pain reliever/fever reducer
Dextromethorphan HBr Cough suppressant
Phenylephrine HCl Nasal decongestant

Uses temporarily relieves these common cold/flu symptoms:

- minor aches and pains • headache • sore throat • fever
- nasal congestion • cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if adult/child takes:

- more than 4doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or a pharmacist before taking this product.

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacists.

Ask a doctor before use if you have

- liver disease • heart disease • thyroid disease • diabetes
- high blood pressure • trouble urinating due to enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema • cough accompanied by excessive phlegm (mucus)

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

When using this product • do not use more than directed (see overdose warning) • avoid alcoholic drinks

Stop use and ask a doctor if • you get nervous, dizzy or sleepless
• new symptoms occur • fever gets worse or lasts more than 3days
• pain or cough gets worse or lasts more than 5days (children) or 7days (adults) • redness or swelling is present
• cough comes back or occurs with rash or headache that lasts.
These could be signs of a serious condition.

If pregnant or breast-feeding ask a health professional before use. Keep this and all drugs out of the reach of children.

Overdose Warning: Taking more than the recommended dose (overdose) could cause serious health problems, including liver damage. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep this and all drugs out of the reach of children.

Keep this and all drugs out of the reach of children.

Directions

• take only as recommended (see overdose warning) • use dosage cup or tablespoon (TBSP) • do not exceed 4doses per 24 hours

age	dose
adults and children 12 years and over	30 mL (2 TBSP) every 4hours
children 6 to under 12 years	15 mL (1 TBSP) every 4 hours
children 4 years to under 6 years	ask a doctor
children under 4 years	do not use

• When using Day Time and Night Time products, carefully read each label to ensure correct dosing.

Other information

• sodium content per tablespoon: 10 mg
• store at room temperature

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN
Failure to follow these warnings could result in serious consequences

Inactive ingredients

citric acid, FD&C Yellow #6, flavor, glycerin, propylene glycol

purified water, saccharin sodium, sodium benzoate, sucrose

Questions? Call weekdays **1-877-798-5944**.

*This product is not manufactured or distributed by Proctor & Gamble, owner of the registered trademark Vicks® NyQuil®.

DISTRIBUTED BY: PHARMACY VALUE ALLIANCE, LLC
407 EAST LANCASTER AVENUE, WAYNE PA 19087

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

Principal Display Panel

Premier® Value

COMPARE TO THE ACTIVE

INGREDIENTS IN

VICKS® DAYQUIL®*

COLD & FLU

Day Time Multi-Symptom Relief

Acetaminophen..... Pain Reliever/Fever Reducer
Dextromethorphan Hsr..... Cough Suppressant
Phenylephrine HCl..... Nasal Decongestant

Non-drowsy

8 FL OZ (237 mL)

INDEPENDENTLY TESTED

PV

SATISFACTION GUARANTEED

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN

Failure to follow these warnings could result in serious consequences

Questions? Call weekdays **1-877-798-5944**.

*This product is not manufactured or distributed by Proctor & Gamble, owner of the registered trademark Vicks® NyQuil®.

DISTRIBUTED BY: PHARMACY VALUE ALLIANCE, LLC
407 EAST LANCASTER AVENUE, WAYNE PA 19087

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

LR-028

REV-01

Product Label - 8 FL OZ

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN
 Failure to follow these warnings could result in serious consequences

Drug Facts

Active Ingredients (in each 15 mL, 1 tablespoon)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses temporarily relieves these common cold/flu symptoms:
 • minor aches and pains • headache • sore throat • fever
 • nasal congestion • cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if adult/child takes:
 • more than 4 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen
 • 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
 If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or a pharmacist before taking this product.
 • with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacists.

Ask a doctor before use if you have
 • liver disease • heart disease • thyroid disease • diabetes

Premier Value

COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® DAYQUIL™

COLD & FLU

Day Time Multi-Symptom Relief

Acetaminophen..... Pain Reliever/Fever Reducer
 Dextromethorphan HBr..... Cough Suppressant
 Phenylephrine HCl..... Nasal Decongestant

Non-drowsy

8 FL OZ (237 mL)



Drug Facts (continued)

- high blood pressure • trouble urinating due to enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema • cough accompanied by excessive phlegm (mucus)

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

When using this product • do not use more than directed (see overdose warning) • avoid alcoholic drinks

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • new symptoms occur • fever gets worse or lasts more than 3 days • pain or cough gets worse or lasts more than 5 days (children) or 7 days (adults) • redness or swelling is present • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding ask a health professional before use. **Keep this and all drugs out of the reach of children.**

Overdose Warning: Taking more than the recommended dose (overdose) could cause serious health problems, including liver damage. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

LR-027
REV 01



8 40986 03476 3

DRUG FACTS CONTINUED ON BACK

Peel Here

Drug Facts (continued)

Directions
 • take only as recommended (see overdose warning) • use dosage cup or tablespoon (TBSP) • do not exceed 4 doses per 24 hours

age	dose
adults and children 12 years and over	30 mL (2 TBSP) every 4 hours
children 6 to under 12 years	15 mL (1 TBSP) every 4 hours
children 4 years to under 6 years	ask a doctor
children under 4 years	do not use

• When using Day Time and Night Time products, carefully read each label to ensure correct dosing.

Other information
 • sodium content per tablespoon: 10 mg
 • store at room temperature

Inactive ingredients
 citric acid, FD&C Yellow # 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sucrose

Questions? Call weekdays 1-877-798-5944.
 *This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks® NyQuil®.
 DISTRIBUTED BY: PHARMACY VALUE ALLIANCE, LLC
 407 EAST LANCASTER AVENUE, WAYNE PA 19087

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

Product Label - 12 FL OZ

Premier Value Cold & Flu Day Time Multi-Symptom Relief

COLD AND FLU DAY TIME MULTI-SYMPTOM RELIEF			
acetaminophen, dextromethorphan hbr, phenylephrine hcl liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-236
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE	5 mg in 15 mL
Inactive Ingredients			
Ingredient Name			Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
GLYCERIN (UNII: PDC6A3C0OX)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			

SODIUM BENZOATE (UNII: OJ245FE5EU)

SUCROSE (UNII: C151H8M554)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-236-08	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	
2	NDC:68016-236-12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/01/2018	

Labeler - Pharmacy Value Alliance, LLC (101668460)

Registrant - AptaPharma Inc. (790523323)

Establishment

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(68016-236)

Revised: 12/2023

Pharmacy Value Alliance, LLC