

TOPCARE PAIN RELIEF- acetaminophen tablet
Topco Associates LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topco Associates LLC. Pain Relief Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
liver disease

Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,
ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"> • take 2 caplets every 6 hours while symptoms last • do not take more than 6 caplets in 24 hours, unless directed by a doctor • do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store at 20-25°C (68-77°F)

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	L484
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-484-76	1 in 1 CARTON	07/15/1987	
1		120 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:36800-484-62	1 in 1 CARTON	07/15/1987	
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:36800-484-71	1 in 1 CARTON	07/15/1987	
3		50 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:36800-484-78	1 in 1 CARTON	07/15/1987	
4		100 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:36800-484-90	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/15/1987	
6	NDC:36800-484-82	2 in 1 CARTON	07/15/1987	
6		100 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:36800-484-47	150 in 1 BOTTLE; Type 0: Not a Combination Product	07/15/1987	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	07/15/1987	

Labeler - Topco Associates LLC (006935977)

Revised: 4/2022

Topco Associates LLC