POVIDONE IODINE- povidone iodine 10% liquid Humco Holding Group, Inc.

Private Label Povidone Iodine

Drug Facts

Active Ingredient

Povidone-Iodine 10% (1% available iodine)

Purpose

First Aid Antiseptic

Use

First aid antiseptic to prevent infection in minor cuts and burns.

Warnings

For External Use Only

Ask a doctor before use if you have

- deep punture wounds
- animal bites
- serious burns

Stop use and consult a doctor if

- the condition persists or gets worse
- irritation and redness develop and persits for more than 72 hours

When using this product do not

- use in eyes
- use on individuals who are allergic or sensitive to iodine or use for longer thn 1 week unless directed by a doctor
- apply over large areas of the body

Keep out of reach of children.

In case of accidental ingestion, seek professional assistance or contact a Poison Control

Center immediately.

Directions

Clean the affected area. Apply a small amount on the area 1 to 3 times daily. May be covered with sterile bandage. If badaged, let it dry first.

Other information

store at room temperature, 20-25C (68-77F)

Inactive Ingredients

Citric Acid, Disodium Phosphate, Glycerin, nonoxynol-9. Sodium Hydroxide and Purified Water.


Sunmark Label

8 floz






Good Neighbor Label







Leader Label



Warnings For external use only.

- animal bites
- serious burns

Stop use and consult a doctor if

- the condition persists or gets worse
- you need to use this product for more than 1 week

When using this product do not

- use in the eyes
- use if you are allergic to povidone-iodine or any other ingredients in this preparation
- apply over large areas of the body

Keep out of reach of children. If swallowed, seek medical help or contact a Poison Control Center (1-800-222-1222) immediately.

Continued >

Purpose



NOC 70000-0402-1

Povidone**lodine**

10% Topical Solution First Aid Antiseptic

Kills Germs Promptly in Minor Cuts, Scrapes and Bums

+ SAFETY SEALED

COMPARE TO BETADINE active ingredient

8 FL OZ (237 mL)

*Warning: Do not use if Tamper Evident Seal is broken or missing. This product is sealed with a

Drug Facts (continued)

Directions

■ clean the affected area ■ apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first.

Other information

store at room temperature, 20-25° C (68-77° F)

Inactive ingredients

Citric Acid, Disodium Phosphale, Glycerin Nonoxynol-9, Sodium Hydroxide and Purified Wate

Questions or comments? 1-800-662-3435

"This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine*



© 2019 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

DISTRIBUTED BY CARDINAL HEALTH DUBLIN, OHIO 43017 www.myleader.com 1-800-200-6315



Purpose

NDC 37205-186-34





*SAFETY SEALED

Povidonelodine Solution

10% Topical Solution

Antiseptic / Germicide



Promptly Kills Germs In Cuts, Burns, Scrapes and Minor Skin Wounds

WARNING: Do not use if Tamper Evident Seal is broken or missing This product is sealed with a foil seal under cap

8 FL OZ (237mL)

Drug Facts

Active ingredient

Povidone-lodine, 10% (Equivalent to 1% titratable iodine),

■ Health-care antiseptic for preparation of the skin prior to surgery

First aid antiseptic to help prevent infection in minor cuts, scrapes and burns.

For external use only. Do not use

Warnings

- in the eyes
- as a first aid antiseptic longer than one week
- on individuals who are allergic or sensitive to iodine
- m over large areas of the body

Stop use and ask a doctor

- if condition persists or gets worse
- if irritation and redness develop and persist for over 72 hours.

Ask a doctor in case of

- deep puncture wounds
- animal bites
- serious burns

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

For preparation of the skin prior to surgery:

- apply to the operative site prior to surgery.
- As a first aid antiseptic:
- clean the affected area
- apply a small amount of this product to the area 1 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first.

Other information Store at room temperature: 15° - 30° C (59° - 86° F)

Inactive ingredients citric acid, disodium phosphate, glycerin, nonoxynol-9, sodium hydroxide, and purified water

This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark BETADINE®.

0 9 0

R052311RLG

All Leader® Brand Products are 100% satisfaction guaranteed or returnto place of purchase for a full retund.

DISTRIBUTED BY: CARDINAL HEALTH DUBLIN, OHIO 43017 CIN 2443554 www.myleader.com 1-800-200-5313

Quality Choice Label

Active ingredient

Purpose

Povidone-lodine 10%......First Aid Antiseptic (*Equivalent to 1% titratable iodine)

Indications

First aid antiseptic to prevent infection in minor cuts, scrapes and burns

Warnings For external use only.

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites

serious burns

Stop use and consult a doctor if

- the condition persists or gets worse
- irritation and redness develop and persists for more than 72 hours

When using this product do not

- use in the eyes
- use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor
- apply over large areas of the body

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center Immediately.

Continued ▶



NDC 63868-230-08

Compare to active ingredient in Betadine

Povidonelodine

Topical Antiseptic Germicide

10% Topical Solution

Helps Prevent Infection In Cuts, Burns, Scrapes and Minor Skin Wounds

For external use only



**This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine®.

Citric Acid, Disodium Phosphale, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water

■ clean the affected area ■ apply a small amount

on the area 1 to 3 times daily may be covered

with sterile bandage
if bandaged let dry first

■ store a room temperature, 20-25° C (68-77° F)

Drug Facts (continued)

Directions

Other information

Inactive ingredients



A46494 04 2022



Distributed by C.D.M.A., Inc.6 43157 W. Nine Mile Novi, MI 48376-0995 www.qualitycholca.com Questions: 248-449-9500

MADE IN USA



*Compare to the active Ingredients of BETADINE®

8 fl OZ (237mL)

NDC# 63868-230-08

Povidone Iodine Solution

TOPICAL ANTISEPTIC GERMICIDE

Helps Prevent Infections in:

- Burns Cuts
- Minor Skin Wounds
- Scrapes

FOR EXTERNAL USE ONLY

8 fl oz (237 mL)

Drug Facts

Active Ingredient

Povidone-lodine, 10% (equivalent 1% titratable iodine)

.....Antimicrobial

Purpose

5

bealth-care antiseptic for preparation of the skin prior to surgery
 first aid antiseptic to help prevent infecton in minor cuts, scrapes and burns

Warnings For external use only

Do not use

- in the eyes or apply over large areas of the body
- as a first aid antiseptic longer than one week unless directed by a doctor
 on individuals who are allergic or sensitive to iodine

Stop use and ask a doctor

- . if condition persists or gets worse
- if irritation and redness develop and persist for over 72 hours

Ask a doctor in case of

· deep or puncture wounds · animal bites · serious burns

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

For preparation of the skin prior to surgery

· apply to the operative site prior to surgery

As a first aid antiseptic

- · clean the affected area
- apply a small amount of this product to the area 1-3 times daily
- may be covered with a sterile bandage
- · if bandaged, let dry first

Inactive ingredients citric acid, disodium phosphate, glycerin, nonoxynol-9, sodium hydroxide, purified water USP.







O DISTRIBUTED BY QUALITY CHOICE
NOVI, MI 48376-0995
(900) 935-2362
MADE IN U.S.A.
QC 98515
www.chaindrug.com



L-QC-PW-8 R02/07



Publix Label



Active ingredient

Purpose

Povidone-Iodine 10%....

..Antiseptic

(*Equivalent to 1% titratable iodine)

Use

First aid antiseptic to prevent infection in minor cuts and burns

Warnings For external use only.

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and consult a doctor if

- the condition persists or gets worse
- irritation and redness develop and persists for more than 72 hours

When using this product do not

- do not use in the eyes
- use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor
- do not apply over large areas of the body

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Continued >



NDC 68016-415-00 **Compare to Active Ingredient in Betadine®

Povidonelodine

10%* TOPICAL SOLUTION USP

Topical Antiseptic Microbicide



Drug Facts (continued)

Directions

■ clean the affected area ■ apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage
if bandaged let dry first

Other information

■ store a room temperature, 20-25° C (68-77° F)

Inactive ingredients

Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water

**This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine®.



Purpose

DISTRIBUTED BY: CHAIN DRUG CONSORTIUM 3301 NW BOCA RATON BLVD SUITE 101, BOCA RATON, FL 33431 MADE IN U.S.A.

Fred's Label

First Aid Antiseptic Solution

10% Povidone-lodine Topical Solution

Kills Germs And Helps Prevent The Risk Of Infection In:

- Minor Cuts
- Scrapes
- Burns



Compare to Active Ingredients In: **BETADINE®*** Povidone-Iodine 8 FL OZ (236 mL)

Drug Facts

Active ingredient Povidone- Iodine USP,

10% w/v (equivalent to 1% titratable iodine)

Use First aid antiseptic to help prevent the risk of infection in minor cuts, scrapes and burns

Warnings For external use only.

Do not

Area left blank for lot and expiration date.

- ■use in the eyes
- ■use on individuals who are allergic or sensitive to iodine
- ■apply over large areas of the body

Discontinue use and ask a doctor

- if condition persists or gets worse
- for use longer than 1 week

Ask a doctor in case of

- ■deep or puncture wounds ■ animal bites
- serious burns
- Keep out of reach of children. If

swallowed, get medical help or contact a Poison Control Center right away.

FRED'S Promise

Not Happy With This Product? Just Bring It Back For a Refund!

Distributed By: Fred's, Inc. 4300 New Getwell Rd., Memphis, TN 38118 www.fredsinc.com

Drug Facts (continued)

Directions

- Clean the affected area ■ Apply a small amount of this product to the area 1 – 3 times daily
- May be covered with a sterile bandage

 If bandaged, let dry first


Other information

Store at room temperature: 15° – 30° C (59° – 86° F)

Inactive ingredients

citric acid, disodium phosphate, glycerin, nonoxynol-9, sodium hydroxide, purified

*This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark, Betadine® R062511RLG



empty package to: discount drug mart, 211 commerce drive, medina ohio 44256

2

Rite Aid Label

8 FL OZ (237mL)






Active Ingredient Povidone-lodine, 10% (equivalent 1% titratable iodine)	Purpose Antiseption
Uses • first aid antiseptic to prevent infection in minor cuts and	
Warnings For external use only	
Ask a doctor before use if you have • deep or puncture wounds • animal bites • serious burns	
Stop use and consult a doctor if • the condition persists or gets worse • irritation and redness develop and persists for more than 72 hor	urs
When using this product do not use • in the eyes • on individuals who are allergic or sensitive to iodine or use for lunless directed by a doctor • apply over large areas of the body	onger than 1 week
Keep out of reach of children. In case of accidental ingestion, see assistance or contact a Poison Control Center immediately.	ek professional
Directions • clean the affected area • apply a small amount on the area 1 to • may be covered with sterile bandage • if bandaged, let dry first	3 times daily
Other Information • store at room temperature, 20-25°C (68-77°F)	
Inactive ingredients Citric Acid, Disodium Phosphate, Glyc Sodium Hydroxide and Purified Water	erin, Nonoxynol-9,



CVS Label



Active ingredient Purpose Povidone-lodine 10%

("Equivalent to 1% litratable iodine)

First aid antiseptic to prevent infection in minor cuts and burns

Warnings

For external use only.

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and consult a doctor if

- the condition persists or gets worse
- irritation and redness develop and persists for more than 72 hours

When using this product do not

- do not use in the eyes
 use on individuals who are allergic or sensitive to iodine or use for longer than " week unless directed by a doctor
- do not apply over large areas of the body

Keep out of reach of children. In case of accidental indestion, seek professional assistance or contact a Po son Control Center immediately. Continued >

Compares to Betadine





10%* TOPICAL SOLUTION USP SOLUCIÓN TÓPICA AL 10% * USP

FIRST AID ANTISEPTIC

 First aid antiseptic to prevent infection in minor cuts and burns.

ANTISÉPTICO PARA PRIMEROS AUXILIOS

 Antiséptico de primeros auxilios para prevenir infecciones en cortaduras y quemaduras leves.

8 FL. OZ. (237 mL) • 8 ONZAS FL (237 mL)

Drug Facts (continued)

Directions

■ clean the affected area ■ apply a small amount on the area 1 to 3 times daily may be covered with sterile bandage if bandaged let dry first

Other information

 store at room temperature, 20-25° C (68-77° F)

Inactive ingredients

Citric Acid. Disodium Phosphate, Nonoxynol-9, Glycerin. Sodium Hydroxide and Purified Water

**This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine⁸.

Distributed by/Distribuido por:

R082413.RLG

MAGE LAN Medley, Forida 33178

Questions or Comments? © 1-800-982-3435



Información sobre el

Medicamento (continuación)

Mantener fuera del alcance de los

niños. En caso de ingestión accidental,

contacte un Centro de Control de Envenenamiento inmediatamente.

■ Limpie el área afectada ■ aplique una

cantidad pequeña en el área de 1 a 3

veces al día e puede cubrirse con un

vendaje estéril, pero deje secar antes

Almacene a temperatura ambiente.

busque asistencia profesional

Información sobre el Medicamento

Ingrediente activo Povidona Yodada at 10%

Propósito Antiséptico

("Equivalente al 1% de yodo titulable)

Antiséptico de primeros auxilios para prevenir infacciones en cortaduras y quemaduras leves.

Advertencias:

Para el uso externo solamente.

Consulte un médico antes de usar si usted tiene

- heridas o punciones profundas
- mordeduras de animales
- quemaduras graves

Suspenda el uso de este producto y consulte un médico si

- la condición persiste o empecra.
- aparecen irritación y enrojecimiento y persisten por más da 72 horas.

Al utilizar este producto no

- No lo usa en los ojos
- NO use en personas alérgicas o sensibles al yodo. NO use por más de 1 semana a menos que sea ordenado por un médico
- No aplique sobre áreas extensas del cuerpo.

Continuación >



Compare con Betadine[®]

POVIDONA-YODADA

SOLUCIÓN TÓPICA AL 10% * USP

ANTISÉPTICO PARA PRIMEROS AUXILIOS

 Antiséptico de primeros auxilios para prevenir infecciones en cortaduras y quemaduras leves.



8 ONZAS FL (237 mL)

Ácido Citrico, Fosfato Glicerina, Nonoxinol-9, Hidróxido de Sodio y Agua Purificada

Este producto no es fabricado ni distribuido por Purdue Producta L.P., propietario de la marca registrada Beladine⁶

Distributed by Distributed nor

Instrucciones

Otra información

20-25 ° C (68-77 ° F)

Ingredientes inactivos

de cubrir.

MAGEJLAN Medley, Honda 33178

¿Preguntas o comentarios? 11-800-862-3435

HEB Label



Purpose

Antiseptic

(*Equivalent to 1% titratable iodine)

First aid antiseptic to prevent infection in minor cuts and burns

Warnings For external use only

Ask a doctor before use if you have

■ deep or puncture wounds ■ animal bites ■ serious burns

■ the condition persists or gets worse ■ irritation and redness develop and persists for more than 72 hours

When using this product do not

■ do not use in the eyes ■ use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor do not apply over large areas of the body

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.


■ clean the affected area ■ apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first


■ store at room temperature, 20-25° C (68-77° F)

citric acid, disodium phosphate, glycerin, nonoxynol-9, sodium hydroxide and purified water

Health Mart Label

 ∞





 clean the affected area = apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged, let


store at room temperature, 20-25° C (68-77° F)

Citric Acid, Discolum Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified

by Purdue Products L.P., owner of the registered trademark Betadine®.

A44102 08 2021





Purdue Betadine Label



Active ingredient

Purpose

Povidone-Iodine 10%.....First Aid Antiseptic (1% available iodine)

Uses

First aid to help prevent infection in minor ■ cuts ■ scrapes ■ burns

Warnings For external use only

Do not use ■ in the eyes ■ over large areas of the body ■ if you are allergic to povidone-iodine or any other ingredients in this preparation

Ask a doctor before use if you have

- deep or puncture wounds serious burns
- animal bites

Stop use and ask a doctor if ■ the condition persists or gets worse ■ you need to use this product for more than 1 week

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

■ clean the affected area ■ apply a small amount of product to the area 1 to 3 times daily may be covered with sterile bandage if bandaged let dry first

Continued >

✓QUALITY GUARANTEED

Purpose



*Safety Sealed

COMPARE TO BETADINE® ACTIVE INGREDIENT

Povidone**lodine**

10% TOPICAL SOLUTION USP FIRST AID ANTISEPTIC

- Kills Germs in Minor Cuts, Scrapes & Burns
- Used by Hospitals, **Doctors & Nurses**
- Water Soluble

*Warning: Do not use if Tamper Evident Seal is broken or missing. This product is sealed with a

Drug Facts (continued)

Other information

■ store at room temperature, 20-25° C (68-77° F)

Inactive ingredients

Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water

Questions or Comments? 1-888-423-0139

† This product is not manufactured or distributed by Purdue Products L.P., distributor of Betadine® First Aid Solution 10% Povidone-lodine Topical Solution DISTRIBUTED BY TOPCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007 © TOPCO HHGA0521 QUESTIONS? 1-888-423-0139 toncare@tonco.com_www.toncarebrand.com



Scan here for more information or call 1-888-423-0139



8 FL OZ (237 mL)

Drug Facts

Active ingredient

Povidone-lodine 10%. .First Aid Antiseptic (1% available iodine)

Uses

First aid to help prevent infection in minor ■ cuts ■ scrapes ■ burns

Warnings For external use only

Do not use ■ In the eyes ■ over large areas of the body if you are allergic to povidone-iodine or any other ingredients in this preparation

Ask a doctor before use if you have

- deep or puncture wounds serious burns
- animal bites

Stop use and ask a doctor if me the condition persists or gets worse # you need to use this product for more than 1 week

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

■ clean the affected area ■ apply a small amount of product to the area 1 to 3 times daily ■ may be covered with sterile bandage III if bandaged let dry first

Continued >

Povidonelodine

10% TOPICAL SOLUTION USP FIRST AID ANTISEPTIC

- · Kills Germs in Minor Cuts, Scrapes & Burns
- Used by Hospitals. **Doctors & Nurses**
- Water Soluble



Safety Sealed *Warning: Do not use if Tamper Evident Seal is broken or missing. This product is sealed with a foil seal under cap.

Drug Facts (continued)

Other information

 store at room temperature, 20-25° C (68-77°F)

Inactive ingredients

Citric Acid, Disodium Phosphate, Nonoxynol-9, Glycerin, Sodium Hydroxide and Purified Water

Questions or Comments? 1-888-423-0139

This product is not manufactured or distributed by Pundue Products L.P., distributor of Betadine® First Aid Solution 10% Powdone-lodine Topical Solution DISTRIBUTED BY TOPCO ASSOCIATES ILC, ELK GROVE VILLAGE, IL 60007 62TOPCO HHGAD219 OUESTIONS21-888-823-0139 topcare@topco.com www.topcarebrand.com

A46495

Scan here 04 2022 for more information

VQUALITY GUARANTEED



8 FL OZ (237 mL)

POVIDONE IODINE

povidone iodine 10% liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0395-9123
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
POVIDONE-IODINE (UNII: 85H0HZU99M) (IODINE - UNII:9679TC07X4)	IODINE	100 mg in 1 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
GLYCERIN (UNII: PDC6A3C0OX)		
NONOXYNOL-9 (UNII: 48Q180SH9T)		
SODIUM HYDROXIDE (UNII: 55X04QC32I)		
WATER (UNII: 059QF0KO0R)		

ı	Packaging				
# Item Code Package Description		Marketing Start Date	Marketing End Date		
	NDC:0395- 9123-98	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/14/2017		
	NDC:0395- 9123-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/14/2017		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	01/01/2008	

Labeler - Humco Holding Group, Inc. (825672884)

Registrant - Pharma Nobis, LLC (118564114)

Establishment			
Name	Address	ID/FEI	Business Operations
Pharma Nobis, LLC		118564114	manufacture(0395-9123), analysis(0395-9123), pack(0395-9123), label(0395-9123)

Revised: 12/2023 Humco Holding Group, Inc.