

SEQUIN 2- hydroquinone liquid
Vivier Pharma, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Seequin 2

Active Ingredient

Hydroquinone (2%)

Indication

Skin Lightening

Indication

To gradually lighten hyperpigmentation of the skin such as age spots, liver spots, freckles, or other areas of unwanted melanin hyperpigmentation that may occur in pregnancy or from the use of oral contraceptives.

Precautions

For external use only. A mild transient stinging may occur for people with sensitive skin. Do not use on broken or irritated skin. Discontinue use if irritation or rash occurs. Avoid contact with eyes and mucous membranes. In case of contact, rinse thoroughly with water. Do not use on children under 12 years of age. Keep out of reach of children. Avoid use around the eyes and lips.

Precautions

keep away from children.

Directions

Remove and discard solid white cap from bottle and replace with dropper cap. Test for skin sensitivity before using by applying 1 or 2 drops of SEEQUIN™ 2 IDS™ Liquid Gel S-erum * to an area of unbroken skin about 25 mm in diameter. If excessive irritation, rash or other reaction develops within 24 hours, discontinue use. To apply, place 1 to 2 drops of SEEQUIN™ 2 IDS™ Liquid Gel S-erum* in the palm of your hand and use fingertips to apply a thin layer to the affected areas. Use both morning and night or as directed by a physician. Depigmentation is a gradual process and results should be expected within 12 weeks of daily use. Discontinue use if no results are apparent after 8 weeks. To maintain results, use several times per week or as directed by a physician. Limit sun exposure during treatment by using protective clothing or a sunscreen protective agent to avoid re-darkening of the skin. Close dropper cap securely after each use. Store at room temperature (15-25°C / 59-77°F).

Non-Medicinal Ingredients

CITRUS GRANDIS (GRAPEFRUIT) EXTRACT, ETHOXYDIGLYCOL, FRAGRANCE/ PARFUM, L-ASCORBIC ACID USP, PROPYLENE GLYCOL, WATER/EAU.

Principal Display Panel

Seequin2_PackageLabel_PrincipalDisplayPanel

ACTIVE INGREDIENT

Hydroquinone 2% USP in a non-medical liquid gel base containing Vitamin C (L-Ascorbic Acid USP).

INGRÉDIENT ACTIF

Hydroquinone 2 % dans un gel liquide non-médical contenant de la vitamine C (Acide L-ascorbique USP).

NON-MEDICINAL

INGREDIENTS /

NON-MÉDICINAUX
CITRUS GRANDIS (GRAPEFRUIT) EXTRACT, ETHOXYDIGLYCOL, FRAGRANCE/ PARFUM, L-ASCORBIC ACID USP, PROPYLENE GLYCOL, WATER/EAU.

Vivier PHARMA
Mid. for Vivier Pharma
Hudson QC Canada J0P 1H0
Champlain NY USA 12919
vivierpharma.com

*Patent / @ / TM Vivier Canada Inc.
- used under license
IDS - IntraDermal System™ IDS™
CDA Patent: 2,396,927 U.S. Patent: 7,342,045 B2
Made in Canada

07-8908-01003

INDICATION

To gradually lighten hyperpigmentation of the skin such as age spots, liver spots, freckles or other areas of unwanted melanin hyperpigmentation that may occur in pregnancy or from the use of oral contraceptives.

DIRECTIONS

Remove and discard solid white cap from bottle and replace with dropper. To apply, place 1 to 2 drops of SEEQUIN™ 2 IDS™ (Skin Lightening) Liquid Gel Serum™ in palm of hand and use fingertips to apply a thin layer to affected areas. Use both morning and night or as directed by a physician. Gradual lightening of the discoloration can be expected in most cases. Close dropper cap securely after each use. Store at room temperature (15-25°C / 59-77°F).

PRECAUTIONS

For external use only.
Keep away from children.
See package insert for more details.

Oil-free
Alcohol-free
Dermatologist tested
Recommended by physicians

INDICATION

Pour l'éclaircissement graduel des décolorations cutanées comme des taches de son, de vieillesse, hépatiques ou la pigmentation cutanée qui peuvent se manifester durant la grossesse ou avec l'emploi des contraceptifs oraux.

MODE D'EMPLOI

Enlever et mettre au rebut le bouchon blanc de la bouteille et remplacer-le avec la pipette. Déposer 1 ou 2 gouttes de SEEQUIN™ 2 IDS™ (Éclaircissant pour la peau) Sérums gel liquide™ dans la paume de votre main et, en utilisant vos doigts, appliquer une mince couche sur la région affectée. Appliquer le matin et le soir ou selon l'avis d'un médecin. Avec une application continue on peut constater un éclaircissement graduel de la peau dans la plupart des cas. Bien fermer le bouchon après chaque usage. Entreposer à température ambiante (15 à 25 °C / 59 à 77 °F).

PRÉCAUTIONS

Pour usage externe seulement.
Garder hors de la portée des enfants. Voir l'info-feuille pour plus de détails.

Sans huile
Sans alcool
Éprouvé par les dermatologues
Recommandé par les médecins



NDC 67226-2620-3

Vivierskin®

SEEQUIN™ 2 IDS™
(HYDROQUINONE 2% USP)

LIQUID GEL SÉRUM*

SÉRUM GEL LIQUIDE*

30 mL / 1 fl oz

SEEQUIN 2

hydroquinone liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67226-2620
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROQUINONE (UNII: XV74C1N1AE) (HYDROQUINONE - UNII:XV74C1N1AE)	HYDROQUINONE	2 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
GRAPEFRUIT (UNII: O82C39RR8C)	
DIETHYLENE GLYCOL MONOETHYL ETHER (UNII: A1A118 X02B)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:67226-2620-3	1 in 1 BOX		
1		30 mL in 1 BOTTLE, DROPPER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part358A	09/01/2013	

Labeler - Vivier Pharma, Inc. (250996550)

Establishment

Name	Address	ID/FEI	Business Operations
Vivier Pharma, Inc.		250996550	manufacture(67226-2620)

Revised: 11/2013

Vivier Pharma, Inc.