

CHERRY SORE THROAT LOZENGE- benzocaine lozenge

TopCo

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Benzocaine 15 mg

Menthol 3.6 mg

Active ingredients (in each lozenge)

Benzocaine 15 mg

Menthol 3.6 mg

Purpose

Oral anesthetic

Oral anesthetic

Uses temporarily relieves these symptoms:

occasional minor irritation, pain, sore mouth and sore throat

Warnings

Methemoglobinemia warning:

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Sore throat warning: If sore throat is severe, persistent for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use

- for teething
- in children under 2 years of age

Stop use and ask doctor if

sore throat is severe or irritation, pain, or redness persists or worsens

sore mouth symptoms do not improve in 7 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact Poison Control Center right away.

Directions

adults and children 5 years and older - allow 1 lozenge to dissolve slowly in mouth. May be repeated every two hours as necessary or as directed by a doctor.

children under 5 years - do not use

Other information

store at room temperature

protect from moisture

Inactive ingredients: Corn starch, FD&C Red #40, glucose syrup, glycerin, medium chain triglycerides, natural & artificial flavors, propylene glycol, soybean oil, sucrose and water.

Questions? Or to report an adverse event call 1-888-423-0139.

TopCare
health™

Sore Throat Lozenges



18 LOZENGES

MENTHOL • BENZOCAINE • FAST ACTING • ORAL ANESTHETIC

TopCare
health™

COMPARE TO CEPACOL®
ACTIVE INGREDIENTS*

Sore Throat Lozenges

MENTHOL • BENZOCAINE
FAST ACTING • ORAL ANESTHETIC

• Fast, Soothing, Temporary Relief of
Minor Sore Throat Pain and Irritation



18 LOZENGES

CHERRY FLAVOR actual size



PN-320209
DISTRIBUTED BY TOPCO ASSOCIATES, LLC
ELK GROVE VILLAGE, IL 60007
©TOPCO B5TACH18
QUESTIONS? 1-888-423-0139
topcare@topco.com
www.topcareand.com
MADE IN USA OF U.S.
AND IMPORTED INGREDIENTS.
GLUTEN FREE
U
QUALITY GUARANTEED

Drug Facts (continued)
Inactive ingredients Corn starch, FD&C Red #40, glucose syrup, glycerin, medium chain triglycerides, natural & artificial flavors, propylene glycol, soybean oil, sucrose and water.
Questions? Or to report an adverse event call 1-888-423-0139

Drug Facts
Active ingredients (in each lozenge)
Benzocaine 15 mg
Menthol 3.6 mg
Purpose Oral anesthetic
Uses temporarily relieves these symptoms:
occasional minor irritation, pain, sore mouth and sore throat
Warnings
Methemoglobinemia warning:
Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: ■ pale, gray, or blue colored skin (cyanosis) ■ headache ■ rapid heart rate ■ shortness of breath ■ dizziness or lightheadedness ■ fatigue or lack of energy
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persists or worsens, see your dentist or doctor promptly.
Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.
Do not use ■ for teething ■ in children under 2 years of age
Stop use and ask doctor if ■ sore throat is severe or irritation, pain or redness persists or worsens ■ sore mouth symptoms do not improve in 7 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away.
Directions adults and children allow 1 lozenge to dissolve slowly in mouth. May be repeated every two hours as necessary or as directed by a doctor children under 5 years do not use
Other information ■ store at room temperature ■ protect from moisture

*This product is not manufactured or distributed by Reckitt Benckiser owner of the registered trademark Cepacol. WARNING: TAMPER-EVIDENT PACKAGING. DO NOT USE IF OUTER BOX OR BLISTER IS OPEN OR DAMAGED.



CHERRY SORE THROAT LOZENGE

benzocaine lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-832
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	3.6 mg
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	15 mg

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	17mm
Flavor	CHERRY	Imprint Code	B
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-832-18	18 in 1 BLISTER PACK; Type 0: Not a Combination Product	12/19/2017	



Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	12/19/2017	

Labeler - TopCo (006935977)

Registrant - Bestco Inc. (002149136)

Establishment

Name	Address	ID/FEI	Business Operations
Bestco Inc.		002149136	manufacture(36800-832)