SHISEIDO URBAN ENVIRONMENT OIL-FREE UV PROTECTOR- octinoxate, octocrylene, and zinc oxide lotion SHISEIDO AMERICA INC.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

SHISEIDO URBAN ENVIRONMENT OIL-FREE UV PROTECTOR CREAM

Drug Facts

ACTIVE INGREDIENTS:	Purpose
OCTINOXATE 4.9%	Sunscreen
OCTOCRYLENE 3.0%	Sunscreen
ZINC OXIDE 12.5%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 40 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- **Sun Protection Measures**. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. -2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor

Inactive Ingredients

WATER, DIMETHICONE, ISODODECANE, SD ALCOHOL 40-B, DIPROPYLENE GLYCOL,

ETHYLHEXYL PALMITATE, PEG-10 DIMETHICONE, GLYCERIN, POLYMETHYLSILSESQUIOXANE, VINYL DIMETHICONE/METHICONE SILSESQUIOXANE CROSSPOLYMER, POLYBUTYLENE GLYCOL/PPG-9/1 COPOLYMER, DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER, XYLITOL, METHYL METHACRYLATE CROSSPOLYMER, THIOTAURINE, SCUTELLARIA BAICALENSIS ROOT EXTRACT, PAEONIA ALBIFLORA ROOT EXTRACT, ONONIS SPINOSA ROOT EXTRACT, ECTOIN, LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE, HYDROGEN DIMETHICONE, BUTYLENE GLYCOL, DISODIUM EDTA, DISTEARDIMONIUM HECTORITE, ALUMINA, BHT, TRIETHOXYCAPRYLYLSILANE, SYZYGIUM JAMBOS LEAF EXTRACT, TOCOPHEROL, PHENOXYETHANOL, BENZOIC ACID, FRAGRANCE, TITANIUM DIOXIDE, MICA,

Other information

• protect this product in this container from excessive heat and direct sun.

Questions or comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICA INC. NEW YORK, N.Y. 10022
SHISEIDO DIST. NEW YORK • PARIS • MILANO

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton

SHISEIDO

GINZA TOKYO

Urban Environment
Oil-Free UV Protector

42

BROAD SPECTRUM SPF 42

For Face

WATER RESISTANT (40 MINUTES)

SUNSCREEN

30mL 1 FL. OZ.







BROAD SPECTRUM SPF 42

Drug Facts (continued)

Directions

- For sunscreen use:
- apply liberally 15 minutes before sun exposure reapply:
- after 40 minutes of swimming or sweating - immediately after towel drying - at least every 2
- hours Sun Protection
 Measures. Spending time in the
 sun increases your risk of skin
 cancer and early skin aging. To
 decrease this risk, regularly use a
 sunscreen with a broad
 spectrum SPF of 15 or higher and
 other sun protection measures
 including: I limit time in the
 sun, especially from 10 a.m. -2
 p.m. wear long-sleeve shirts,
- pants, hats, and sunglasses
 children under 6 months: Ask
 a doctor



http://s1872.com/14431AA See our website for more information.

SHISEIDO GINZA TOKYO

Urban Environment Oil-Free UV Protector

42

BROAD SPECTRUM SPF 42

For Face

WATER RESISTANT (40 MINUTES) SUNSCREEN

30mL1FL.OZ.

Drug Facts (continued)

Inactive ingredients

WATER - DIMETHICONE - ISODODECANE - SD ALCOHOL 40-B - DIPROPYLENE GLYCOL - ETHYLHEXYL PALMOTATE - PEG-10 DIMETHICONE - GLYCERIN - POLYMETHICSUSSOLIDIZANE - VINAL DIMETHICONE/WETHICONE SILSESOLIDIZANE CROSSPOLYMER - DIMETHICONE/WHYL DIMETHICONE CONSTRUYMER - XYLTDI - XYLTDI - SD ALCOHOLOR - ST ALCOHOLOR -


METHYL METHACRYLATE CROSSPOLYMER + Thiotaurine -

SCUTELLARIA BAICALENSIS ROOT EXTRACT PAEONIA ALBIRLORA ROOT EXTRACT ONONIS SPINOSA ROOT EXTRACT - ECTCIN LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL
DIMETHICONE -

HYDROGEN DIMETHICONE BUTYLENE GLYCOL - DISCOUM EDTA DISTEARCIMONIUM HECTORITE - ALLMINA BHT - TRIETHOXYCAPRYLYI SILANE SYZYGIUM JAMBOS LEAF EXTRACT TOCOPHEROL - PHENOXYETHANOL BENZOC ACID - FRAGRANCE TITANIUM DICKIDE - MICA -

Other Information

protect this product in this container from excessive heat and direct sun.



SHISEIDO URBAN ENVIRONMENT OIL-FREE UV PROTECTOR

octinoxate, octocrylene, and zinc oxide lotion

Prod	uct Ii	nform	ation
1 1 0 0	uct 11	111 1 1 1 1 1 1	auvu

Product TypeHUMAN OTC DRUGItem Code (Source)NDC:58411-346

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1557 mg in 31.77 g	
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	953 mg in 31.77 g	
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	3971 mg in 31.77 g	

Inactive Ingredients	
Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICO NE (UNII: 92RU3N3Y1O)	
ISODODECANE (UNII: A8289P68Y2)	
DIPRO PYLENE GLYCOL (UNII: E107L85C40)	
ETHYLHEXYL PALMITATE (UNII: 2865993309)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8 PR7V1SVM0)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
XYLITOL (UNII: VCQ006KQ1E)	
THIO TAURINE (UNII: NQZ2D7AO62)	
SCUTELLARIA BAICALENSIS ROOT (UNII: 7J95K7ID2S)	
PAEONIA LACTIFLORA ROOT (UNII: 3Z3866YW6P)	
ONONIS SPINOSA ROOT (UNII: FD2FMC53M1)	
ECTO INE (UNII: 7GXZ3858RY)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
EDETATE DISO DIUM (UNII: 7FLD9 1C86K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
BUTYLATED HYDRO XYTO LUENE (UNII: 1P9 D0 Z171K)	
TRIETHO XYCAPRYLYLSILANE (UNII: LDC331P08E)	
SYZYGIUM JAMBOS LEAF (UNII: 407Z4W5LFF)	
TOCOPHEROL (UNII: R0ZB2556P8)	
PHENO XYETHANOL (UNII: HIE49 2ZZ3T)	
BENZOIC ACID (UNII: 8 SKN0 B0 MIM)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	
MICA (UNII: V8A1AW0880)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-346- 10	1 in 1 CARTON	10/01/2012	
1		31.77 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:58411-346- 80	2.12 g in 1 POUCH; Type 0: Not a Combination Product	10/01/2012	
3	NDC:58411-346- 11	1 in 1 CARTON	10/01/2012	

.3	7.41 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product			
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH NOT FI	NAL part352	10/01/2012		

Labeler - SHISEIDO AMERICA INC. (193691821)

Establishment			
Name	Address	ID/FEI	Business Operations
SHISEIDO AMERICA INC.		782677132	MANUFACTURE(58411-346), ANALYSIS(58411-346)

Revised: 12/2019 SHISEIDO AMERICA INC.