

MOTION SICKNESS- dimenhydrinate tablet
Strategic Sourcing Services, LLC (Health Mart)

Health Mart 44-198 Delisted

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery
- avoid alcoholic beverages

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

Other information

- **each tablet contains:** calcium 35 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- protect from moisture
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?

1-800-426-9391

Principal display panel

Health Mart®

NDC 62011-0341-1

Compare to Dramamine® Original Formula
active ingredient*

ORIGINAL FORMULA

Motion Sickness

Dimenhydrinate 50 mg
Antiemetic

Prevents Nausea, Vomiting & Dizziness
for Children & Adults

Actual Size

12 Tablets

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR
SHOWS ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by Medtech Products
Inc., owner of the registered trademark Dramamine® Original Formula.
50844 REV0518B19802

McKESSON

Distributed by McKesson
6555 State Highway 161
Irving, TX 75039
healthmart.com



HealthMart.



NDC 62011-0341-1

Compare to Dramamine® Original Formula active ingredient*

ORIGINAL FORMULA

Motion Sickness

Dimenhydrinate 50 mg
Antiemetic

12 Tablets

Prevents Nausea, Vomiting & Dizziness
for Children & Adults



HealthMart.

NDC 62011-0341-1

Compare to Dramamine® Original Formula active ingredient*

ORIGINAL FORMULA

Motion Sickness

Dimenhydrinate 50 mg
Antiemetic

Actual Size
12 Tablets

Prevents Nausea, Vomiting & Dizziness
for Children & Adults

LEBG749M

MCKESSON
Distributed by McKesson Corporation
6555 State Highway 161
Irving, TX 75039
www.healthmart.com


*This product is not manufactured or distributed by Mediatech Products Inc., owner of the registered trademark Dramamine® Original Formula.
50844 REV0518818902

8-1241-196-02-R
REV0518818902



TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR
SHOWS ANY SIGNS OF TAMPERING

Drug Facts	Active ingredient (in each tablet) Purpose Dimenhydrinate 50 mg. Antiemetic
Uses Associated with motion sickness: nausea ■ vomiting ■ dizziness	Warnings Do not use for children under 2 years of age unless directed by a doctor. ■ Ask a doctor before use if you have ■ breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. ■ When using this product ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ marked drowsiness may occur ■ use caution when driving a motor vehicle or operating machinery ■ avoid alcoholic beverages ■ If pregnant or breast-feeding, ask a health professional before use. ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions ■ To prevent motion sickness, the first dose should be taken one-half to one hour before starting activity. adults and children 1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor. children 6 to under 12 years 1/2 to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor. children 2 to under 6 years 1/2 tablet every 6-8 hours; do not exceed 1 1/2 tablets in 24 hours, or as directed by a doctor.	Other information ■ each tablet contains: calcium 35 mg ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture ■ see end flap for expiration date and lot number Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid Questions or comments? 1-800-426-9391



KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION



Health mart 44-198

MOTION SICKNESS

dimenhydrinate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62011-0341
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, 8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;198
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0341-1	2 in 1 CARTON	12/01/1992	09/15/2024
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	12/01/1992	09/15/2024

Labeler - Strategic Sourcing Services, LLC (Health Mart) (116956644)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(62011-0341)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(62011-0341) , pack(62011-0341)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(62011-0341)

Revised: 10/2023

Strategic Sourcing Services, LLC (Health Mart)