

## **HEADACHE RELIEF EXTRA STRENGTH- acetaminophen, aspirin, caffeine tablet**

### **TARGET Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

## **Drug Facts**

### **Active ingredients (in each geltab)**

**Acetaminophen 250 mg**

**Aspirin 250 mg (NSAID)\***

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

## **Purposes**

**Pain reliever**

**Pain reliever**

Pain reliever aid

## **Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - a cold
  - arthritis
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps

## **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

### **Stop use and ask doctor if**

- an allergic reaction occurs. Seek medical help right away.

- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red and swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 gels every 6 hours; do not take more than 8 gels in 24 hours
- children under 12 years: do not use unless directed by a doctor

### **Other information**

- store between 20°-25°C (68°-77°F)

### **Inactive ingredients**

black printing ink, corn starch, D&C yellow #10 aluminum lake, edetate disodium, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hydroxypropyl cellulose, hydroxypropyl cellulose-low substituted, hypromellose, iron oxide black, maltitol, povidone, purified water, stearic acid, talc, titanium dioxide

### **Questions?**

Call **1-800-910-6874**

### **Principal Display Panel**

Compare to active ingredients in Excedrin® Extra Strength†  
extra strength

**headache relief**

acetaminophen, 250 mg pain reliever

aspirin (NSAID), 250 mg pain reliever

caffeine, 65 mg pain reliever aid

GELTABS

†This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare Holdings (US) LLC, distributor of Excedrin® Extra Strength.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS  
BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT  
INFORMATION.**

Distributed by Target Corporation

Minneapolis, MN 55403

**Package Label**

Exp. Date:

Lot No.:

PLD-DS14C F0008571

3 59726 58680 1

094 01 0434 R03 C-000703-01-041  
Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2023 Target Brands, Inc.

## Drug Facts

**Active ingredients (in each geltab)** **Purposes**  
Acetaminophen 250 mg.....Pain reliever  
Aspirin 250 mg (NSAID)\*.....Pain reliever  
Caffeine 65 mg.....Pain reliever aid  
\*nonsteroidal anti-inflammatory drug

## Uses

- temporarily relieves minor aches and pains due to
  - headache
  - a cold
  - arthritis
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions.

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Liver warnings:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks every day while using this product
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains

Compare to active ingredients in  
Excedrin® Extra Strength†

NDC 11673-686-80

# extra strength headache relief

acetaminophen, 250 mg pain reliever  
aspirin (NSAID), 250 mg pain reliever  
caffeine, 65 mg pain reliever aid



80 GELTABS



ACTUAL SIZE

80

GELTABS

## Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 geltabs every 6 hours; do not take more than 8 geltabs in 24 hours
- children under 12 years: do not use unless directed by a doctor

## Other information

- store between 20°-25°C (68°-77°F)

**Inactive ingredients** black printing ink, corn starch, D&C yellow #10 aluminum lake, edictate disodium, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hydroxypropyl cellulose, hydroxypropyl cellulose-low substituted, hypromellose, iron oxide black, maltitol, povidone, purified water, stearic acid, talc, titanium dioxide

## Questions? Call 1-800-910-6874

This product is not manufactured or distributed by  
GlaxoSmithKline Consumer Healthcare Holdings  
(US) LLC, distributor of Excedrin® Extra Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY  
SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUT OF REACH OF CHILDREN.  
KEEP OUTEN GARTING FOR COMPLETE  
QUANTITIES AND PRODUCT INFORMATION.

## TARGET Extra Strength Headache Relief

### HEADACHE RELIEF EXTRA STRENGTH

acetaminophen, aspirin, caffeine tablet

#### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-686
Route of Administration	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

#### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 9XZ8H6N6OH)	
<b>FERROSOFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>MALTITOL</b> (UNII: D65DG142WK)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)	
<b>HYDROXYPROPYL CELLULOSE, LOW SUBSTITUTED</b> (UNII: 2165RE0K14)	

#### Product Characteristics

Color	white (green)	Score	no score
Shape	CAPSULE (Round shaped)	Size	13mm
Flavor		Imprint Code	P
Contains			

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-686-20	1 in 1 BOX	01/31/2018	
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:11673-686-80	1 in 1 BOX	01/31/2018	
2		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final		part343	01/31/2018	

**Labeler** - TARGET Corporation (006961700)

Revised: 7/2023

TARGET Corporation