

**EXTRA PAIN RELIEVER- acetaminophen, aspirin and caffeine tablet, film coated**  
**Geri-Care Pharmaceutical Corp**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**GC 226**

**Active ingredients (in each caplet)**

ACETAMINOPHEN 250 MG

ASPIRIN 250 MG (nsaid\*)

CAFFEINE 65 MG

\*nonsteroidal anti-inflammatory drug

**Purposes**

PAIN RELIEVER - PAIN RELIEVER AID

**USES**

- temporarily relieves minor aches and pains

**WARNINGS**

**REYE'S SYNDROME:**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**ALLERGY ALERT**

Aspirin may cause a severe allergic reaction which may include:

hives

facial swelling

asthma (sheezing)

shock

**LIVER WARNING:**

This product contains acetaminophen. Severe liver damage may occur if you take more than 8 tablets in 24 hours, which is the maximum daily amount;

with other drugs containing acetaminophen;

3 or more alcoholic drinks every day while using this product

**STOMACH BLEEDING WARNING:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

are age 60 or older;

have had stomach ulcers or bleeding problems

take a blood thinning (anticoagulant) or steroid drug

take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

take more or for a longer time than directed

### **CAFFEINE WARNING:**

The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine containing medications, foods or beverages which while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness and occasionally, rapid heart beat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are taking**

- any other drug containing an NSAID (prescription or nonprescription)
- a blood thinning (anticoagulant) or steroid drug
- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - ringing in the ears or loss of hearing occurs
  - painful area is red or swollen
  - pain gets worse or lasts for more than 10 days
  - fever gets worse or lasts for more than 3 days
  - any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

Overdose Warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed (see overdose warning)
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12: ask a doctor

**INACTIVE INGREDIENTS**

carnauba wax, cellulose, hypromellose, mineral oil, povidone, silica, sodium starch glycolate, starch, stearic acid, titanium dioxide

**Other information**

- store at room temperature 15°- 25°C (59°- 77°F)

**Package label**



NDC 57896-226-01

THE SENSIBLE CHOICE  
FOR PRICE AND HEALTH

**Extra  
Strength**

# Pain-Reliever

Acetaminophen 250 mg  
Aspirin 250 mg (NSAID)  
Caffeine 65 mg

COMPARE TO THE ACTIVE  
INGREDIENTS IN EXCEDRIN®\*

100 Caplets

**Tamper Evident:** Do not use if imprinted seal under cap is missing or broken.

## Drug Facts

### Active ingredients (in each caplet)

Acetaminophen 250 mg ..... Pain reliever  
Aspirin 250 mg (NSAID)\* ..... Pain reliever  
Caffeine 65 mg ..... Pain reliever aid  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ Temporarily relieves minor aches and pains

### Warnings

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than directed ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

\*This product is not manufactured or distributed by the owner of the registered trademark Excedrin®.

Distributed By: Gert-Care Pharmaceuticals Corp.  
1650 63rd Street, Brooklyn, NY 11204

REV0715



3 57896 22601 5

PEEL HERE FOR MORE DRUG FACTS

## Drug Facts (continued)

**Do not use** ■ If you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor or pharmacist before use if** ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

**Ask a doctor or pharmacist before use if you are taking** ■ any other drug containing an NSAID (prescription or nonprescription) ■ a blood thinner (anticoagulant) or steroid drug ■ a prescription drug for diabetes, gout, or arthritis ■ any other drug, or are under a doctor's care for any serious condition

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding:  feel faint  vomit blood  have bloody or black stools

■ you experience any of the following signs of stomach bleeding:  feel faint  vomit blood  have bloody or black stools

■ tingling in the ears or loss of hearing occurs  
■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**  
**Overdose Warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ do not use more than directed (see overdose warning)  
■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours  
■ children under 12: ask a doctor

**Other information**  
■ store at room temperature 15° - 25°C (59° - 77°F)  
**Inactive ingredients** camellia wax, cellulose, hypromellose, mineral oil, povidone, silica, sodium starch glycolate, starch, stearic acid, titanium dioxide.

STOP PEELING

## EXTRA PAIN RELIEVER

acetaminophen, aspirin and caffeine tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-226
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg	
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg	

Inactive Ingredients			
Ingredient Name	Strength		
STEARIC ACID (UNII: 4ELV7Z65AP)			
CARNAUBA WAX (UNII: R12CBM0EIZ)			

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (CAPSULE SHAPED TABLET)	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL370
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-226-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2015	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	07/01/2015	

**Labeler** - Geri-Care Pharmaceutical Corp (611196254)

**Registrant** - Geri-Care Pharmaceutical Corp (611196254)

### Establishment

Name	Address	ID/FEI	Business Operations
Geri-Care Pharmaceutical Corp		611196254	repack(57896-226)

Revised: 12/2017

Geri-Care Pharmaceutical Corp