

CYANOCOBALAMIN- cyanocobalamin injection, solution
Somerset Therapeutics, LLC

CYANOCOBALAMIN INJECTION, USP
Rx only

DESCRIPTION


Cyanocobalamin Injection, USP is a sterile solution of cyanocobalamin for intramuscular or subcutaneous injection. Each mL contains 1000 mcg cyanocobalamin.

Each vial also contains Sodium Chloride, 0.9%. Benzyl Alcohol, 1.5%, is present as a preservative. Hydrochloric acid and/or sodium hydroxide may have been added during manufacture to adjust the pH (range 4.5-7.0).

Cyanocobalamin appears as red crystalline or small crystals. It is very hygroscopic in the anhydrous form, and sparingly soluble in water (1:80). It is stable to autoclaving for short periods at 121°C. The vitamin B₁₂ coenzymes are very unstable in light.

The chemical name is 5,6-dimethyl-benzimidazolyl cyanocobamide; the molecular formula is C₆₃H₈₈CoN₁₄O₁₄P. The cobalt content is 4.34%. The molecular weight is 1355.39.

The structural formula is represented below.



CLINICAL PHARMACOLOGY

Vitamin B₁₂ is essential to growth, cell reproduction, hematopoiesis, and nucleoprotein and myelin synthesis.

Cyanocobalamin is quantitatively and rapidly absorbed from intramuscular and subcutaneous sites of injection; the plasma level of the compound reaches its peak within 1 hour after intramuscular injection. Absorbed vitamin B₁₂ is transported via specific B₁₂ binding proteins, transcobalamin I and II to the various tissues. The liver is the main organ for vitamin B₁₂ storage.

Within 48 hours after injection of 100 or 1000 mcg of vitamin B₁₂, 50 to 98% of the injected dose may appear in the urine. The major portion is excreted within the first eight hours. Intravenous administration results in even more rapid excretion with little opportunity for liver storage.

Gastrointestinal absorption of vitamin B₁₂ depends on the presence of sufficient intrinsic factor and calcium ions. Intrinsic factor deficiency causes pernicious anemia, which may be associated with subacute combined degeneration of the spinal cord. Prompt parenteral administration of vitamin B₁₂ prevents progression of neurologic damage.

The average diet supplies about 5 to 15 mcg/day of vitamin B₁₂ in a protein-bound form that is available for absorption after normal digestion. Vitamin B₁₂ is not present in foods of plant origin, but is abundant in foods of animal origin. In people with normal absorption, deficiencies have been reported only in strict vegetarians who consume no products of animal origin (including no milk products or eggs).

Vitamin B₁₂ is bound to intrinsic factor during transit through the stomach; separation occurs in the terminal ileum in the presence of calcium, and vitamin B₁₂ enters the mucosal cell for absorption. It is then transported by the transcobalamin binding proteins. A small amount (approximately 1% of the total amount ingested) is absorbed by simple diffusion, but this mechanism is adequate only with very large doses. Oral absorption is considered too undependable to rely on in patients with pernicious anemia or other conditions resulting in malabsorption of vitamin B₁₂.

Cyanocobalamin is the most widely used form of vitamin B₁₂, and has hematopoietic activity apparently identical to that of the antianemia factor in purified liver extract. Hydroxycobalamin is equally as effective as cyanocobalamin, and they share the cobalamin molecular structure.

INDICATIONS AND USAGE

Cyanocobalamin is indicated for vitamin B₁₂ deficiencies due to malabsorption which may be associated with the following conditions:

It may be possible to treat the underlying disease by surgical correction of anatomic lesions leading to small bowel bacterial overgrowth, expulsion of fish tapeworm, discontinuation of drugs leading to vitamin malabsorption (see **Drug Interactions**), use of a gluten-free diet in nontropical sprue, or administration of antibiotics in tropical sprue. Such measures remove the need for long-term administration of cyanocobalamin.

Requirements of vitamin B₁₂ in excess of normal (due to pregnancy, thyrotoxicosis, hemolytic anemia, hemorrhage, malignancy, hepatic and renal disease) can usually be met with oral supplementation.

Cyanocobalamin Injection, USP is also suitable for the vitamin B₁₂ absorption test (Schilling test).

CONTRAINDICATIONS

Sensitivity to cobalt and/or vitamin B₁₂ is a contraindication.

WARNINGS:

Patients with early Leber's disease (hereditary optic nerve atrophy) who were treated with cyanocobalamin suffered severe and swift optic atrophy.

Hypokalemia and sudden death may occur in severe megaloblastic anemia which is treated intensely.

Anaphylactic shock and death have been reported after parenteral vitamin B₁₂ administration. An intradermal test dose is recommended before Cyanocobalamin Injection, USP is administered to patients suspected of being sensitive to this drug.

This product contains Benzyl Alcohol. Benzyl Alcohol has been reported to be associated with a fatal "Gasping Syndrome" in premature infants.

This product contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired.

Premature neonates are particularly at risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.

Research indicates that patients with impaired kidney function, including premature neonates, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

PRECAUTIONS

General Precautions: Vitamin B₁₂ deficiency that is allowed to progress for longer than 3 months may produce permanent degenerative lesions of the spinal cord. Doses of folic acid greater than 0.1 mg per day may result in hematologic remission in patients with vitamin B₁₂ deficiency. Neurologic manifestations will not be prevented with folic acid, and if not treated with vitamin B₁₂, irreversible damage will result.

Doses of cyanocobalamin exceeding 10 mcg daily may produce hematologic response in patients with folate deficiency. Indiscriminate administration may mask the true diagnosis.

Information for Patients: Patients with pernicious anemia should be informed that they will require monthly injections of vitamin B₁₂ for the remainder of their lives. Failure to do so will result in return of the anemia and in development of incapacitating and irreversible damage to the nerves of the spinal cord. Also, patients should be warned about the danger of taking folic acid in place of vitamin B₁₂, because the former may prevent anemia but allow progression of subacute combined degeneration.

A vegetarian diet which contains no animal products (including milk products or eggs) does not supply any vitamin B₁₂. Patients following such a diet, should be advised to take oral vitamin B₁₂ regularly. The need for vitamin B₁₂ is increased by pregnancy and lactation. Deficiency has been recognized in infants of vegetarian mothers who were breast fed, even though the mothers had no symptoms of deficiency at the time.

Contains no more than 57 mcg/L of aluminum.

Laboratory Tests: During the initial treatment of patients with pernicious anemia, serum potassium must be observed closely the first 48 hours and potassium replaced if necessary.

Hematocrit, reticulocyte count, vitamin B₁₂, folate and iron levels should be obtained prior to treatment. Hematocrit and reticulocyte counts should be repeated daily from the fifth to seventh days of therapy and then frequently until the hematocrit is normal. If folate levels are low, folic acid should also be administered. If reticulocytes have not increased after treatment or if reticulocyte counts do not continue at least twice normal as long as the hematocrit is less than 35%, diagnosis or treatment should be reevaluated. Repeat determinations of iron and folic acid may reveal a complicating illness that might inhibit the response of the marrow.

Patients with pernicious anemia have about 3 times the incidence of carcinoma of the stomach as the general population, so appropriate tests for this condition should be carried out when indicated.

Drug/Laboratory Test Interactions: Persons taking most antibiotics, methotrexate and pyrimethamine invalidate folic acid and vitamin B₁₂ diagnostic blood assays.

Colchicine para-aminosalicylic acid and heavy alcohol intake for longer than 2 weeks may produce malabsorption of vitamin B₁₂.

Carcinogenesis, Mutagenesis, Impairment of Fertility: Long term studies in animals to evaluate carcinogenic potential have not been done. There is no evidence from long-term use in patients with pernicious anemia that cyanocobalamin is carcinogenic. Pernicious anemia is associated with an increased incidence of carcinoma of the stomach, but this is believed to be related to the underlying pathology and not to treatment with cyanocobalamin.

Pregnancy: Teratogenic Effects. Pregnancy Category C: Adequate and well-controlled studies have not been done in pregnant women. However, vitamin B₁₂ is an essential vitamin and requirements are increased during pregnancy. Amounts of vitamin B₁₂ that are recommended by the Food and Nutrition Board, National Academy of Science-National Research Council for pregnant women (4 mcg daily) should be consumed during pregnancy.

Nursing Mothers: Vitamin B₁₂ is known to be excreted in human milk. Amounts of vitamin B₁₂ that are recommended by the Food and Nutrition Board, National Academy of Science-National Research

Council for lactating women (4 mcg daily) should be consumed during lactation.

Pediatric Use: Intake in children should be in the amount (0.5 to 3 mcg daily) recommended by the Food and Nutrition Board, National Academy of Science-National Research Council.

ADVERSE REACTIONS

Generalized: Anaphylactic shock and death have been reported with administration of parenteral vitamin B₁₂ (see **WARNINGS**).

Cardiovascular: Pulmonary edema and congestive heart failure early in treatment; peripheral vascular thrombosis.

Hematological: Polycythemia vera

Gastrointestinal: Mild transient diarrhea

Dermatological: Itching; transitory exanthema

Miscellaneous: Feeling of swelling of entire body

OVERDOSAGE

No overdosage has been reported with this drug.

DOSAGE AND ADMINISTRATION

Avoid using the intravenous route. Use of this product intravenously will result in almost all of the vitamin being lost in the urine.

Pernicious Anemia: Parenteral vitamin B₁₂ is the recommended treatment and will be required for the remainder of the patient's life. The oral form is not dependable. A dose of 100 mcg daily for 6 or 7 days should be administered by intramuscular or deep subcutaneous injection. If there is clinical improvement and if a reticulocyte response is observed, the same amount may be given on alternate days for seven doses, then every 3 to 4 days for another 2 to 3 weeks. By this time hematologic values should have become normal. This regimen should be followed by 100 mcg monthly for life. Folic acid should be administered concomitantly if needed.

Patients with Normal Intestinal Absorption: Where the oral route is not deemed adequate, initial treatment similar to that for patients with pernicious anemia may be indicated depending on the severity of the deficiency. Chronic treatment should be with an oral B₁₂ preparation. If other vitamin deficiencies are present, they should be treated.

Schilling Test: The flushing dose is 1000 mcg.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

HOW SUPPLIED

Cyanocobalamin Injection, USP 1000 mcg/mL

NDC Number	Cyanocobalamin mcg/mL	Volume
70069- 171 -05	1,000	30 mL
70069- 171 -10		

30 mL vials are multiple dose vial, packaged as;

- 5 vials per tray and
- 10 vials per tray.

Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) [See USP Controlled Room Temperature].

PROTECT THE PRODUCT FROM LIGHT.

For Product Inquiry call 1-800-417-9175.

Issued: October, 2019

Manufactured for:

Somerset Therapeutics, LLC

Hollywood, FL 33024

Made in India

Code No.:KR/DRUGS/KTK/28/289/97

ST-CYC13/P/01

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Container label

<p>NDC 70069-171-01 Rx only</p>	<p>Each mL contains: Cyanocobalamin 1,000 mcg, Benzyl Alcohol 1.5%, Sodium Chloride 0.9%, Water for Injection q.s. pH adjusted with Hydrochloric Acid and/or Sodium Hydroxide.</p>	<p>ST-CYC13/L/01</p>
<p>Cyanocobalamin Injection, USP</p>	<p>Contains no more than 625 mcg/L of aluminum.</p>	
<p>30,000 mcg/30 mL</p>	<p>WARNING: PROTECT FROM LIGHT. Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature].</p>	
<p>(1,000 mcg/mL)</p>	<p>Usual Dosage: See package insert.</p>	<p>No Varnish Area 38 x 12mm</p>
<p>For IM or SC Use Only</p>	<p>Manufactured for: Somerset Therapeutics, LLC Hollywood, FL 33024</p>	
<p>30 mL Multiple Dose Vial</p>	<p>Made in India Code No.:KR/DRUGS/KTK/28/289/97</p>	

Carton label – 5 x 30 mL

<p>Each mL contains: Cyanocobalamin 1,000 mcg, Benzyl Alcohol 1.5%, Sodium Chloride 0.9%, Water for Injection q.s. pH (range 4.5-7.0) adjusted with Hydrochloric Acid and/or Sodium Hydroxide.</p> <p>Contains no more than 625 mcg/L of aluminum.</p> <p>WARNING: PROTECT FROM LIGHT. Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) [See USP Controlled Room Temperature].</p> <p>Usual Dosage: See package insert.</p> <p>Manufactured for: Somerset Therapeutics, LLC Hollywood, FL 33024 Customer Care # 1-800-417-9175</p> <p>Made in India Code No.:KR/DRUGS/KTK/28/289/97 ST-CYC13/C5/00</p>	<p>Rx only</p> <p style="text-align: right;">NDC 70069-171-05 5 x 30 mL Multiple Dose Vials</p> <h2 style="margin: 0;">Cyanocobalamin Injection, USP</h2> <div style="background-color: #00a651; color: white; padding: 5px; display: inline-block; margin: 5px 0;">30,000 mcg/30 mL</div> <p>(1,000 mcg/mL)</p> <p>For Intramuscular or Subcutaneous Use Only</p>
---	--

Carton label – 10 x 30 mL

<p>Each mL contains: Cyanocobalamin 1,000 mcg, Benzyl Alcohol 1.5%, Sodium Chloride 0.9%, Water for Injection q.s. pH (range 4.5-7.0) adjusted with Hydrochloric Acid and/or Sodium Hydroxide.</p> <p>Contains no more than 625 mcg/L of aluminum.</p> <p>WARNING: PROTECT FROM LIGHT. Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) [See USP Controlled Room Temperature].</p> <p>Usual Dosage: See package insert.</p>	<p>Manufactured for: Somerset Therapeutics, LLC Hollywood, FL 33024 Customer Care # 1-800-417-9175</p> <p>Made in India Code No.:KR/DRUGS/KTK/28/289/97 ST-CYC13/C10/01</p>	<p>Rx only</p> <p style="text-align: right;">NDC 70069-171-10 10 x 30 mL Multiple Dose Vials</p> <h2 style="margin: 0;">Cyanocobalamin Injection, USP</h2> <div style="background-color: #00a651; color: white; padding: 5px; display: inline-block; margin: 5px 0;">30,000 mcg/30 mL</div> <p>(1,000 mcg/mL)</p> <p>For Intramuscular or Subcutaneous Use Only</p>	<p style="font-size: small;">3 70069 17110 3</p>
---	--	---	--

<h2 style="margin: 0;">CYANOCOBALAMIN</h2> <p style="margin: 0;">cyanocobalamin injection, solution</p>			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70069-171
Route of Administration	INTRAMUSCULAR, SUBCUTANEOUS		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
CYANOCOBALAMIN (UNII: P6 YC3EG204) (CYANOCOBALAMIN - UNII:P6 YC3EG204)	CYANOCOBALAMIN	1000 ug in 1 mL	
Inactive Ingredients			
Ingredient Name	Strength		
BENZYL ALCOHOL (UNII: LKG8494WBH)	15 mg in 1 mL		
SODIUM CHLORIDE (UNII: 451W47IQ8X)	9 mg in 1 mL		
HYDROCHLORIC ACID (UNII: QTT17582CB)			
SODIUM HYDROXIDE (UNII: 55X04QC32I)			
WATER (UNII: 059QF0K00R)			
Product Characteristics			
Color	RED (Red Colored Clear Solution)	Score	
Shape		Size	
Flavor		Imprint Code	

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70069-171-05	5 in 1 CARTON	12/18/2018	
1	NDC:70069-171-01	30 mL in 1 VIAL; Type 0: Not a Combination Product		
2	NDC:70069-171-10	10 in 1 CARTON	12/18/2018	
2	NDC:70069-171-01	30 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209429	12/18/2018	

Labeler - Somerset Therapeutics, LLC (079947873)**Registrant** - Somerset Therapeutics, LLC (079947873)**Establishment**

Name	Address	ID/FEI	Business Operations
Wintac Limited		677236695	ANALYSIS(70069-171) , LABEL(70069-171) , MANUFACTURE(70069-171) , PACK(70069-171)

Revised: 3/2020

Somerset Therapeutics, LLC