

MAXI-TUSS JR- dextromethorphan hydrobromide and phenylephrine hydrochloride liquid
MCR American Pharmaceuticals, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Maxi-Tuss Jr

Drug Facts

<i>Active Ingredients (in each 5 mL teaspoonful)</i>	<i>Purpose</i>
Dextromethorphan HBr 5 mg	Cough Suppressant
Phenylephrine HCl 2.5 mg	Nasal Decongestant

Uses

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other respiratory allergies:

- cough due to minor throat and bronchial irritation as may occur with a cold
- nasal congestion due to a cold, hay fever or other upper respiratory allergies

Warnings

Do not exceed recommended dosage.

Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts or as occurs with asthma
- difficulty in urination due to enlargement of the prostate gland

- A persistent cough may be a sign of a serious condition. If cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache, consult a doctor.
- Do not take this product for persistent or chronic cough such as occurs with smoking, asthma, or emphysema, or if cough is accompanied by excessive phlegm (mucus) unless directed by a doctor.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

Directions**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	2 teaspoonfuls (10 mL) every 4 hours, not to exceed 24 teaspoonfuls in 24 hours or as directed by a doctor
Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 4 hours, not to exceed 12 teaspoonfuls in 24 hours or as directed by a doctor
Children under 6 years of age:	Consult a physician

Other information

Store at 59°-86°F (15°-30°C) [see USP Controlled Room Temperature]

Inactive ingredients

Citric acid, grape flavor, methylparaben, monoammonium glycyrrhizinate, potassium citrate, propylparaben, propylene glycol, purified water, sorbitol, sucralose

Questions or comments?

Call 352.754.8587

PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label

NDC 58605-308-16

Maxi-Tuss Jr

Cough Suppressant ■ Nasal Decongestant

Sugar Free ■ Alcohol Free ■ Dye Free

Each teaspoonful (5 mL) for oral administration contains:

Dextromethorphan HBr 5 mg

Phenylephrine HCl 2.5 mg

Grape Flavor

Tamper evident by foil seal under cap.

Do not use if foil seal is broken or missing.

Manufactured for:

MCR American Pharmaceuticals, Inc.

Brooksville, FL 34604

16 fl oz (473 mL)

NDC 58605-308-16

Maxi-Tuss Jr

Cough Suppressant ■ Nasal Decongestant

Sugar Free ■ Alcohol Free ■ Dye Free

Each teaspoonful (5 mL) for oral administration contains:
 Dextromethorphan HBr 5 mg
 Phenylephrine HCl 2.5 mg

Grape Flavor

Tamper evident by foil seal under cap.
 Do not use if foil seal is broken or missing.

Manufactured for:
 MCR American Pharmaceuticals, Inc.
 Brooksville, FL 34604



16 fl oz (473 mL)

Lot:
Exp. Date:

Drug Facts

Active Ingredients (in each 5 mL teaspoonful)

Dextromethorphan HBr 5 mg	Cough Suppressant
Phenylephrine HCl 2.5 mg	Nasal Decongestant

Uses temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other respiratory allergies:
 ■ cough due to minor throat and bronchial irritation as may occur with a cold ■ nasal congestion due to a cold, hay fever or other upper respiratory allergies

Warnings
Do not exceed recommended dosage.

Do not use this product
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if you have
 ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ cough that occurs with too much phlegm (mucus) ■ chronic cough that lasts or as occurs with asthma ■ difficulty in urination due to enlargement of the prostate gland

■ A persistent cough may be a sign of a serious condition. If cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache, consult a doctor.
 ■ Do not take this product for persistent or chronic cough such as occurs with smoking, asthma, or emphysema, or if cough is accompanied by excessive phlegm (mucus) unless directed by a doctor.

Drug Facts (continued)

Stop use and ask a doctor if
 ■ nervousness, dizziness, or sleeplessness occur
 ■ symptoms do not improve within 7 days or are accompanied by fever
 ■ new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
 In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

Directions

Do not exceed recommended dosage.

Adults and children 12 years of age and over:	2 teaspoonfuls (10 mL) every 4 hours, not to exceed 24 teaspoonfuls in 24 hours or as directed by a doctor
---	--

Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 4 hours, not to exceed 12 teaspoonfuls in 24 hours or as directed by a doctor
--------------------------------------	--

Children under 6 years of age:	Consult a physician
--------------------------------	---------------------

Other information

Store at 59°-86°F (15°-30°C) [see USP Controlled Room Temperature]

Inactive ingredients

Citric acid, grape flavor, methylparaben, monoammonium glycyrrhizinate, potassium citrate, propylparaben, propylene glycol, purified water, sorbitol, sucralose

Questions or comments?

Call 352.754.8587

Rev. 05/20

MAXI-TUSS JR

dextromethorphan hydrobromide and phenylephrine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58605-308
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Dextromethorphan Hydrobromide (UNII: 9D2RT9KYH) (Dextromethorphan - UNII:7355X3ROTS)	Dextromethorphan Hydrobromide	5 mg in 5 mL
Phenylephrine Hydrochloride (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine Hydrochloride	2.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
Methylparaben (UNII: A2I8C7HI9T)	
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
Potassium Citrate (UNII: EE90ONI6FF)	
Propylparaben (UNII: Z8IX2SC1OH)	
Propylene Glycol (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
Sorbitol (UNII: 506T60A25R)	

Sucralose (UNII: 96K6UQ3ZD4)

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58605-308-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	08/01/2020	

Labeler - MCR American Pharmaceuticals, Inc. (783383011)

Establishment

Name	Address	ID/FEI	Business Operations
MCR American Pharmaceuticals, Inc.		783383011	MANUFACTURE(58605-308)

Revised: 6/2020

MCR American Pharmaceuticals, Inc.