

**SALICYLIC ACID- salicylic acid liquid**  
**Rochester Pharmaceuticals**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).*

-----


**DESCRIPTION**

Rx Only

Salicylic Acid, 27.5% Wart Remover

FOR TOPICAL USE ONLY. NOT FOR OPHTHALMIC, ORAL OR INTRA-VAGINAL USE.

Salicylic Acid, 27.5% Wart Remover is a topical preparation containing 27.5% salicylic acid in a film-forming vehicle composed of acrylates copolymer, isopropyl alcohol, isobutyrate, n-butyl acetate, and polyvinyl butyral. The pharmacologic activity of this medication is generally attributed to the keratolytic activity of salicylic acid which is incorporated into a film-forming vehicle designed to cover the wart without the need for a bandage. The chemical structural formula of salicylic acid is:



**CLINICAL PHARMACOLOGY**

Although the exact mode of action of salicylic acid in the treatment of warts is not known, its activity appears to be associated with its keratolytic action which results in mechanical removal of epidermal cells infected with wart viruses.

**INDICATIONS & USAGES**

Salicylic Acid, 27.5% Wart Remover is indicated for the topical treatment and removal of common warts and plantar warts.

**CONTRAINDICATIONS**

Salicylic Acid, 27.5% Wart Remover should not be used by diabetics or patients with impaired blood circulation. Salicylic Acid, 27.5% Wart Remover should not be used on moles, birthmarks, unusual warts with hair growing from them or warts on the face.

**PRECAUTIONS**

Salicylic Acid, 27.5% Wart Remover is for external use only. Do not permit Salicylic Acid, 27.5% Wart Remover to contact eyes or mucous membranes. If contact with eyes or mucous membranes occurs, immediately flush with water for 15 minutes. Salicylic Acid, 27.5% Wart Remover should not be allowed to contact normal skin surrounding the wart. Treatment should be discontinued if excessive

irritation occurs. Salicylic Acid, 27.5% Wart Remover is flammable and should be kept away from fire or flame. Keep bottle tightly capped when not in use.

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

## **ADVERSE REACTIONS**

A localized irritant reaction may occur if Salicylic Acid, 27.5% Wart Remover is applied to the normal skin surrounding the wart. Any irritation may normally be controlled by temporarily discontinuing use and by applying the medication only to the wart site when treatment is resumed.

## **DOSAGE & ADMINISTRATION**

Prior to application of Salicylic Acid, 27.5% Wart Remover, soak wart in warm water for five minutes. Remove any loosened tissue by gently rubbing with a brush, wash cloth, emery board or pumice stone. Dry thoroughly. Using the brush applicator supplied, apply twice to entire wart surface, allowing the first application to dry before applying the second. Treatment should be once or twice a day and should continue as directed by physician. Be careful not to apply to surrounding skin. Clinically visible improvement will normally occur during the first or second week of therapy. Maximum resolution may be expected after four to six weeks of drug use.

## **HOW SUPPLIED**

Salicylic Acid, 27.5% Wart Remover is supplied in 10 mL amber glass bottles with a brush applicator - NDC 49908-237-10

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Brief exposure to temperatures up to 40°C (104°F) may be tolerated provided the mean kinetic temperature does not exceed 25°C (77°F); however, such exposure should be minimized. Protect from freezing.

Manufactured for: Rochester Pharmaceuticals • Doylestown, PA 18901 • 1-866-458-1772  
[www.RochesterPharm.com](http://www.RochesterPharm.com)

SA27-PI-0915

## **INSTRUCTIONS FOR USE**

Your health care provider has prescribed Salicylic Acid, 27.5% Wart Remover, a topical prescription preparation for the treatment of common and plantar warts. In order for Salicylic Acid, 27.5% Wart Remover to work correctly and ensure maximum benefit, the following instructions should be followed carefully. Of course, as with any medication, always consult your health care provider if you experience any discomfort or unexpected reactions.

### **STEP 1: WASH**

- Soak wart area in warm water for about 5 minutes.
- Remove any loosened tissue by gently rubbing with a brush, wash cloth, emery board, or pumice stone.
- Dry thoroughly.

### **STEP 2: BRUSH**

- Using the brush applicator supplied, apply twice to entire wart surface, allowing the first application to dry before applying the second.
- Be careful not to apply to surrounding skin.
- Leave Salicylic Acid, 27.5% Wart Remover on wart area after it dries.

Repeat Step 1 and Step 2 once or twice daily as advised by your health care provider. Clinically visible

improvement will normally occur during the first or second week of therapy.

## Label and Carton

NDC 49908-237-10

**Indications and Usage:** See insert for full Prescribing Information.  
Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Protect from freezing.  
**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

**Rx Only**  
**For topical use only**  
**Not for ophthalmic use**  
**Salicylic Acid, 27.5%**  
**Wart Remover**  
10 mL

Manufactured for:  
Rochester Pharmaceuticals • Doylestown, PA 18901  
1-866-458-1772 • [www.RochesterPharm.com](http://www.RochesterPharm.com)

**Rochester**  
PHARMACEUTICALS

09/15

10 mL

**Salicylic Acid, 27.5%**  
**Wart Remover**

Not for ophthalmic use  
For topical use only  
**Rx Only**

NDC 49908-237-10

NDC 49908-237-10

**Rx Only**  
**For topical use only**  
**Not for ophthalmic use**

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Protect from freezing.

Manufactured for:  
Rochester Pharmaceuticals  
Doylestown, PA 18901  
1-866-458-1772  
[www.RochesterPharm.com](http://www.RochesterPharm.com)

**Salicylic Acid, 27.5%**  
**Wart Remover**

**Rochester**  
PHARMACEUTICALS

3 49908 23710 2

**Ingredients:**  
Each mL of Salicylic Acid, 27.5% Wart Remover contains salicylic acid, 27.5% in a film-forming vehicle composed of acrylates copolymer, isopropyl alcohol, isobutyrate, n-butyl acetate, and polyvinyl butyral.

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

**Dosage and Administration:**  
Use as directed by your physician. See insert for full Prescribing Information.

10 mL

09/15

# SALICYLIC ACID

salicylic acid liquid

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:49908-237
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	275 mg in 1 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
ISOPROPYL ALCOHOL (UNII: ND2M416302)	

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:49908-237-10	1 in 1 CARTON	11/07/2011	
1		10 mL in 1 BOTTLE, WITH APPLICATOR; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
unapproved drug other		11/07/2011	

**Labeler** - Rochester Pharmaceuticals (134473771)

Revised: 7/2017

Rochester Pharmaceuticals