

PROFEND NASAL DECOLONIZATION- povidone iodine usp, 10% w/w swab
Professional Disposables International, Inc.

Profend Nasal Decolonization Kit

Active Ingredient

Povidone-Iodine USP 10% w/w (equivalent to 1.0% titratable iodine)

Purpose

Antiseptic

Uses

- Antiseptic for preparation of skin
- Helps reduce bacteria that can potentially cause skin infection
- A broad spectrum antiseptic for topical application for preparation of skin prior to surgery

Warnings

For external use only

Do not use

- in the eyes or mouth
- on individuals who are allergic or sensitive to iodine
- over large areas of the body

Stop use and ask doctor if irritation and redness develop or if condition persists for more than 72 hours.

Ask a doctor before use if you have

- Deep or puncture wounds
- animal bites
- serious burns

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

For nasal application:

Step 1: Use tissue to clean inside of both nostrils, including inside tips of nostrils.
Discard tissue.

Using total of four swabsticks, perform Steps 2-6 two times for right nostril and two times for left nostril.

Step 2: Hold applicator tube between thumb and forefinger at blue band with tube in vertical position with tip/handle up. With the other hand, place thumb and forefinger at base of handle (“thumb-to-thumb”).

Step 3: Bend and snap open along break line.

Step 4: Bend and snap in opposite direction to disconnect swabstick.

Step 5: Remove swabstick from tube. Discard tube.

Step 6: Insert swabstick comfortably into one nostril. For a total of 15 seconds, rotate swabstick around circumference of nostril and then rotate in the anterior nares for a minimum of 6 complete revolutions with slight pressure, covering all surfaces. Discard swabstick.

Note: Do not blow nose. If solution drips, gently wipe with a tissue.

Other information

- For hospital and professional use
- Store at room temperature (15-29 °C; 59-84 °F)
- Avoid excessive heat

Inactive Ingredients

Purified Water, Sodium Hydroxide

Questions?

call 1-800-999-6423 (M-F 9 a.m.-5 p.m. EST)

Principal Display Panel

Profend


Nasal Decolonization Kit

Povidone-Iodine USP, 10% w/w

Antiseptic

Non-sterile solution

Swabstick



box



5BX120IN

Profend® NASAL SWABSTICKS
 Reorder No. X12048 NDC 10819-3888-1

Drug Facts

Active ingredient **Purpose**
 Povidone-iodine USP, 10% w/w
 (equivalent to 1% titratable iodine).....Antiseptic

Uses
 ■ Antiseptic for preparation of skin
 ■ Helps reduce bacteria that can potentially cause skin infection
 ■ A broad spectrum antiseptic for topical application for preparation of skin prior to surgery

Warnings
For external use only.

Do not use
 ■ In the eyes or mouth
 ■ On individuals who are allergic or sensitive to iodine
 ■ Over large areas of the body

Stop use and ask doctor if irritation and redness develop or if condition persists for more than 72 hours.

Ask a doctor before use if you have
 ■ Deep or puncture wounds
 ■ Animal bites
 ■ Serious burns

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions
 For nasal application:

- **Step 1:** Use tissue to clean inside of both nostrils, including inside tips of nostrils. Discard tissue. Using total of four swabsticks, perform Steps 2-6 two times for right nostril and two times for left nostril.
- **Step 2:** Hold applicator tube between thumb and forefinger at blue band with tube in vertical position with tip/handle up. With the other hand, place thumb and forefinger at base of handle ("thumb-to-thumb").
- **Step 3:** Bend and snap open along break line.
- **Step 4:** Bend and snap in opposite direction to disconnect swabstick.
- **Step 5:** Remove swabstick from tube. Discard tube.
- **Step 6:** Insert swabstick comfortably into one nostril. For a total of 15 seconds, rotate swabstick around circumference of nostril and then rotate in the anterior nares for a minimum of 6 complete revolutions with slight pressure, covering all surfaces. Discard swabstick.

■ **Note:** Do not blow nose. If solution drips, gently wipe with a tissue.

Other information
 ■ For hospital and professional use.
 ■ Store at room temperature (15-29°C; 59-84°F).
 ■ Avoid excessive heat.

Inactive ingredients
 Purified Water, Sodium Hydroxide

Questions? call 1-800-999-6423 (M-F 9 a.m.- 5 p.m. EST)

PROFEND NASAL DECOLONIZATION

povidone iodine usp, 10% w/w swab

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:10819-3888
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POVIDONE-IODINE (UNII: 85H0HZU99M) (IODINE - UNII:9679TC07X4)	IODINE	10 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10819-3888-3	12 in 1 CARTON	02/16/2018	
1	NDC:10819-3888-2	4 in 1 BOX		
1	NDC:10819-3888-1	0.75 mL in 1 TUBE, WTH APPLICATOR; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	02/16/2018	

Labeler - Professional Disposables International, Inc. (800777117)

Establishment

Name	Address	ID/FEI	Business Operations
Professional Disposables International, Inc.		800777117	manufacture(10819-3888)

Revised: 2/2024

Professional Disposables International, Inc.