

ASPIRIN LOW STRENGTH ADULT- aspirin tablet, coated

Sam's West Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Simply Right 44-600A

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

Do not use

if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn

- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

NDC 68196-600-51

**SIMPLY RIGHT™
Health Care**

**low strength adult
aspirin 81 mg
pain reliever (NSAID)**

Compare to Bayer® Low Dose 81 mg Aspirin active ingredient†

**aspirin regimen
enteric safety coated**

365 tablets actual size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Bayer® Low Dose Aspirin.

50844 REV0613B60051

We would like to hear from you with any comments or suggestions. In the continental US or Canada, you can call us toll-free at 1.800.426.9391 from 8:30 am to 4:00 pm ET, Monday-Friday. You can also send us an email using the feedback icon at [SamsClub.com](https://www.SamsClub.com).

Distributed by:
SAM'S West, Inc.
Bentonville, AR 72716

STOP PEELING

Drug Facts (continued)

- redness or swelling is present

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

- use by expiration date on package

Inactive ingredients corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?
1-800-426-9391

BASE (S3)

Hinge will not peel away

Drug Facts (continued)

- take a blood thinning (anticoagulant) or steroid drug
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days

TOP BACK (S2)

PRINTS MIRROR IMAGE ON 231 CLEAR LAM


Hinge will not peel away

NDC 68196-600-51

SIMPLY RIGHT
Health Care

low strength adult
aspirin 81 mg
pain reliever (NSAID)

Compare to Bayer® Low Dose 81 mg Aspirin active ingredient*
aspirin regimen
enteric safety
coated

actual size

365 tablets

We would like to hear from you with any comments or suggestions. In the continental US or Canada, you can call us toll-free at 1-800-426-9391 from 8:30 am to 4:00 pm ET, Monday-Friday. You can also send us an email using the feedback icon at SamsClub.com.

NOT FOR INDIVIDUAL SALE

Distributed by:
SAM'S West, Inc.
Bentonville, AR 72716

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 81 mg (NSAID)*,	Pain reliever *nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling ■ shock ■ hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- take more or for a longer time than directed

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Low Dose Aspirin. 50844 REV0613B60051

PEEL HERE FOR MORE DRUG FACTS

TOP (S1)

0.063 CR (4)

Simply Right 44-600A

ASPIRIN LOW STRENGTH ADULT			
aspirin tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68 196-600
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg
Inactive Ingredients			
	Ingredient Name		Strength
	STARCH, CORN (UNII: O8232NY3SJ)		
	D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		

HYPROMELLOSES (UNII: 3NXW29V3WO)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	L
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68196-600-51	365 in 1 BOTTLE; Type 0: Not a Combination Product	02/12/2008	06/19/2018

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	02/12/2008	06/19/2018

Labeler - Sam's West Inc (051957769)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(68196-600)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(68196-600)