

IBUPROFEN- ibuprofen tablet, film coated
Sam's West Inc

Rite Aid 44-329

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - the common cold
 - menstrual cramps
 - backache
 - toothache
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- are age 60 or older
- take more or for a longer time than directed

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- taking any other drug
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - leg swelling
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- use by expiration date on package
- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

Compare to ADVIL®
active ingredient†

NDC 68196-291-78

Member's Mark™

ibuprofen

tablets USP, 200 mg

**pain reliever
fever reducer
(NSAID)**

actual size


600 Tablets

†This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil®. 50844 ORG122129178


DISTRIBUTED BY: SAM'S WEST, INC.,
BENTONVILLE, AR 72716

Compare to ADVIL®
active ingredient†

NDC 68196-291-78

 Member's Mark™

ibuprofen
tablets USP, 200 mg
pain reliever
fever reducer
(NSAID)


actual size
600
Tablets

Drug Facts

Active ingredient
(in each brown tablet)
ibuprofen USP,
200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purpose

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ the common cold ■ menstrual cramps ■ backache ■ toothache ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ blisters ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ are age 60 or older ■ take more or for a longer time than directed ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if

Drug Facts (continued)
you use more than directed or for longer than directed.
Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery
Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you have problems or serious side effects from taking pain relievers or fever reducers
Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug ■ under a doctor's care for any serious condition
When using this product ■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ slurred speech ■ leg swelling ■ trouble breathing ■ chest pain ■ weakness in one part or side of body ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

Drug Facts (continued)
No Print/No Varnish Lot & EXP
Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor
Other information ■ use by expiration date on package ■ store between 20°-25°C (68°-77°F) ■ avoid excessive heat 40°C (104°F)
Inactive ingredients carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide
Questions or comments? Call 1-800-426-9381 8:30 AM-4:00 PM ET, Monday-Friday

This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil®. 50844 ORG122129178

how2recycle.info

PLASTIC BOTTLE

1 93968 32812 2

DISTRIBUTED BY: SAM'S WEST, INC.,
BENTONVILLE, AR 72716

Members Mark 44-291

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68196-291
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	

MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;291
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68196-291-78	600 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	
2	NDC:68196-291-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	10/09/2015

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/24/1988	

Labeler - Sam's West Inc (051957769)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(68196-291)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(68196-291)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(68196-291)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		868734088	manufacture(68196-291) , pack(68196-291)
-------------------------	--	-----------	--

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(68196-291)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(68196-291)

Revised: 9/2023

Sam's West Inc