


MECLIZINE HYDROCHLORIDE- meclizine hydrochloride tablet
NuCare Pharmaceuticals, Inc.

MECLIZINE HYDROCHLORIDE TABLETS, USP

Rx only

DESCRIPTION

Chemically, Meclizine HCl is 1-(*p*-chloro- α -phenylbenzyl)-4-(*m*-methylbenzyl) piperazine dihydrochloride monohydrate.



20

M .W . 481 .88

Meclizine HCl Tablets, USP are available in two different strengths: 12.5 mg and 25 mg. In addition each tablet contains the following inactive ingredients: Colloidal Silicon Dioxide, Croscarmellose Sodium, Lactose Monohydrate, Magnesium Stearate, Microcrystalline Cellulose. Also, Meclizine HCl Tablets USP, 12.5 mg contains FD&C Blue #1 Aluminum Lake (11-13%) and Meclizine HCl Tablets USP, 25 mg contains D&C Yellow #10 Aluminum Lake (15-20%).

CLINICAL PHARMACOLOGY

Meclizine Hydrochloride is an antihistamine which shows marked protective activity against nebulized histamine and lethal doses of intravenously injected histamine in guinea pigs. It has a marked effect in blocking the vasodepressor response to histamine, but only a slight blocking action against acetylcholine. Its activity is relatively weak in inhibiting the spasmogenic action of histamine on isolated guinea pig ileum.

INDICATIONS AND USAGE

For the management of nausea and vomiting, and dizziness associated with motion sickness.

CONTRAINDICATIONS

Meclizine Hydrochloride is contraindicated in individuals who have shown a previous hypersensitivity to it.

WARNINGS

Since drowsiness may, on occasion, occur with use of this drug, patients should be warned of this possibility and cautioned against driving a car or operating dangerous machinery.

Patients should avoid alcoholic beverages while taking this drug.

Due to its potential anticholinergic action, this drug should be used with caution in patients with asthma, glaucoma or enlargement of the prostate gland.

PRECAUTIONS

PREGNANCY, Teratogenic Effects

Pregnancy Category B. Reproduction studies in rats have shown cleft palates at 25-50 times the human dose. Epidemiological studies in pregnant women, however, do not indicate that medicine increases the risk of abnormalities when administered during pregnancy. Despite the animal findings, it would appear that the possibility of fetal harm is remote. Nevertheless, meclizine, or any other medication, should be used during pregnancy only if clearly necessary.

Pediatric Use

Clinical studies establishing safety and effectiveness in children have not been done; therefore, usage is not recommended in children under 12 years of age.

ADVERSE REACTIONS

Drowsiness, dry mouth and, on rare occasions, blurred vision have been reported.

DOSAGE AND ADMINISTRATION

Motion Sickness

The initial dose of 25 to 50 mg of Meclizine HCl should be taken one hour prior to travel for protection against motion sickness. Thereafter, the dose may be repeated every 24 hours for the duration of the journey.

HOW SUPPLIED

12.5 mg (Blue, oval-shaped, scored, debossed with TL122)

NDC 68071-3308-2 Bottles of 20

NDC 68071-3308-3 Bottles of 30

NDC 68071-3308-6 Bottles of 60

NDC 68071-3308-9 Bottles of 90

Store at 20-25°C (68-77°F) (See USP Controlled Room Temperature).

Manufactured By:

Jubilant Cadista Pharmaceuticals Inc.

Salisbury, MD 21801, USA.

Revised 03/11

PRINCIPAL DISPLAY PANEL

NuCare Pharmaceuticals, Inc.

NDC: 68071-3308-3
Meclizine HCl 12.5mg
#30 Tablets

Each tablet contains 12.5mg of Meclizine HCl, USP

Oval Shaped Blue Scored Tablet
 Debossed: "TL 122" on the scored side

Product #: P0138030
Rx Only

Meclizine HCl 12.5mg
 Lot: 000000 NDC: 68071-3308-03
 MFR NDC: 59746-122-06 Exp.: 00-00

Meclizine HCl 12.5mg
 Lot: 000000 NDC: 68071-3308-03
 MFR NDC: 59746-122-06 Exp.: 00-00

GTIN 00368071330830
 Serial# 00000000002
 Exp. Date 00-00
 LOT#: 000000

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

WARNING: KEEP OUT OF REACH OF CHILDREN

STORE AT CONTROLLED TEMPERATURE 68-77°F.

MECLIZINE HYDROCHLORIDE

meclizine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68071-3308(NDC:59746-122)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	12.5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
ALUMINUM OXIDE (UNII: LMI26O6933)	

Product Characteristics

Color	blue	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	TL122
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68071-3308-2	20 in 1 BOTTLE; Type 0: Not a Combination Product	01/17/2018	
2	NDC:68071-3308-3	30 in 1 BOTTLE; Type 0: Not a Combination Product	01/17/2018	
3	NDC:68071-3308-6	60 in 1 BOTTLE; Type 0: Not a Combination Product	01/17/2018	
4	NDC:68071-3308-9	90 in 1 BOTTLE; Type 0: Not a Combination Product	01/17/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040659	06/04/2010	

Labeler - NuCare Pharmaceuticals, Inc. (010632300)

Establishment

Name	Address	ID/FEI	Business Operations
NuCare Pharmaceuticals, Inc.		010632300	repack(68071-3308)

Revised: 2/2021

NuCare Pharmaceuticals, Inc.