

STAFAC 50- virginiamycin powder
Phibro Animal Health

Stafac® 50
(virginiamycin)
TYPE A MEDICATED ARTICLE

Active Drug Ingredient

Virginiamycin 11%

(Contains 50 g virginiamycin activity per lb)

Ingredients:

Processed grain by-products, roughage products, calcium carbonate,
carboxymethylcellulose, mineral oil

Caution:

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive drug to use by or on the order of a licensed veterinarian.

For further manufacture only.

Follow Directions on Back Panel

Stafac is registered trademark of Phibro Animal Health Corporation

Phibro Animal Health Corporation, Teaneck, NJ 07666

Net Weight 50 lb (22.6 kg)

Distributed by:

Phibro Animal Health Corporation

Teaneck, NJ 07666

Approved by FDA under NADA # 091-467

9910000

101-9107-10

(50 g virginiamycin activity per lb)

For use in swine and poultry feeds as specified below

Directions for Use

Mixing Directions—Swine

Stafac 50 Type A Medicated Article must be thoroughly mixed in the feed to assure even distribution. Determine the appropriate amount of Stafac 50 required from the table

below. Dilute Stafac 50 with a portion of one of the feed ingredients (enough to make a quantity equal to 10 lb per ton) before final mixing.

Swine	Required level of virginiamycin activity per ton of Type C Medicated Feed (complete feed)	Required amount of Stafac 50 per ton of Type C Medicated Feed (complete feed)	Amount of virginiamycin activity provided to animal per lb of body weight per day
Treatment of swine dysentery in nonbreeding swine weighing over 120 lb	100 g for 2 weeks	2 lb	2.27 mg
Treatment and control of swine dysentery in swine weighing up to 120 lb	100 g for 2 weeks followed by 50 g	2 lb 1 lb	2.27 mg 1.14 mg
Aid in control of swine dysentery in swine weighing up to 120 lb For use in animals or on premises with a history of swine dysentery but where symptoms have not yet occurred	25 g	8 oz	0.57 mg
CAUTION: Not for use in breeding swine over 120 lbs.			

Mixing Directions—Poultry

Mix 1 lb of Stafac 50 with 9 lb of feed ingredients to make 10 lb of an intermediate mix to contain 5 g/lb virginiamycin. Add 1 lb of the intermediate feed premix for each 5 g of virginiamycin needed to medicate a ton of feed.

Poultry	Required level of virginiamycin activity per ton of Type C Medicated Feed (complete feed)	Required amount of Stafac 50 per ton of Type C Medicated Feed (complete feed)
For use in broiler chickens		
Prevention of necrotic enteritis caused by <i>Clostridium perfringens</i> susceptible to virginiamycin	20 g	180 g
Not for Use in Laying Chickens		

Storage and Handling

Feed Continuously as Sole Ration

Store At Or Below 25°C (77°F), Excursions Permitted Up to 40°C (104°F)

Close Container After Use

Restricted Drug (California) - Use Only As Directed

Stafac 50

Suppl. 50:508.BB Suppl. 975-94 BT-1.0 10xds30-0674092



virginiamycin powder

Product Type

VFD TYPE A MEDICATED ARTICLE ANIMAL
DRUG

**Item Code
(Source)**

NDC:66104-8815

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name

Basis of Strength

Strength

VIRGINIAMYCIN (UNII: C49WS9N75L) (VIRGINIAMYCIN - UNII:C49WS9N75L)

VIRGINIAMYCIN

50 g in 0.45 kg

Inactive Ingredients

Ingredient Name	Strength
RICE BRAN (UNII: R60QEP13IC)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66104-8815-0	22.6 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA091467	03/24/2010	

Labeler - Phibro Animal Health (006989008)

Registrant - Phibro Animal Health (006989008)

Revised: 11/2023

Phibro Animal Health