

ESTRUMATE- cloprostenol sodium injection
Merck Sharp & Dohme Corp.

Estrumate®
(cloprostenol injection)


250 mcg cloprostenol/mL (equivalent to 263 mcg cloprostenol sodium/mL)

A sterile solution of a prostaglandin F2 α analogue for intramuscular injection in beef cows, lactating dairy cows, and replacement beef and dairy heifers

Caution: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION:

Estrumate® (cloprostenol injection) is a synthetic prostaglandin analogue structurally related to prostaglandin F2 α (PGF2 α). Each mL of the sterile colorless aqueous solution contains 250 mcg cloprostenol (equivalent to 263 mcg cloprostenol sodium), 6.1 mg sodium citrate, 0.56 mg anhydrous citric acid, 6.7 mg sodium chloride, 20 mg benzyl alcohol, and water for injection, q.s.



INDICATIONS FOR USE:

1. For unobserved or non-detected estrus in beef cows, lactating dairy cows, and replacement beef and dairy heifers
2. For treatment of pyometra or chronic endometritis in beef cows, lactating dairy cows, and replacement beef and dairy heifers
3. For treatment of mummified fetus in beef cows, lactating dairy cows, and replacement beef and dairy heifers
4. For treatment of luteal cysts in beef cows, lactating dairy cows, and replacement beef and dairy heifers
5. For abortion of beef cows, lactating dairy cows, and replacement beef and dairy heifers
6. For estrus synchronization in beef cows, lactating dairy cows, and replacement beef and dairy heifers
7. For use with Fertagyl® (gonadorelin) to synchronize estrous cycles to allow for fixed time artificial insemination (FTAI) in lactating dairy cows.

Estrumate causes functional and morphological regression of the *corpus luteum* (luteolysis) in cattle. In normal, non-pregnant cycling animals, this effect on the life span of the *corpus luteum* usually results in estrus 2 to 5 days after treatment. In animals with prolonged luteal function (pyometra, mummified fetus, and luteal cysts), the induced luteolysis usually results in resolution of the condition and return to cyclicity. Pregnant animals may abort depending on the stage of gestation.

DOSAGE AND ADMINISTRATION:

Two mL of Estrumate (500 mcg cloprostenol) should be administered by *INTRAMUSCULAR INJECTION* using the specific dosage regimen for the indication.

20 mL bottle size: Use within 28 days of first puncture.

100 mL bottle size: Use within 28 days of first puncture and puncture a maximum of 12 times. Use only with automatic injection equipment or repeater syringe. Discard bottle after one stopper puncture with draw-off spike.

1. For unobserved or non-detected estrus in beef cows, lactating dairy cows, and replacement beef and dairy heifers

Cows and heifers which are not detected in estrus, although ovarian cyclicity continues, can be treated with Estrumate if a mature *corpus luteum* is present. Estrus is expected to occur 2 to 5 days following injection, at which time animals may be inseminated. Treated cattle should be inseminated at the usual time following detection of estrus. If estrous detection is not desirable or possible, treated animals may be inseminated twice at about 72 and 96 hours post-injection.

2. For treatment of pyometra or chronic endometritis in beef cows, lactating dairy cows, and replacement beef and dairy heifers

Damage to the reproductive tract at calving or postpartum retention of the placenta often leads to infection and inflammation of the uterus (endometritis). Under certain circumstances, this may progress into chronic endometritis with the uterus becoming distended with purulent matter. This condition, commonly referred to as pyometra, is characterized by a lack of cyclical estrous behavior and the presence of a persistent *corpus luteum*. Induction of luteolysis with Estrumate usually results in evacuation of the uterus and a return to normal cyclical activity within 14 days after treatment. After 14 days post-treatment, recovery rate of treated animals will not be different than that of untreated cattle.

3. For treatment of mummified fetus in beef cows, lactating dairy cows, and replacement beef and dairy heifers

Death of the conceptus during gestation may be followed by its degeneration and dehydration. Induction of luteolysis with Estrumate usually results in expulsion of the mummified fetus from the uterus. (Manual assistance may be necessary to remove the fetus from the vagina). Normal cyclical activity usually follows.

4. For treatment of luteal cysts in beef cows, lactating dairy cows, and replacement beef and dairy heifers

A cow or heifer may be noncyclic due to the presence of a luteal cyst (a single, anovulatory follicle with a thickened wall which is accompanied by no external signs and by no changes in palpable consistency of the uterus). Treatment with Estrumate can restore normal ovarian activity by causing regression of the luteal cyst.

5. For abortion of beef cows, lactating dairy cows, and replacement beef

and dairy heifers

Unwanted pregnancies can be safely and efficiently terminated from 1 week after mating until about 5 months of gestation. The induced abortion is normally uncomplicated and the fetus and placenta are usually expelled about 4 to 5 days after the injection with the reproductive tract returning to normal soon after the abortion. The ability of Estrumate to induce abortion decreases beyond the fifth month of gestation while the risk of dystocia and its consequences increases. Estrumate has not been sufficiently tested under feedlot conditions; therefore, recommendations cannot be made for its use in heifers placed in feedlots.

6. For estrus synchronization in beef cows, lactating dairy cows, and replacement beef and dairy heifers

The luteolytic action of Estrumate can be utilized to schedule estrus and ovulation for an individual cycling animal or a group of animals. This allows control of the time at which cycling cows or heifers can be bred. Estrumate can be used in a breeding program with the following methods:

- **Single Estrumate injection:** Only animals with a mature *corpus luteum* should be treated to obtain maximum response to the single injection. However, not all cycling cattle should be treated since a mature *corpus luteum* is present for only 11 to 12 days of the 21-day cycle. Prior to treatment, cattle should be examined rectally and found to be anatomically normal, be non-pregnant, and have a mature *corpus luteum*. If these criteria are met, estrus is expected to occur 2 to 5 days following injection, at which time animals may be inseminated. Treated cattle should be inseminated at the usual time following detection of estrus. If estrous detection is not desirable or possible, treated animals may be inseminated either once at about 72 hours or twice at about 72 and 96 hours post-injection. With a single injection program, it may be desirable to assess the cyclicity status of the herd before Estrumate treatment. This can be accomplished by heat detecting and breeding at the usual time following detection of estrus for a 6-day period, all prior to injection. If by the sixth day the cyclicity status appears normal (approximately 25%-30% detected in estrus), all cattle not already inseminated should be palpated for normality, non-pregnancy, and cyclicity, then injected with Estrumate. Breeding should then be continued at the usual time following signs of estrus on the seventh and eighth days. On the ninth and tenth days, breeding may continue at the usual time following detection of estrus, or all cattle not already inseminated may be bred either once on the ninth day (at about 72 hours post-injection) or on both the ninth and tenth days (at about 72 and 96 hours post-injection).
- **Double Estrumate injections:** prior to treatment, cattle should be examined rectally and found to be anatomically normal, non-pregnant, and cycling (the presence of a mature *corpus luteum* is not necessary when the first injection of a double injection regimen is given). A second injection should be given 11 days after the first injection. In normal, cycling cattle, estrus is expected 2 to 5 days following the second injection. Treated cattle should be inseminated at the usual time following detection of estrus. If estrous detection is not desirable or possible, treated animals may be inseminated either once at about 72 hours or twice at about 72 and 96 hours following the second Estrumate injection. Many animals will come into estrus following the first injection; these animals can be inseminated at the usual time following detected estrus. Animals not inseminated should receive a second injection 11 days after the first injection. Animals receiving both injections may be inseminated at the usual time following

detection of estrus or may be inseminated either once at about 72 hours or twice at about 72 and 96 hours post second injection.

Any breeding program recommended should be completed by either:

- observing animals (especially during the third week after injection) and inseminating or hand mating any animals returning to estrus, or
- turning in clean-up bull(s) 5 to 7 days after the last injection of Estrumate to cover any animals returning to estrus.

Management considerations for use of Estrumate for estrus synchronization:

A variety of programs can be designed to best meet the needs of individual management systems. A breeding program should be selected which is appropriate for the existing circumstances and management practices. Before a breeding program is planned, the producer's objectives must be examined and the producer must be made aware of the projected results and limitations. The producer and the consulting veterinarian should review the operation's breeding history, herd health, and nutritional status and agree that a breeding program is practical in the producer's specific situation. For any successful breeding program:

- cows and heifers must be normal, non-pregnant, and cycling (rectal palpation should be performed);
- cows and heifers must be in sound breeding condition and on an adequate or increasing plane of nutrition;
- proper program planning and record keeping are essential;
- if artificial insemination is used, it must be performed by competent inseminators using high-quality semen.

It is important to understand that Estrumate is effective only in animals with a mature *corpus luteum* (ovulation must have occurred at least 5 days prior to treatment). This must be considered when breeding is intended following a single Estrumate injection.

There is no difference in the fertility achieved following the single or double dosage regimen when breeding occurs at induced estrus, or at 72 and 96 hours post-treatment. Conception rates may be lower than expected in those fixed time breeding programs employing Estrumate alone which omit the second insemination (ie, the insemination at or near 96 hours). This is especially true if a fixed time insemination is used following a single Estrumate injection.

7. For use with Fertagyl® (gonadorelin) to synchronize estrous cycles to allow for fixed time artificial insemination (FTAI) in lactating dairy cows

Use in reproductive synchrony programs similar to the following:

- Administer the first Fertagyl® injection (2 mL; 86 mcg gonadorelin, as gonadorelin acetate) by intramuscular injection on Day 0.
- Administer 2 mL of Estrumate by intramuscular injection 6 to 8 days after the first Fertagyl® injection.
- Administer the second Fertagyl® injection (2 mL; 86 mcg gonadorelin, as gonadorelin acetate) 30 to 72 hours after the Estrumate injection.
- Perform FTAI 8 to 24 hours after the second Fertagyl® injection, or inseminate

cows on detected estrus using standard herd practices.

CONTRAINDICATIONS:

Do not use this drug product in pregnant cattle, unless abortion is desired.

WARNINGS AND PRECAUTIONS:

WITHDRAWAL PERIODS AND RESIDUE WARNINGS:

No milk discard or pre-slaughter drug withdrawal period is required when used according to labeling. Use of this product in excess of the approved dose may result in drug residues.

USER SAFETY WARNINGS: Not for use in humans. Keep this and all drugs out of the reach of children.

Women of childbearing age, asthmatics, and persons with bronchial and other respiratory problems should exercise extreme caution when handling this product.

Estrumate is readily absorbed through the skin and can cause abortion and/or bronchospasms. Direct contact with the skin should therefore be avoided. Accidental spillage on the skin should be washed off immediately with soap and water.

To obtain a copy of the Safety Data Sheet (SDS) or for technical assistance, contact Merck Animal Health at 1-800-211-3573 or <http://www.merck.com>

ANIMAL SAFETY WARNINGS:

As with all parenteral products, careful aseptic techniques should be employed to decrease the possibility of post-injection bacterial infection. Severe localized clostridial infections associated with injection of Estrumate have been reported. In rare instances, such infections have resulted in death. Aggressive antibiotic therapy should be employed at the first sign of infection at the injection site, whether localized or diffuse.

At 50 and 100 times the recommended dose, mild side effects may be detected in some cattle. These include increased uneasiness, slight frothing, and milk let-down.

CONTACT INFORMATION:

To report suspected adverse drug experiences, call Merck Animal Health at 1-800-211-3573. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or at <http://www.fda.gov/reportanimalae>

HOW SUPPLIED:

20 mL and 100 mL multidose vials

STORAGE, HANDLING, AND DISPOSAL:

1. Protect from light.

2. Store in carton.
3. Store at 2-30°C (36-86°F).

See FDA's website <http://www.fda.gov/safesharpsdisposal> for information on safe disposal of needles and other sharps.

Approved by FDA under NADA # 113-645

Copyright © 2017-2022 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved. Formulated in Germany

Rev. 06/2022

397732 R1

PRINCIPAL DISPLAY PANEL - 20 mL Vial Carton

NDC 0061-5995-01

10 Doses

20 mL


Estrumate®
(cloprostenol injection)

250 mcg cloprostenol/mL
(equivalent to 263 mcg
cloprostenol sodium/mL)

A sterile solution of a
prostaglandin F_{2α} analogue
for intramuscular injection
in beef cows, lactating dairy
cows, and replacement beef
and dairy heifers.

CAUTION: Federal (USA) law
restricts this drug to use by or
on the order of a licensed
veterinarian.

MERCK
Animal Health



ESTRUMATE
cloprostenol sodium injection

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:0061-5995
---------------------	--------------------------	---------------------------	---------------

Route of Administration INTRAMUSCULAR

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
cloprostenol sodium (UNII: 886SAV9675) (cloprostenol - UNII:4208238832)	cloprostenol	263 ug in 1 mL

Inactive Ingredients

Ingredient Name	Strength
Trisodium Citrate Dihydrate (UNII: B22547B95K)	
Anhydrous Citric Acid (UNII: XF417D3PSL)	
Sodium Chloride (UNII: 451W47IQ8X)	
Benzyl alcohol (UNII: LKG8494WBH)	
Water (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0061-5995-01	1 in 1 CARTON		
1		20 mL in 1 VIAL, MULTI-DOSE		
2	NDC:0061-5995-02	1 in 1 CARTON		
2		100 mL in 1 VIAL, MULTI-DOSE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA113645	02/02/1982	

Labeler - Merck Sharp & Dohme Corp. (001317601)

Establishment

Name	Address	ID/FEI	Business Operations
Vet Pharma Friesoythe GmbH		341934053	MANUFACTURE

Establishment

Name	Address	ID/FEI	Business Operations
Piramal Healthcare Uklimited		215586998	API MANUFACTURE

Establishment

Name	Address	ID/FEI	Business Operations
Cayman Pharma s.r.o.		495076551	API MANUFACTURE