

## **STOMACH RELIEF- bismuth subsalicylate tablet, chewable**

**Meijer Distribution Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Meijer Distribution, Inc. Stomach Relief Drug Facts**

#### **Active ingredient (in each tablet)**

Bismuth subsalicylate 262 mg

#### **Purposes**

Upset stomach reliever and antidiarrheal

#### **Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

#### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

#### **Do not use**

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are**

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

**When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- chew or dissolve in mouth
- adults and children 12 years and over: 2 tablets every ½ to 1 hour as needed
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**

- **each tablet contains:** calcium 190 mg
- salicylate 104 mg
- very low sodium
- store at 20-25°C (68-77°F). Avoid excessive heat over 104°F (40°C).

**Inactive ingredients**

calcium carbonate, D&C red #27 aluminum lake, flavor, magnesium stearate, mannitol, saccharin sodium, stearic acid

Questions or comments?

1-800-719-9260

**Principal Display Panel**

GREAT SIZE FOR TRAVEL

CHEWABLE

stomach relief

Upset Stomach Reliever | Anti-diarrheal

Bismuth Subsalicylate 262 mg

5 SYMPTOM RELIEF OF: Heartburn | Indigestion | Nausea | Upset Stomach | Diarrhea

Compare to Pepto-Bismol® active ingredient

6 Tablets | actual size



Tablets protected by printed inner wrap. Do not use if torn or missing.

### Drug Facts

#### Active ingredient (in each tablet)

Bismuth subsalicylate 262 mg.....Upset stomach reliever and antidiarrheal

#### Purposes

#### Uses

relieves  
 ■ travelers' diarrhea ■ diarrhea  
 ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion  
 ■ nausea ■ gas ■ belching ■ fullness

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning the blood) ■ arthritis ■ diabetes ■ gout

### Drug Facts (continued)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

#### Stop use and ask a doctor if

■ symptoms get worse  
 ■ ringing in the ears or loss of hearing occurs  
 ■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### Directions

■ chew or dissolve in mouth  
 ■ adults and children 12 years and over: 2 tablets every ½ to 1 hour as needed  
 ■ do not exceed 8 doses (16 tablets) in 24 hours  
 ■ use until diarrhea stops but not more than 2 days  
 ■ children under 12 years: ask a doctor  
 ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

#### Other information

■ each tablet contains: calcium 190 mg  
 ■ salicylate 104 mg ■ very low sodium  
 ■ store at 20-25 °C (68-77 °F). Avoid excessive heat over 104 °F (40 °C).

**Inactive ingredients** calcium carbonate, D&C red #27 aluminum lake, flavor, magnesium stearate, mannitol, saccharin sodium, stearic acid

PID 4358589



### Drug Facts (continued)

#### Questions or comments?

1-800-719-9260

\*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Pepto-Bismol®.

## STOMACH RELIEF

bismuth subsalicylate tablet, chewable

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-469
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CDI18YE)	BISMUTH SUBSALICYLATE	262 mg

### Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	

### Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	17mm
Flavor	WINTERGREEN	Imprint Code	L469
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-469-91	1 in 1 CARTON	09/30/2018	
1		6 in 1 CELLO PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	09/30/2018	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 10/2018

Meijer Distribution Inc