

UV PROTECT FACE SPF 50- zinc oxide, titanium dioxide cream
Naos USA, Inc.

UV PROTECT FACE SPF 50

Drug Facts

Active Ingredients

Zinc Oxide 20%

Titanium Dioxide 8.82%

Purpose

Sunscreen

Uses

Help prevent sunburn.

Warnings

For external use only.

Do not use

on broken skin.

When using this product

avoid contact with eyes. If contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if

rash occurs

Keep out reach of children.

If swallowed, call a poison control centre or get medical help right away.

Directions

Adults 18 years and older:

- Apply liberally and evenly 15 minutes before sun exposure.
- Reapply at least every 2 hours.
- Use a water resistant sunscreen if swimming or sweating.
- Sun protection measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad

Spectrum SPF value of 15 or higher and other sun protection measures including:

- Limit time in the sun, especially from 10 a.m. - 2 p.m.
- Wear long-sleeve shirts, pants, hats and sunglasses.
- Children under 6 months of age: Ask a doctor.

Other information

- Keep the product between 15-30°C (59-86°F)
- Protect the product in this container from excessive heat and direct sun.

Inactive Ingredients

Coco-caprylate/caprate, Caprylic/Capric Triglyceride, Methyl Methacrylate Crosspolymer, Dicaprylyl Ether, Dicaprylyl Carbonate, Silica Dimethyl Silylate, Oryza Sativa (Rice) Bran Oil, Polyhydroxystearic Acid, Tocopheryl Acetate, Stearic Acid, Alumina, Fragrance (Parfum), Stearyl Glycyrhettinate, Tocopherol, Farnesol.

Questions?

Call to 1 877 443 3762

Package Labeling:

INSTITUT
ESTHEDERM
PARIS

Pour vivre une expérience de
soin unique, des protocoles
d'exception Institut Esthederm
sont à découvrir dans nos
centres de beauté partenaires.

www.esthederm.ca
www.esthederm.us



INSTITUT
ESTHEDERM
PARIS

Discover exceptional
Institut Esthederm
protocols in partnering
beauty centers and enjoy a
unique skincare experience.

www.esthederm.ca
www.esthederm.us

**Drug Facts (continued) /
Info-médicament (suite)**

Directions / Mode d'emploi

Adults 18 years and older:

- Apply liberally and evenly 15 minutes before sun exposure.
- Reapply at least every 2 hours.
- Use a water resistant sunscreen if swimming or sweating.
- Sun protection measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: • Limit time in the sun, especially from 10 a.m. - 2 p.m.; and
- Wear long-sleeve shirts, pants, hats and sunglasses.
- Children under 6 months of age: ask a doctor. /

Adultes 18 ans et plus :

- Appliquer généreusement et uniformément 15 minutes avant l'exposition au soleil.
- Appliquer de nouveau au moins toutes les 2 heures.
- Utiliser un écran solaire résistant à l'eau lors d'une activité aquatique ou s'il y a transpiration.
- Mesures de protection contre le soleil : Passer du temps au soleil augmente le risque de cancer de la peau et du vieillissement prématuré de la peau. Afin de diminuer ce risque, il convient d'utiliser régulièrement un écran solaire avec un FPS à large spectre d'au moins 15 et d'appliquer d'autres mesures de protection contre le soleil telles que :
- Limiter le temps passé au soleil, notamment entre 10h - 14h ; et

**Drug Facts (continued) /
Info-médicament (suite)**

- Porter des chemises à manches longues, des pantalons, un chapeau et des lunettes de soleil.
- Pour les enfants de moins de 6 mois : demander l'avis d'un médecin.

Other information / Autres renseignements

- Keep the product between 15-30°C (59-86°F).
- Protect the product in this container from excessive heat and direct sun. / • Garder le produit entre 15-30°C (59-86°F).
- Tenir loin de la chaleur excessive ou des rayons du soleil.

**Inactive ingredients /
Ingrédients inactifs**

Coco-caprylate/caprate,
Caprylic/capric Triglyceride, Methyl Methacrylate Crosspolymer, Dicaprylyl Ether, Dicaprylyl Carbonate, Silica Dimethyl Silylate, Oryza Sativa (Rice) Bran Oil, Polyhydroxystearic Acid, Tocopheryl Acetate, Stearic Acid, Alumina, Fragrance (Parfum), Stearyl Glycyrhettinate, Tocopherol, Farnesol.

Questions?

Call / Composez le 1-877-443-3762

Réf : V9000 00CA

Importé et distribué par/ Imported and distributed by:
NAOS CANADA INC.
Montréal (QC) H2W 2R2 Canada
NAOS USA INC.
Wilmington, DE 19808-1609 USA

NAOS

NAOS - Human Care Enterprise
13290 Aix-en-Provence
Made in France



INSTITUT
ESTHEDERM
PARIS

UV PROTECT

CREAM / CRÈME

100% mineral filters

100% filtres minéraux

SPF / FPS 50+

Broad Spectrum
Sunscreen
Écran solaire
à large spectre

PROTECTION UVA-UVB PROTECTION

Care first. NAOS

1 fl.oz.

30 mL

NPN : 80080706

**Drug Facts /
Info-médicament**

**Active ingredients (w/w)/
Ingrédients actifs (p/p)**

Zinc Oxide 20%
Titanium Dioxide 8,82%

Purpose / Utilité

Sunscreen / Écran solaire

Use / Usage

Helps prevent sunburn. / Aide à
prévenir les coups de soleil.

Warnings / Mises en garde

For external use only.

Pour usage externe seulement.

Do not use on broken skin.
Ne pas utiliser sur une peau
érayée.

When using this product avoid
contact with eyes. If contact occurs,
rinse thoroughly with water.

Lorsque vous utilisez ce produit,
éviter tout contact avec les yeux. Le
cas échéant, bien rincer avec de
l'eau.

Stop use and ask a doctor if rash
occurs.


Cessez d'utiliser et consultez un
médecin si une éruption cutanée se
développe.

Keep out of reach of children. If
swallowed, call a poison control
centre or get medical help right
away.

Garder hors de la portée des
enfants. En cas d'ingestion, appeler
immédiatement un centre antipoison
ou obtenir une assistance médicale.



Q1ET9000CAV4



UV PROTECT FACE SPF 50

zinc oxide, titanium dioxide cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:14324-011
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC CATION - UNII:13S1S8SF37)	ZINC CATION	200 mg in 1 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	88.2 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
COCO-CAPRYLATE/CAPRATE (UNII: 8D9H4QU99H)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
DICAPRYLYL ETHER (UNII: 77JZM5516Z)	
DICAPRYLYL CARBONATE (UNII: 609A3V1SUA)	

SILICA DIMETHYL SILYLATE (UNII: EU2PSP0G0W)	
WHITE RICE (UNII: A195V20H7A)	
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
STEARYL GLYCYRRHETINATE (UNII: 3YYE6VJS0P)	
TOCOPHEROL (UNII: R0ZB2556P8)	
FARNESOL (UNII: EB41QIU6JL)	
POLYHYDROXYSTEARIC ACID (2300 MW) (UNII: YXH47AOU0F)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:14324-011-00	1 in 1 BOX	01/31/2018	09/30/2025
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	01/31/2018	09/30/2025

Labeler - Naos USA, Inc. (080727572)

Revised: 11/2023

Naos USA, Inc.