

CETIRIZINE HYDROCHLORIDE - cetirizine hydrochloride tablet
Rising Pharma Holdings, Inc.

Cetirizine Hydrochloride Tablets USP
5 mg, Allergy

ACTIVE INGREDIENTS (IN EACH TABLET)

Cetirizine HCl USP 5 mg

PURPOSE

Antihistamine

USES

Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

WARNINGS:

DO NOT USE

Do Not Use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

ASK DOCTOR

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

ASK DOCTOR/PHARMACIST

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

WHEN USING THIS PRODUCT

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery.

STOP USE

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

IF PREGNANT OR BREAST-FEEDING

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact Poison Control Center right away. (1-

800-222-1222)

DIRECTIONS

| | |
|--|--|
| Adults and children 6 years and over | 1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours |
| Adults 65 years and over | 1 tablet once a day; do not take more than 1 tablet in 24 hours |
| Children under 6 years of age | Ask a doctor |
| Consumers with liver or kidney disease | Ask a doctor |

OTHER INFORMATION

Store at 20° to 25°C (68° to 77°F)

[See USP Controlled Room Temperature].

INACTIVE INGREDIENTS

hypromellose, lactose, magnesium stearate, maize starch, polyethylene glycol, povidone, titanium dioxide.

QUESTIONS

Call 1-844-874-7464

Manufactured by:

Unique Pharmaceutical Labs,
(A Div. of J. B. Chemicals & Pharmaceuticals Ltd.),
Mumbai 400 030, India.

Distributed by:

Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816

M.L. G/1430 July 2020

129575



Cetirizine Hydrochloride Tablets USP 5 mg


Rising NDC 16571-401-10

Original Prescription Strength

Cetirizine Hydrochloride
Tablets 5 mg

100 Tablets

|  <p>NDC 16571-401-10</p> <p>Original Prescription Strength</p> <h1>Cetirizine Hydrochloride Tablets USP</h1> <p>5mg</p> <p>100 Tablets</p> | <p>24 Hour Relief of:</p> <ul style="list-style-type: none"> • Sneezing • Runny Nose • Itchy, Watery Eyes • Itchy Throat or Nose <p>Antihistamine ALLERGY Allergies</p> | <p>126407</p> <p>Manufactured by: Unique Pharmaceutical Labs. (A Div. of J.B.Chemicals & Pharmaceuticals Ltd.), Mumbai 400 030, India</p> <p>Distributed by: Rising Pharma Holdings, Inc. East Brunswick, NJ 08816</p> <p>M. L. G/1430 Jul. 2020</p> <p>Lot No.: Exp.:</p> |  <p>N 16571-40110 9</p> | | | | |
|---|--|--|--|------------------------------------|---------|-------------------------|---------------|
| <p>Drug Facts (continued)</p> <p>Other information:</p> <ul style="list-style-type: none"> ■ Store at 20° to 25° C (68° to 77° F) ■ Use USP Controlled Room Temperature. <p>Inactive ingredients hypromellose, lactose, magnesium stearate, maize starch, polyethylene glycol, povidone, titanium dioxide</p> <p>Questions? call 1-844-874-7464</p> | | <p>Do not use if imprinted foil inner seal on bottle is broken or missing</p> <p>Drug Facts</p> <table border="1"> <tr> <th>Active Ingredient (in each tablet)</th> <th>Purpose</th> </tr> <tr> <td>Cetirizine HCl USP 5 mg</td> <td>Antihistamine</td> </tr> </table> <p>Uses: Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat</p> <p style="text-align: right;">OPEN FOR FULL INFORMATION →</p> | | Active Ingredient (in each tablet) | Purpose | Cetirizine HCl USP 5 mg | Antihistamine |
| Active Ingredient (in each tablet) | Purpose | | | | | | |
| Cetirizine HCl USP 5 mg | Antihistamine | | | | | | |

|  <p>NDC 16571-401-10</p> <p>Original Prescription Strength</p> <h1>Cetirizine Hydrochloride Tablets USP</h1> <p>5mg</p> <p>100 Tablets</p> | <p>24 Hour Relief of:</p> <ul style="list-style-type: none"> • Sneezing • Runny Nose • Itchy, Watery Eyes • Itchy Throat or Nose <p>Antihistamine ALLERGY Allergies</p> | <p>Drug Facts (continued)</p> <p>Directions</p> <table border="1"> <tr> <td>Adults and children 6 years and over</td> <td>1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours.</td> </tr> <tr> <td>Adults 65 years and over</td> <td>1 tablet once a day; do not take more than 1 tablet in 24 hours.</td> </tr> <tr> <td>Children under 6 years of age</td> <td>ask a doctor</td> </tr> <tr> <td>Consumers with liver or kidney disease</td> <td>ask a doctor</td> </tr> </table> | Adults and children 6 years and over | 1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours. | Adults 65 years and over | 1 tablet once a day; do not take more than 1 tablet in 24 hours. | Children under 6 years of age | ask a doctor | Consumers with liver or kidney disease | ask a doctor | <p>Do not use if imprinted foil inner seal on bottle is broken or missing</p> <p>Drug Facts</p> <table border="1"> <tr> <th>Active Ingredient (in each tablet)</th> <th>Purpose</th> </tr> <tr> <td>Cetirizine HCl USP 5 mg</td> <td>Antihistamine</td> </tr> </table> <p>Uses: Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat</p> <p style="text-align: right;">OPEN FOR FULL INFORMATION →</p> | Active Ingredient (in each tablet) | Purpose | Cetirizine HCl USP 5 mg | Antihistamine |
|--|--|---|--------------------------------------|---|--------------------------|--|-------------------------------|--------------|--|--------------|--|------------------------------------|---------|-------------------------|---------------|
| Adults and children 6 years and over | 1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours. | | | | | | | | | | | | | | |
| Adults 65 years and over | 1 tablet once a day; do not take more than 1 tablet in 24 hours. | | | | | | | | | | | | | | |
| Children under 6 years of age | ask a doctor | | | | | | | | | | | | | | |
| Consumers with liver or kidney disease | ask a doctor | | | | | | | | | | | | | | |
| Active Ingredient (in each tablet) | Purpose | | | | | | | | | | | | | | |
| Cetirizine HCl USP 5 mg | Antihistamine | | | | | | | | | | | | | | |

| | |
|--|---|
| <p>Drug Facts (continued)</p> <p>Warnings: Do Not Use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.</p> <p>Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.</p> <p>Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. →</p> | <p>When using this product</p> <ul style="list-style-type: none"> ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery <p>Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.</p> <p>If pregnant or breast-feeding: ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) →</p> |
|--|---|

CETIRIZINE HYDROCHLORIDE

cetirizine hydrochloride tablet

| Product Information | | | |
|--|--------------------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:16571-401 |
| Route of Administration | ORAL | | |
| Active Ingredient/Active Moiety | | | |
| Ingredient Name | Basis of Strength | Strength | |
| Cetirizine Hydrochloride (UNII: 64O047KTOA) (Cetirizine - UNII:YO7261ME24) | Cetirizine Hydrochloride | 5 mg | |
| Inactive Ingredients | | | |
| Ingredient Name | Strength | | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | | | |
| lactose (UNII: J2B2A4N98G) | | | |
| magnesium stearate (UNII: 70097M6I30) | | | |
| starch, corn (UNII: O8232NY3SJ) | | | |
| polyethylene glycol (UNII: 3WJQ05DW1A) | | | |
| povidone (UNII: FZ989GH94E) | | | |
| titanium dioxide (UNII: 15FIX9V2JP) | | | |
| Product Characteristics | | | |
| Color | WHITE (White) | Score | no score |
| Shape | BULLET (Barrel Shaped) | Size | 7mm |
| Flavor | | Imprint Code | CTN;5 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:16571-401-10 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 10/01/2009 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA077829 | 10/01/2009 | |

Labeler - Rising Pharma Holdings, Inc. (835513529)**Registrant** - Unique Pharmaceutical Laboratories (917165052)**Establishment**

| Name | Address | ID/FEI | Business Operations |
|------------------------------------|---------|-----------|--|
| Unique Pharmaceutical Laboratories | | 650434645 | ANALYSIS(16571-401) , MANUFACTURE(16571-401) |

Revised: 11/2021

Rising Pharma Holdings, Inc.