

**KROGER NITETIME FLU PLUS SEVERE COLD AND COUGH- acetaminophen, diphenhydramine hydrochloride, and phenylephrine hydrochloride powder, for solution**  
**Kroger Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Kroger® NiteTime Flu + Severe Cold & Cough**

***Drug Facts***

<b><i>Active ingredients (in each packet)</i></b>	<b><i>Purposes</i></b>
Acetaminophen 650 mg	Pain reliever/fever reducer
Diphenhydramine hydrochloride 25 mg	Antihistamine/cough suppressant
Phenylephrine hydrochloride 10 mg	Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to a cold:
  - minor aches and pains
  - minor sore throat pain
  - headache
  - nasal and sinus congestion
  - runny nose
  - sneezing
  - itchy nose or throat
  - itchy, watery eyes due to hay fever
  - cough due to minor throat and bronchial irritation
- temporarily reduces fever

**Warnings**

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

**Do not use**

- in a child under 4 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

**Ask a doctor or pharmacist before use if you are taking**

- sedatives or tranquilizers
- the blood thinning drug warfarin

**When using this product**

- **do not exceed recommended dosage**
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health care professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not use more than directed**
- take every 4 hours; do not take more than 5 packets in 24 hours unless directed by a doctor

<b>Age</b>	<b>Dose</b>
children under 4 years of age	<b>do not use</b>
children 4 to under 12 years of age	do not use unless directed by a doctor
adults and children 12 years of age and over	one packet

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

**Other information**

- **each packet contains:** potassium 4 mg, sodium 27 mg
- **phenylketonurics:** contains phenylalanine 34 mg per packet
- store at controlled room temperature 20°-25°C (68°-77°F). Protect product from heat and moisture.

**Inactive ingredients**

acesulfame potassium, aspartame, citric acid anhydrous, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, flavors, maltodextrin, sodium citrate anhydrous, sucrose, and pregelatinized starch.

**Questions or Comments?**

Call **1-800-632-6900**

DISTRIBUTED BY  
THE KROGER CO.  
CINCINNATI, OHIO 45202

**PRINCIPAL DISPLAY PANEL - 6 Packet Carton**

**COMPARE TO** the active ingredients of **THERAFLU®**  
**NIGHTTIME SEVERE COLD & COUGH** \*See side panel

SEE NEW WARNINGS  
INFORMATION & DIRECTIONS

**NDC 30142-113-07**

**Kroger®**

FROM OUR FAMILY  
TO YOURS

**NiteTime**

**Flu + Severe  
Cold & Cough**

**Acetaminophen**, Pain Reliever/Fever Reducer  
Diphenhydramine HCl, Antihistamine/Cough Suppressant  
Phenylephrine HCl, Nasal Decongestant

- Nasal Congestion
- Sore Throat Pain
- Runny Nose
- Body Ache
- Fever
- Headache
- Cough
- Sneezing

• **Our Pharmacists** •  
**Recommend**

**6**  
PACKETS

**HONEY**  
**LEMON**  
INFUSED WITH  
CHAMOMILE &  
WHITE TEA FLAVORS



FROM OUR FAMILY TO YOURS



# Flu + Severe Cold & Cough

Acetaminophen, Pain Reliever/Fever Reducer  
Diphenhydramine HCl, Antihistamine/Cough Suppressant  
Phenylephrine HCl, Nasal Decongestant

COMPARE TO the active ingredients of THERAFLU® NIGHTTIME SEVERE COLD & COUGH \*See side panel SEE NEW WARNINGS INFORMATION & DIRECTIONS

NDC 30142-113-07



FROM OUR FAMILY TO YOURS



## NiteTime

# Flu + Severe Cold & Cough

Acetaminophen, Pain Reliever/Fever Reducer  
Diphenhydramine HCl, Antihistamine/Cough Suppressant  
Phenylephrine HCl, Nasal Decongestant

- Nasal Congestion
- Sore Throat Pain
- Runny Nose
- Body Ache
- Fever
- Headache
- Cough
- Sneezing



6 PACKETS

**HONEY LEMON**  
INFUSED WITH CHAMOMILE & WHITE TEA FLAVORS



TAMPER EVIDENT INNER PACKET  
DO NOT USE IF SEALED PACKET IS TORN OR BROKEN

DISTRIBUTED BY  
THE KROGER CO.  
CINCINNATI, OHIO 45202

Made in Israel

QUALITY GUARANTEE  
[www.kroger.com](http://www.kroger.com)



\*Theraflu® is a registered trademark of Novartis Consumer Health, Inc., Parsippany, NJ 07054. Novartis Consumer Health, Inc. is not affiliated with The Kroger Co. or this product.



READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.  
KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

## Drug Facts

### Active ingredients (in each packet)

Acetaminophen 650 mg.....Pain reliever/fever reducer  
Diphenhydramine hydrochloride 25 mg.....Antihistamine/cough suppressant  
Phenylephrine hydrochloride 10 mg.....Nasal decongestant

### Purposes

### Uses

- temporarily relieves these symptoms due to a cold:
  - minor aches and pains
  - minor sore throat pain
  - headache
  - nasal and sinus congestion
- runny nose
- sneezing
- itchy nose or throat
- itchy, watery eyes due to hay fever
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

### Warnings

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

- Do not use**
- in a child under 4 years of age
  - if you are allergic to acetaminophen
  - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
  - with any other product containing diphenhydramine, even one used on the skin
  - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

### Ask a doctor or pharmacist before use if you are taking

- sedatives or tranquilizers
- the blood thinning drug warfarin

### When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

## Drug Facts (continued)

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health care professional before use.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- do not use more than directed
- take every 4 hours; do not take more than 5 packets in 24 hours unless directed by a doctor

Age	Dose
<b>children under 4 years of age</b>	<b>do not use</b>
children 4 to under 12 years of age	do not use unless directed by a doctor
adults and children 12 years of age and over	one packet

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

### Other information

- each packet contains: potassium 4 mg, sodium 27 mg
- phenylketonurics: contains phenylalanine 34 mg per packet
- store at controlled room temperature 20°-25°C (68°-77°F). Protect product from heat and moisture.

**Inactive ingredients** acesulfame potassium, aspartame, citric acid anhydrous, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, flavors, maltodextrin, sodium citrate anhydrous, sucrose, and pregelatinized starch.

### Questions or Comments?

1-800-632-6900

20661-1214-2  
790

# KROGER NITETIME FLU PLUS SEVERE COLD AND COUGH

acetaminophen, diphenhydramine hydrochloride, and phenylephrine hydrochloride powder, for solution

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:30 142-113
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Acetaminophen</b> (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	650 mg
<b>Diphenhydramine hydrochloride</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	Diphenhydramine hydrochloride	25 mg
<b>Phenylephrine hydrochloride</b> (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine hydrochloride	10 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>acesulfame potassium</b> (UNII: 23OV73Q5G9)	
<b>aspartame</b> (UNII: Z0H242BBR1)	
<b>anhydrous citric acid</b> (UNII: XF417D3PSL)	
<b>D&amp;C yellow no. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C blue no. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C red no. 40</b> (UNII: WZB9127XOA)	
<b>maltodextrin</b> (UNII: 7CVR7L4A2D)	
<b>anhydrous trisodium citrate</b> (UNII: RS7A450LGA)	
<b>sucrose</b> (UNII: C151H8M554)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30 142-113-07	6 in 1 CARTON; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	06/28/2013	

**Labeler** - Kroger Company (006999528)