

**K-TAB- potassium chloride tablet, film coated, extended release
AbbVie Inc.**

**K-TAB®
(potassium chloride extended-release tablets, USP)**

DESCRIPTION

K-TAB (potassium chloride extended-release tablets) is a solid oral dosage form of potassium chloride containing 8 mEq, 10 mEq and 20 mEq of potassium chloride, USP, equivalent to 600 mg, 750 mg and 1500 mg of potassium, respectively, in a film-coated (not enteric-coated), wax matrix tablet. These formulations are intended to slow the release of potassium so that the likelihood of a high localized concentration of potassium chloride within the gastrointestinal tract is reduced. The expended inert, porous, wax/polymer matrix is not absorbed and may be excreted intact in the stool.

K-TAB tablets are an electrolyte replenisher. The chemical name is potassium chloride, and the structural formula is KCl. Potassium chloride, USP, occurs as a white, granular powder or as colorless crystals. It is odorless and has a saline taste. Its solutions are neutral to litmus. It is freely soluble in water and insoluble in alcohol.

Inactive Ingredients

8 mEq and 10 mEq Tablets

Castor oil, cellulosic polymers, colloidal silicon dioxide, D&C Yellow No. 10, magnesium stearate, paraffin, polyvinyl acetate, titanium dioxide, vanillin and vitamin E.

20 mEq Tablets

Castor oil, cellulosic polymers, colloidal silicon dioxide, magnesium stearate, paraffin, polyvinyl acetate, titanium dioxide, vanillin and vitamin E.

CLINICAL PHARMACOLOGY

Potassium ion is the principal intracellular cation of most body tissues. Potassium ions participate in a number of essential physiological processes including the maintenance of intracellular tonicity, the transmission of nerve impulses, the contraction of cardiac, skeletal and smooth muscle, and the maintenance of normal renal function.

The intracellular concentration of potassium is approximately 150 to 160 mEq per liter. The normal adult plasma concentration is 3.5 to 5 mEq per liter. An active ion transport system maintains this gradient across the plasma membrane.

Potassium is a normal dietary constituent and under steady state conditions the amount of potassium absorbed from the gastrointestinal tract is equal to the amount excreted in the urine. The usual dietary intake of potassium is 50 to 100 mEq per day.

Potassium depletion will occur whenever the rate of potassium loss through renal excretion and/or loss from the gastrointestinal tract exceeds the rate of potassium intake. Such depletion usually develops as a consequence of therapy with diuretics, primary or secondary hyperaldosteronism, diabetic ketoacidosis, or inadequate replacement of potassium in patients on prolonged parenteral nutrition. Depletion can develop rapidly with severe diarrhea, especially if associated with vomiting. Potassium depletion due to these causes is usually accompanied by a concomitant loss of chloride and is manifested by hypokalemia and metabolic alkalosis. Potassium depletion may produce weakness, fatigue, disturbances of cardiac rhythm (primarily ectopic beats), prominent U-waves in the electrocardiogram, and, in advanced cases, flaccid paralysis and/or impaired ability to concentrate urine.

If potassium depletion associated with metabolic alkalosis cannot be managed by correcting the fundamental cause of the deficiency, e.g., where the patient requires long term diuretic therapy, supplemental potassium in the form of high potassium food or potassium chloride may restore normal potassium levels.

In rare circumstances, (e.g., patients with renal tubular acidosis) potassium depletion may be associated with metabolic acidosis and hyperchloremia. In such patients potassium replacement should be accomplished with potassium salts other than the chloride, such as potassium bicarbonate, potassium citrate, potassium acetate, or potassium gluconate.

INDICATIONS AND USAGE

BECAUSE OF REPORTS OF INTESTINAL AND GASTRIC ULCERATION AND BLEEDING WITH CONTROLLED-RELEASE POTASSIUM CHLORIDE PREPARATIONS, THESE DRUGS SHOULD BE RESERVED FOR THOSE PATIENTS WHO CANNOT TOLERATE OR REFUSE TO TAKE LIQUID OR EFFERVESCENT POTASSIUM PREPARATIONS, OR FOR PATIENTS WITH WHOM THERE IS A PROBLEM OF COMPLIANCE WITH THESE PREPARATIONS.

1. For the treatment of patients with hypokalemia with or without metabolic alkalosis, in digitalis intoxication, and in patients with hypokalemic familial periodic paralysis. If hypokalemia is the result of diuretic therapy, consideration should be given to the use of a lower dose of diuretic, which may be sufficient without leading to hypokalemia.
2. For the prevention of hypokalemia in patients who would be at particular risk if hypokalemia were to develop, e.g., digitalized patients or patients with significant cardiac arrhythmias.

The use of potassium salts in patients receiving diuretics for uncomplicated essential hypertension is often unnecessary when such patients have a normal dietary pattern, and when low doses of the diuretic are used. Serum potassium should be checked periodically, however, and, if hypokalemia occurs, dietary supplementation with potassium-containing foods may be adequate to control milder cases. In more severe cases and if dose adjustment of the diuretic is ineffective or unwarranted supplementation with potassium salts may be indicated.

CONTRAINDICATIONS

Potassium supplements are contraindicated in patients with hyperkalemia since a further increase in serum potassium concentration in such patients can produce cardiac arrest. Hyperkalemia may complicate any of the following conditions: chronic renal failure, systemic acidosis such as diabetic acidosis, acute dehydration, extensive tissue breakdown as in severe burns, adrenal insufficiency, or the administration of a potassium-sparing diuretic, e.g., spironolactone, triamterene, or amiloride (see **OVERDOSAGE**).

K-TAB tablets are contraindicated in patients with known hypersensitivity to any ingredient in this product.

Controlled-release formulations of potassium chloride have produced esophageal ulceration in certain cardiac patients with esophageal compression due to an enlarged left atrium. Potassium supplementation, when indicated in such patients, should be given as a liquid preparation.

All solid oral dosage forms of potassium chloride are contraindicated in any patient in whom there is structural, pathological, e.g., diabetic gastroparesis, or pharmacologic (use of anticholinergic agents or other agents with anticholinergic properties at sufficient doses to exert anticholinergic effects) cause for arrest or delay in tablet passage through the gastrointestinal tract.

WARNINGS

Hyperkalemia (see OVERDOSAGE)

In patients with impaired mechanisms for excreting potassium, the administration of potassium salts can produce hyperkalemia and cardiac arrest. This occurs most commonly in patients given potassium intravenously, but may also occur in patients given potassium orally. Potentially fatal hyperkalemia can develop rapidly and can be asymptomatic. The use of potassium salts in patients with chronic renal disease, or any other condition which impairs potassium excretion, requires particularly careful monitoring of the serum potassium concentration and appropriate dosage adjustment.

Interaction with Potassium-Sparing Diuretics

Hypokalemia should not be treated by the concomitant administration of potassium salts and a potassium-sparing diuretic, e.g., spironolactone, triamterene, or amiloride, since the simultaneous administration of these agents can produce severe hyperkalemia.

Interaction with Angiotensin Converting Enzyme Inhibitors

Angiotensin converting enzyme (ACE) inhibitors (e.g., captopril, enalapril) will produce some potassium retention by inhibiting aldosterone production. Potassium supplements should be given to patients receiving ACE inhibitors only with close monitoring.

Gastrointestinal Lesions

Solid oral dosage forms of potassium chloride can produce ulcerative and/or stenotic lesions of the gastrointestinal tract. Based on spontaneous adverse reaction reports, enteric-coated preparations of potassium chloride are associated with an increased frequency of small bowel lesions (40-50 per 100,000 patient years) compared to sustained-release wax matrix formulations (less than one per 100,000 patient years). Because of the lack of extensive marketing experience with microencapsulated products, a comparison between such products and wax matrix or enteric-coated products is not available. K-TAB tablets consist of a wax matrix formulated to provide a controlled rate of release potassium chloride and thus to minimize the possibility of a high local concentration of potassium near the gastrointestinal wall.

Prospective trials have been conducted in normal human volunteers in which the upper gastrointestinal tract was evaluated by endoscopic inspection before and after one week of solid oral potassium chloride therapy. The ability of this model to predict events occurring in usual clinical practice is unknown. Trials which approximated usual clinical practice did not reveal any clear differences between the wax matrix and microencapsulated dosage forms. In contrast, there was a higher incidence of gastric and duodenal lesions in subjects receiving a high dose of a wax matrix controlled-release formulation under conditions which did not resemble usual or recommended clinical practice, i.e., 96 mEq per day in divided doses of potassium chloride administered, to fasted patients in the presence of an anticholinergic drug to delay gastric emptying. The upper gastrointestinal lesions observed by endoscopy were asymptomatic and were not accompanied by evidence of bleeding (hemoccult testing). The relevance of these findings to the usual conditions, i.e., nonfasting, no anticholinergic agent, and smaller doses, under which controlled-release potassium chloride products are used is uncertain. Epidemiologic studies have not identified an elevated risk, compared to microencapsulated products, for upper gastrointestinal lesions in patients receiving wax matrix formulations. K-TAB tablets should be discontinued immediately and the possibility of ulceration, obstruction or perforation considered if severe vomiting, abdominal pain, distention, or gastrointestinal bleeding occurs.

Metabolic Acidosis

Hypokalemia in patients with metabolic acidosis should be treated with an alkalinizing potassium salt such as potassium bicarbonate, potassium citrate, potassium acetate, or potassium gluconate.

PRECAUTIONS

General

The diagnosis of potassium depletion is ordinarily made by demonstrating hypokalemia in a patient with a clinical history suggesting some cause for potassium depletion. In interpreting the serum potassium level, the physician should bear in mind that acute alkalosis *per se* can produce hypokalemia in the absence of a deficit in total body potassium, while acute acidosis *per se* can increase the serum potassium concentration to within the normal range even in the presence of a reduced total body potassium. The treatment of potassium depletion, particularly in the presence of cardiac disease, renal disease, or acidosis, requires careful attention to acid-base balance and appropriate monitoring of serum electrolytes, the electrocardiogram, and the clinical status of the patient.

Information for Patients

Physicians should consider reminding the patient of the following:

To take each dose with meals and with a full glass of water or other liquid.

To take this medicine following the frequency and amount prescribed by the physician. This is especially important if the patient is also taking diuretics and/or digitalis preparations.

To check with the physician if there is trouble swallowing tablets or if the tablets seem to stick in the throat.

To check with the physician at once if tarry stools or other evidence of gastrointestinal bleeding is noticed.

To take each dose without crushing, chewing or sucking the tablets.

Laboratory Tests

When blood is drawn for analysis of plasma potassium it is important to recognize that artifactual elevations can occur after improper venipuncture technique or as a result of *in vitro* hemolysis of the sample.

Drug Interactions

Potassium-sparing diuretics, angiotensin converting enzyme inhibitors (see **WARNINGS**).

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity, mutagenicity and fertility studies in animals have not been performed. Potassium is a normal dietary constituent.

Pregnancy Category C

Animal reproduction studies have not been conducted with K-TAB tablets. It is unlikely that potassium supplementation that does not lead to hyperkalemia would have an adverse effect on the fetus or would affect reproductive capacity.

Nursing Mothers

The normal potassium ion content of human milk is about 13 mEq per liter. Since oral potassium becomes part of the body potassium pool, as long as body potassium is not excessive, the contribution of potassium chloride supplementation should have little or no effect on the level in human milk.

Pediatric Use

Safety and effectiveness in children have not been established.

Geriatric Use

Clinical Studies of K-Tab tablets did not include sufficient numbers of subjects aged 65 and over to

determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

ADVERSE REACTIONS

One of the most severe adverse effects is hyperkalemia (see **CONTRAINDICATIONS**, **WARNINGS**, and **OVERDOSAGE**). There also have been reports of upper and lower gastrointestinal conditions including obstruction, bleeding, ulceration, and perforation (see **CONTRAINDICATIONS** and **WARNINGS**).

The most common adverse reactions to oral potassium salts are nausea, vomiting, flatulence, abdominal pain/discomfort, and diarrhea. These symptoms are due to irritation of the gastrointestinal tract and are best managed by taking the dose with meals, or reducing the amount taken at one time.

Skin rash has been reported rarely.

OVERDOSAGE

The administration of oral potassium salts to persons with normal excretory mechanisms for potassium rarely causes serious hyperkalemia. However, if excretory mechanisms are impaired or if intravenous administration is too rapid, potentially fatal hyperkalemia can result (see **CONTRAINDICATIONS** and **WARNINGS**). It is important to recognize that hyperkalemia is usually asymptomatic and may be manifested only by an increased serum potassium concentration (6.5-8.0 mEq/L) and characteristic electrocardiographic changes (peaking of T-waves, loss P-waves, depression of S-T segments, and prolongation of the QT intervals). Late manifestations include muscle paralysis and cardiovascular collapse from cardiac arrest (9-12 mEq/L).

Treatment measures for hyperkalemia include the following:

1. Elimination of foods and medications containing potassium and of any agents with potassium-sparing properties;
2. Intravenous administration of 300 to 500 mL/hr of 10% dextrose solution containing 10-20 units of crystalline insulin per 1,000 mL;
3. Correction of acidosis, if present, with intravenous sodium bicarbonate;
4. Use of exchange resins, hemodialysis, or peritoneal dialysis.

In treating hyperkalemia, it should be recalled that in patients who have been stabilized on digitalis, lowering the serum potassium concentration too rapidly can produce digitalis toxicity.

The extended release feature means that absorption and toxic effects may be delayed for hours. Consider standard measures to remove any unabsorbed drug.

DOSAGE AND ADMINISTRATION

The usual dietary potassium intake by the average adult is 50 to 100 mEq per day. Potassium depletion sufficient to cause hypokalemia usually requires the loss of 200 or more mEq of potassium from the total body store.

Dosage must be adjusted to the individual needs of each patient. The dose for the prevention of

hypokalemia is typically in the range of 20 mEq per day. Doses of 40-100 mEq per day or more are used for the treatment of potassium depletion. Dosage should be divided if more than 20 mEq per day is given such that no more than 20 mEq is given in a single dose.

K-TAB tablets provide 8 mEq, 10 mEq and 20 mEq of potassium chloride.

K-TAB tablets should be taken with meals and with a glass of water or other liquid. This product should not be taken on an empty stomach because of its potential for gastric irritation (see **WARNINGS**).

NOTE: K-TAB tablets are to be swallowed whole without crushing, chewing or sucking the tablets.

HOW SUPPLIED

K-TAB (potassium chloride extended-release tablets, USP) contain 600 mg, 750 mg and 1500 mg of potassium chloride (equivalent to 8 mEq, 10 mEq and 20 mEq, respectively). K-TAB tablets are provided as extended-release Filmtab[®] tablets.

K-TAB 600 mg tablets (8 mEq) are round in shape, yellow in color, debossed with the trademark K-TAB on one side, and are supplied as follows:

- Bottles of 100 – NDC 0074-3058-41
- Bottles of 1000 – NDC 0074-3058-46

K-TAB 750 mg tablets (10 mEq) are ovaloid in shape, yellow in color, and are supplied and debossed as follows:

- Bottles of 100 – NDC 0074-3278-13 (10 on one side and the trademark K-TAB on the other side)
- Bottles of 100 – NDC 0074-7804-13 (the “a” logo on one side and the trademark K-TAB on the other side)
- Bottles of 1000 – NDC 0074-3278-19 (10 on one side and the trademark K-TAB on the other side)
- Bottles of 1000 – NDC 0074-7804-19 (the “a” logo on one side and the trademark K-TAB on the other side)

K-TAB 1500 mg tablets (20 mEq) are ovaloid in shape, white in color, debossed with the trademark K-TAB on one side, and are supplied as follows:

- Bottles of 100 – NDC 0074-3023-13
- Bottles of 500 – NDC 0074-3023-53

Recommended Storage

Store at room temperature 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature].

Filmtab[®] - Film-sealed tablets, AbbVie Inc.

©2017 AbbVie Inc.

Manufactured by:

AbbVie LTD

Barceloneta, PR 00617

For:

AbbVie Inc.

North Chicago, IL 60064

03-B463 January, 2017

NDC 0074-7804-19

K-Tab[®]

(potassium chloride)

extended-release
tablets, USP)
10 mEq (750 mg)
1000 Filmtab®
Tablets
Rx only abbvie

Exp.:
Lot:
Product of:
SN:

GTIN: 00300747804191

04-C140-R23

04C140

3 0074780419 1

NDC 0074-7804-19

K-Tab®
(potassium chloride
extended-release
tablets, USP)

10 mEq (750 mg)

1000 Filmtab®
Tablets

Rx only



abbvie

Do not accept if seal over bottle opening is broken or missing.
Each tablet contains 750 mg potassium chloride (equivalent to 10 mEq).
Store at room temperature 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature].
Dispense in a USP tight container.
See package insert for full prescribing information.
Filmtab® film-sealed tablets, AbbVie Inc.
Mfd. by AbbVie LTD
Barceloneta, PR 00617
For AbbVie Inc.
North Chicago, IL 60064, USA
See opposite side of label for Customs Country of Origin.
©AbbVie

NDC 0074-3278-13
K-Tab®
(potassium chloride
extended-release
tablets, USP)
10 mEq (750 mg)
100 Filmtab®
Tablets
Rx only abbvie

Exp. Lot SN

GTIN: 00300743278132

04C379

3 00743 27813 2

NDC 0074-3278-13

K-Tab[®]
 (potassium chloride
 extended-release
 tablets, USP)

10 mEq (750 mg)

100 Filmtab[®]
 Tablets

Rx only



abbvie

Do not accept if seal over bottle opening is broken or missing.
 Each tablet contains 750 mg potassium chloride (equivalent to 10 mEq).
Store at room temperature 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature].
 Dispense in a USP tight container.
See package insert for full prescribing information.
 Filmtab[®] film-sealed tablets, AbbVie Inc.
 Mfd. by AbbVie LTD
 Barceloneta, PR 00617
 For AbbVie Inc.
 North Chicago, IL 60064 USA
 Product of the United Kingdom
 04-C379-R1

NDC 0074-3023-13

K-Tab[®]

(potassium chloride
 extended-release
 tablets, USP)

20 mEq (1500 mg)

100 Filmtab[®]

Tablets

Rx only abbvie

Exp. Lot SN

GTIN: 00300743023138

04-B917-R3

048917

3 00743 02313 8

NDC 0074-3023-13

K-Tab[®]
 (potassium chloride
 extended-release
 tablets, USP)

20 mEq (1500 mg)

100 Filmtab[®]
 Tablets

Rx only



abbvie

Do not accept if seal over bottle opening is broken or missing.
 Each tablet contains 1500 mg potassium chloride (equivalent to 20 mEq).
Store at room temperature 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature].
 Dispense in a USP tight container.
See package insert for full prescribing information.
 Filmtab[®] film-sealed tablets, AbbVie Inc.
 Mfd. by AbbVie LTD
 Barceloneta, PR 00617
 For AbbVie Inc.
 North Chicago, IL 60064 USA
 Product of the United Kingdom
 ©AbbVie

NDC 0074-3058-41

K-Tab[®]

(potassium chloride
 extended-release
 tablets, USP)
 8 mEq (600 mg)
 100 Filmtab[®]
 Tablets
 Rx only abbvie

NDC 0074-3058-41

K-Tab[®]
 (potassium chloride
 extended-release
 tablets, USP)

8 mEq (600 mg)

100 Filmtab[®]
 Tablets

Rx only 

Do not accept if seal over bottle opening is broken or missing.
 Each tablet contains 600 mg potassium chloride (equivalent to 8 mEq).
Store at room temperature 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature].
 Dispense in a USP tight container.
See package insert for full prescribing information.
 Filmtab[®] film-sealed tablets, AbbVie Inc.
 Mfd. by AbbVie LTD
 Barceloneta, PR 00617
 For AbbVie Inc.
 North Chicago, IL 60064 USA
 Product of the United Kingdom
 04-B918-R4

Exp. Lot SN
 048918
 GTIN: 00300743058413
 3 00743 05841 3

K-TAB			
potassium chloride tablet, film coated, extended release			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0074-3278
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
POTASSIUM CHLORIDE (UNII: 660YQ98I10) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM CHLORIDE	750 mg	
Inactive Ingredients			
Ingredient Name	Strength		
CASTOR OIL (UNII: D5340Y2I9G)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
PARAFFIN (UNII: I9O0E3H2ZE)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
VANILLIN (UNII: CHI530446X)			
ALPHA-TOCOPHEROL (UNII: H4N855PNZ1)			
ETHYLCELLULOSES (UNII: 7Z8S9VYZ4B)			

POLYVINYL ACETATE (UNII: 32K497ZK2U)

Product Characteristics

Color	YELLOW	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	KTAB;10
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0074-3278-13	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2017	
2	NDC:0074-3278-19	1000 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018279	06/01/2017	

K-TAB

potassium chloride tablet, film coated, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0074-3023
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POTASSIUM CHLORIDE (UNII: 660YQ98110) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM CHLORIDE	1500 mg

Inactive Ingredients

Ingredient Name	Strength
CASTOR OIL (UNII: D5340Y2I9G)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
PARAFFIN (UNII: I9O0E3H2ZE)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
VANILLIN (UNII: CHI530446X)	
ALPHA-TOCOPHEROL (UNII: H4N855PNZ1)	
ETHYLCELLULOSES (UNII: 7Z8S9VYZ4B)	
POLYVINYL ACETATE (UNII: 32K497ZK2U)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	KTAB
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0074-3023-13	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/09/1980	
2	NDC:0074-3023-53	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/09/1980	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018279	06/09/1980	

K-TAB

potassium chloride tablet, film coated, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0074-3058
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POTASSIUM CHLORIDE (UNII: 660YQ98110) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM CHLORIDE	600 mg

Inactive Ingredients

Ingredient Name	Strength
CASTOR OIL (UNII: D5340Y2I9G)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
PARAFFIN (UNII: I9O0E3H2ZE)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
VANILLIN (UNII: CHI530446X)	
ALPHA-TOCOPHEROL (UNII: H4N855PNZ1)	
ETHYLCELLULOSES (UNII: 7Z8S9VYZ4B)	
POLYVINYL ACETATE (UNII: 32K497ZK2U)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	KTAB
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0074-3058-41	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/09/1980	
2	NDC:0074-3058-46	1000 in 1 BOTTLE; Type 0: Not a Combination Product	06/09/1980	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018279	06/09/1980	

K-TAB

potassium chloride tablet, film coated, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0074-7804
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POTASSIUM CHLORIDE (UNII: 660YQ98110) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM CHLORIDE	750 mg

Inactive Ingredients

Ingredient Name	Strength
CASTOR OIL (UNII: D5340Y2I9G)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
PARAFFIN (UNII: I9O0E3H2ZE)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
VANILLIN (UNII: CHI530446X)	
ALPHA-TOCOPHEROL (UNII: H4N855PNZ1)	
ETHYLCELLULOSES (UNII: 7Z8S9VYZ4B)	
POLYVINYL ACETATE (UNII: 32K497ZK2U)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	OVAL	Size	15mm

Flavor		Imprint Code	KTAB;a
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0074-7804-13	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/09/1980	
2	NDC:0074-7804-19	1000 in 1 BOTTLE; Type 0: Not a Combination Product	06/09/1980	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018279	06/09/1980	

Labeler - AbbVie Inc. (078458370)

Revised: 7/2017

AbbVie Inc.