EASYLAX- polyethylene glycol 3350 powder, for solution VALU MERCHANDISERS COMPANY

BC PEG

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 grams (cap filled to the indicated "17 GRAMS" line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cup marked to contain 17 grams of powder when filled

to the indicated "17 GRAMS" line

- adults and children 17 years of age and older:
 - fill to indicated "17 GRAMS" line in cap which is marked to indicate the correct dose 17 grams
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- ensure that the powder is fully dissolved before drinking.
- do not drink if there are any clumps.
- use once a day
- use no more than 7 days
- children 16 years of age or younger: ask a doctor

Other information

- store at 20° 25°C (68° 77°F)
- tamper-evident: do not use if foil seal under cap printed with "SEALED for YOUR PROTECTION" is missing, open or broken.

Inactive ingredients

none

For serious adverse events call:

1-888-710-0006

Package label



Powder for Solution

Osmotic Laxative

Relieves Occasional Constipation (irregularity) Softens Stool

- · Dissolves in any beverage
- · Sugar Free
- · Unflavored grit-free powder

NET WT 8.3 oz (238 g)

14 ONCE DAILY DOSES *This product is not manufactured or distributed by the owner of the registered trademark MiraLAX®.

PROUDLY DISTRIBUTED BY: VALUEMERCHANDISERS CO. 5000 KANSAS AVE, KANSAS CITY, KS 66106



Rev 01/2020 7116-0238-Q7-EC



Tamper-evident: do not use if foil seal under cap printed with "SEALED for YOUR PROTECTION" is missing, open or broken.

Lot No.: E.

Drug Facts (continued)

Other information

store at 20° - 25°C (68° - 77°F) tamper-evident: do not use if foil seal under cap printed with "SEALED for YOUR PROTECTION" is missing.

Inactive ingredients none

For serious adverse events call: 1-888-710-0006

This product is not manufactured or distributed by the owner of the registered trademark MiraLAX®.

PROUDLY DISTRIBUTED BY: VALU MERCHANDISERS, CO. 5000 KANSAS AVE, KANSAS CITY, KS 66106



7116-0238-Q7-EC

Lift Here

Purpose

Osmotic

Laxative

3.0000"

Polyethylene Glycol 3350 increases the frequency of bowel movements and softens the stool

Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350. 17 grams (cap filled to the indicated "17 GRAMS" line)

- relieves occasional constipation
- (irregularity) generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a

Drug Facts (continued)

Ask a doctor before use if you have

- nausea, vomiting, or abdominal pain
 a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if ou are taking a prescription drug

When using this product you may have loose, watery, more frequent stools.

Stop use and ask a doctor if

- you have rectal bleeding or your nausea bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed unless advised by your doctor
- the bottle cap is a measuring cup marked to contain 17 grams of powder when filled to the indicated "17 GRAMS" line
- adults and children 17 years of age and older:
- fill to the indicated "17 GRAMS" line in cap which is marked to indicate the correct dose 17 grams
 stir and dissolve in any 4 to 8
- ounces of beverage (cold, hot or room temperature) then drink
- ensure that the powder is fully dissolved before drinking. do not drink if there are any
- clumps.
- use once a day
 use no more than 7 days
- children 16 years of age or younger ask a doctor

EASYLAX

polyethylene glycol 3350 powder, for solution

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:63941-489

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL **POLYETHYLENE** 3350 - UNII:G2M7P15E5P)

17 g GLYCOL 3350 in 17 g

Product Characteristics

Color white Score

Shape	Size
Flavor	Imprint Code
Contains	

Pa	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
	NDC:63941-489- 14	238 g in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2018		

Marketing I	arketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
ANDA	ANDA090812	01/01/2018				

Labeler - VALU MERCHANDISERS COMPANY (868703513)

Registrant - Geri-Care Pharmaceutical Corp (611196254)

Revised: 11/2022 VALU MERCHANDISERS COMPANY