

PAIN RELIEF ASPIRIN CHEWABLE, LOW DOSE- aspirin tablet, chewable
Rite Aid Corporation

Rite Aid 44-397A

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)

- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrans hydrated, flavor, maltodextrin, sucralose

Questions or comments?

1-800-426-9391

Principal Display Panel

NDC 11822-0397-7

Talk to your doctor or other healthcare provider before using this product for your heart.

**Compare to the active ingredient in

Bayer® Chewable Low Dose Aspirin

CHEWABLE LOW DOSE

PAIN RELIEF ASPIRIN

ASPIRIN 81 mg

CHEWABLE TABLETS

PAIN RELIEVER (NSAID)

Chew or crush tablets completely before swallowing.

CHERRY FLAVOR

ACTUAL

SIZE

36 CHEWABLE TABLETS

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Chewable Low Dose Aspirin. 50844 REV0122D39707

DISTRIBUTED BY: RITE AID,
200 NEWBERRY COMMONS
ETTERS, PA 17319 **www.riteaid.com**

**SATISFACTION
GUARANTEE**

If you're not satisfied, we'll happily refund your money.



Rite Aid 44-397A

PAIN RELIEF ASPIRIN CHEWABLE, LOW DOSE

aspirin tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0397
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Drug Facts KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredient (in each chewable tablet) **Purpose**
 Aspirin 81 mg (NSAID) *..... Pain reliever, nonsteroidal anti-inflammatory drug

Uses For the temporary relief of minor aches and pains or as recommended by your doctor.

Warnings
 Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:
 ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ take a blood thinning (anticoagulant) or steroid drug ■ are age 60 or older ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Drug Facts (continued)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ you have asthma ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools ■ feel faint ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing occurs ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur

These could be signs of a serious condition.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients corn starch, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrate, flavor, maltodextrin, sucralose

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Chew or crush tablets completely before swallowing

Drug Facts (continued)
Questions or comments?
 1-800-426-9391

***This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Chewable Low Dose Aspirin. 50844 REV0122D39707

DISTRIBUTED BY: RITE AID,
 200 NEUBERRY COMMONS
 ETTERS, PA 17319 www.riteaid.com

SATISFACTION GUARANTEE
If you're not satisfied, we'll happily refund your money.

No Print Area
Lot no. & Exp. Date

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

0 11822 36153 8

Ingredient Name		Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	81 mg	
Inactive Ingredients				
Ingredient Name		Strength		
STARCH, CORN (UNII: O8232NY3SJ)				
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)				
D&C RED NO. 30 (UNII: 2S42T2808B)				
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)				
MALTODEXTRIN (UNII: 7CVR7L4A2D)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
Product Characteristics				
Color	pink (Light Pink)	Score	no score	
Shape	ROUND	Size	8mm	
Flavor	CHERRY	Imprint Code	44;397	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0397-7	1 in 1 CARTON	09/04/2001	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:11822-0397-9	3 in 1 CARTON	09/04/2001	04/17/2026
2		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	09/04/2001		

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(11822-0397)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-0397)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(11822-0397)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11822-0397)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-0397)

Revised: 6/2024

Rite Aid Corporation