

**AMPROMED FOR CALVES- amprolium solution**  
**Bimeda Inc., Division of Cross Vetpharm Group**

-----  
**AmproMed™ FOR CALVES**

**(amprolium)**

**9.6% Oral Solution**

**Coccidiostat**

**FOR ORAL USE IN ANIMALS ONLY**

**KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN**

**Restricted Drug (California) – Use Only as Directed**

ANADA 200-482, Approved by FDA

**Net Contents:**

**1 Gallon (128 fl oz) (3.785 L)**

**ACTIVE INGREDIENT: amprolium.....9.6%**

**INDICATIONS:** An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

**DOSAGE AND ADMINISTRATION:**

**USE DIRECTIONS:** There are **96 mg** of Amprolium in every **1 mL** of AmproMed™ FOR CALVES

---

5 Day Treatment Protocol	21 Day Prevention Protocol
Daily Dosage: 10 mg amprolium/kg (10 mg per 2.2 lb body weight)	Daily Dosage: 5 mg amprolium/kg (5 mg per 2.2 lb body weight)
1 fl oz = 29.57 mL	8 fl oz = ½ pint (236.56 mL)
1 fl oz = 2 measuring tablespoonfuls	16 fl oz = 1 pint (473.12 mL)

---

**IN DRINKING WATER**

Mix AmproMed™ FOR CALVES daily with fresh drinking water.

**5-DAY TREATMENT:** Add AMPROMED™ FOR CALVES to drinking water at the rate of 16 fl oz/100 gallon. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add AMPROMED™ FOR CALVES to drinking water at the rate of 8 fl oz/100 gallon. At usual rates of water consumption, this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

**AS A DRENCH:** AMPROMED™ FOR CALVES drench solutions may be stored in a clean, labeled container for up to 3 days.

**5-DAY TREATMENT:** Add 3 fl oz AMPROMED™ FOR CALVES to 1 pt (16 oz) of water and, with a dosage syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2) body weight. Give daily for 5 days. Use on a herd basis only: when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of AMPROMED™ FOR CALVES to 1 pt (16 fl oz) of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

**RESIDUE WARNING:** Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**WARNING: Not for Human Use. Keep this and all drugs out of reach of children.**

The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reaction in users, to obtain more information, or to obtain a MSDS contact Bimeda at 1-888-524-6332.

**PRECAUTIONS: MAY CAUSE EYE IRRITATION.** For irritation, flush with plenty of water; get medical attention. **Sodium Benzoate 0.1% add as preservative.**

**STORAGE:** Store container in an upright position. Store between 5°C - 25°C (41°F - 77°F), with brief excursions to 40°C.

**INDICATIONS:** An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.


**DOSE AND ADMINISTRATION:**  
**USE DIRECTIONS:** There are 96 mg of amprolium in every 1 mL of AmproMed™ FOR CALVES.

<p><b>5-Day Treatment Protocol</b>  Daily Dosage: 10 mg amprolium/kg (10 mg per 2.2 lb body weight)</p> <p>1 fl oz = 29.57 mL  1 fl oz = 2 measuring tablespoons</p>	<p><b>21-Day Prevention Protocol</b>  Daily Dosage: 5 mg amprolium/kg (5 mg per 2.2 lb body weight)</p> <p>8 fl oz = 1/2 pint (236.56 mL)  16 fl oz = 1 pint (473.12 mL)</p>
--	--

**IN DRINKING WATER**  
Mix AmproMed™ FOR CALVES daily with fresh drinking water.

**5-DAY TREATMENT:** Add AmproMed™ FOR CALVES to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add AmproMed™ FOR CALVES to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption, this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.



**FOR ORAL USE IN ANIMALS ONLY**  
**KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN**

**Restricted Drug (California) – Use Only as Directed**

ANADA 200-482, Approved by FDA

**Net Contents:**  
**1 Gallon (128 fl oz) (3.785 L)**

**ACTIVE INGREDIENT:** amprolium, 9.6%

Manufactured by: Bimeda, Inc. Le Sueur, MN 56058 www.bimeda.com

**AS A DRENCH:** AmproMed™ FOR CALVES drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

**5-DAY TREATMENT:** Add 3 fl oz AmproMed™ FOR CALVES to 1 pt (16 oz) of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of AmproMed™ FOR CALVES to 1 pt (16 fl oz) of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

**RESIDUE WARNING:** Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**WARNING: Not for Human Use. Keep this and all drugs out of reach of children.** The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information, or to obtain a MSDS contact Bimeda at 1-888-524-6332.

**PRECAUTIONS: MAY CAUSE EYE IRRITATION.** For irritation, flush with plenty of water; get medical attention. **Sodium Benzoate 0.1% added as preservative.**

**STORAGE:** Store container in an upright position. Store between 5°C-25°C (41°F-77°F), with brief excursions to 40°C.

8 99855 40018 5

Lot:  
Exp:

## AMPROMED FOR CALVES

amprolium solution

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:61133-8251
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
AMPROLIUM (UNII: 95CO6N199Q) (AMPROLIUM ION - UNII:H2T307KMZR)	AMPROLIUM ION	96 mg in 1 mL

**Product Characteristics**

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:6 1133-8251-3	3785 mL in 1 JUG		
2	NDC:6 1133-8251-4	473 mL in 1 BOTTLE		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200482	07/30/2012	

**Labeler** - Bimeda Inc., Division of Cross Vetpharm Group (060492923)

**Registrant** - Bimeda, Inc., Division of Cross Vetpharm Group (060492923)

**Establishment**

Name	Address	ID/FEI	Business Operations
Bimeda, Inc., Division of Cross Vetpharm Group		060492923	manufacture

**Establishment**

Name	Address	ID/FEI	Business Operations
Zhejiang Jiuzhou		654682061	api manufacture

Revised: 5/2015

Bimeda Inc., Division of Cross Vetpharm Group