CAREALL ACETAMINOPHEN EXTRA STRENGTH- acetaminophen tablet New World Imports, Inc

Drug Facts

Acteminophen 500 mg

Pain reliever / Fever reducer

In case of accidental overdose, contact a doctor or Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

For the temporary relief of minor aches and pains associated with: the common cold, headache, toothache, muscular aches, backache, minor pain from arthritis, premenstural and menstrual cramps

Temporarily reduces fever

Liver Warning:This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include skin reddening, blisters, rash.

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if:

- Pain gets worse or lasts for more than 10 days
- Fever gets worse or lasts for more than 3 days
- New symptoms occur
- Redness or swelling is present

These could be signs of a serious condition

If pregnant of breast-feeding, ask a health professional before use.

Do not exceed recommended dosage.

Do not use more than directed (see overdose warnings)

Adults and children 12 years of age and older:

- Take 2 caplets every 6 hours while symptoms last.
- Do not take more than 6 caplets in 24 hours, unless directed by a doctor.
- Do not use more than 10 days unless directed by a doctor.

Children under 12 years of age: consult a doctor

Corn Starch, povidone, pregelatinized starch, sodium starch glycolate, stearic acid



Allergy alert. Acetaminophen may cause severe skin reactions. liver damage may occur if you take more than 4,000mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while Liver warning: This product contains acetaminophen. Severe premenstrual and menstrual cramps For the temporary relief of minor aches and pains toothache blisters associated with: ■ the common cold ■ headache UNDER CAP IS BROKEN OR MISSING Active ingredient (in each Caplet) skin reddening ■ backache ■ minor pain from arthritis temporarily reduces fever Symptoms may include: **Drug Facts** muscular aches using this product

Children under 12 years of age

days unless directed by a doctor do not take for more than hours, unless directed by a doctor take more than 6 caplets in 24 while symptoms last ■ do not ■ take 2 caplets every 6 hours

children 12 Adults and

ears and over

Other information

■ store between 20°-25°C

nactive ingredients

corn starch, povidone,

starch, sodium

This product is not manufactured or distributed by Johnson **Oruq Facts** (continued on inside) If a skin reaction occurs, stop use and seek medical help right away Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. & Johnson Consumer Inc, owner of the registered trademark Extra Strength Tylenol®

> Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well

Keep out of reach of children.

as for children even if you do not notice any signs or symptoms

Directions ■ do not take more than directed (see

3 days ■ new symptoms occur ■ redness or swelling Stop use and ask a doctor if pain gets worse or lasts

he blood thinning drug warfarin

is present. These could be signs of a serious condition more than 10 days ■ fever gets worse or lasts more than

pregnant or breast-feeding, ask a health professional

CareALL Products, 160 Athens Way, Nashville, TN 37228 www.careallprodcuts.com Distributed By:

> Drug Facts (continued) ■ if you are allergic to acetaminophen or any of the

nactive ingredients in this product

 ∞

 ∞

Ask a doctor or pharmacist before use if you are taking Ask a doctor before use if you have liver disease

CAREALL ACETAMINOPHEN EXTRA STRENGTH

acetaminophen tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51824-099
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Acetaminophen (UNII: 36209ITL9D) (Acetaminophen - UNII:36209ITL9D)	Acetaminophen	500 mg	

Inactive Ingredients			
Ingredient Name	Strength		
STARCH, CORN (UNII: O8232NY3SJ)			
Stearic Acid (UNII: 4ELV7Z65AP)			
Povidone (UNII: FZ989GH94E)			
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)			

Product Characteristics			
Color	white (White)	Score	no score
Shape	OVAL (Oval)	Size	18mm
Flavor		Imprint Code	AZ;328
Contains			

F	Packaging			
#	tem Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51824-099- 24	24 in 1 CASE	11/28/2023	
1	NDC:51824-099- 01	100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/28/2023	

Labeler - New World Imports, Inc (075372276)

Revised: 10/2023 New World Imports, Inc