

**MULTI-SYMPTOM NITETIME- acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled**  
**SPIRIT PHARMACEUTICALS LLC**

-----  
**NITE TIME COLD&FLU CAPSULES**

***Drug Facts***

<b><i>Active ingredients (in each softgel)</i></b>	<b><i>Purpose</i></b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan Hydrobromide 15 mg	Cough suppressant
Doxylamine Succinate 6.25 mg	Antihistamine

***Uses***

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches/pains
- fever
- runny nose and sneezing

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses in 24 hours, which is the maximum daily amount for this product.
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, or occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood-thinning drug warfarin.

### **When using this product**

- **do not use more than directed**
- excitability may occur especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash, or headache that lasts. These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

**Overdose warning:** Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

### **Directions**

- Take only as recommended—see **Overdose warning.**
- do not exceed 4 doses per 24 hours

adults and children 12 years of age and over	swallow 2 softgels with water every 6 hours
children 4 to under 12 years of age	ask a doctor
children under 4 years of age	<b>do not use</b>

**If taking Nighttime and Daytime softgels carefully read each label to insure correct dosing**

**Other information**

- store at room temperature 15°-30°C (59°-86°F) and avoid excessive heat
- this product does not contain phenylpropanolamine

***Inactive ingredients***

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide

***Questions or comments?***

**1-888-333-9792**

**PRINCIPAL DISPLAY PANEL**

CABINET

Compare to the active ingredients

in VICKS® NyQUIL®\*

MULTI-SYMP TOM

NITE TIME

COLD&FLU NIGHTTIME RELIEF

ACETAMINOPHEN, DEXTROMETHORPHAN HBr, DOXYLAMINE SUCCINATE

48 SOFTGELS



## MULTI-SYMPTOM NITETIME

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68210-4086
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (SOFTGELS)	<b>Size</b>	20mm

<b>Flavor</b>		<b>Imprint Code</b>	215
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68210-4086-1	4 in 1 CARTON	07/02/2020	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/02/2020	

**Labeler - SPIRIT PHARMACEUTICALS LLC (179621011)**

Revised: 12/2023

SPIRIT PHARMACEUTICALS LLC