

MARCS TWIN PACK- sodium phosphate, dibasic, unspecified form and sodium phosphate, monobasic, unspecified form enema
Marc Glassman Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Marc's® Enema Twin Pack

Drug Facts

<i>Active ingredients (in each 118 mL delivered dose)</i>	<i>Purpose</i>
Dibasic Sodium Phosphate 7 g	Saline laxative
Monobasic Sodium Phosphate 19 g	Saline laxative

Use

- relieves occasional constipation
- this product generally produces bowel movement in 1 to 5 minutes

Warnings

For rectal use only.

Dosage warning

Using more than one enema in 24 hours can be harmful.

Do not use

- on children under 2 years of age

Ask a doctor before use if you

- have already used a laxative for more than 3 days
- have kidney disease, have heart problems, or are dehydrated
- are 55 years of age or older
- are on a sodium-restricted diet
- have abdominal pain, nausea, or vomiting
- have a sudden change in bowel habits lasting more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

When using this product

- **do not use more than directed. Serious side effects may occur from excess dosage**
- do not use for more than 3 days, without asking a doctor

Stop use and ask a doctor if

- you have rectal bleeding
- you have no bowel movement within 30 minutes of enema use
- you have symptoms of dehydration (thirstiness, dizziness, vomiting, urinating less often than normal)

These symptoms may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.

Directions (or as directed by a doctor)

Single daily dosage (per 24 hours)

Do not use if taking another sodium phosphate product.

Do not use more unless directed by a doctor. See Warnings.

Adults and children 12 years of age and older	1 bottle once daily
Children 2 to under 12 years of age	Use Pediatric Enema
Children under 2 years of age	DO NOT USE

Other information

- **each 118 ml contains** 4.3 g sodium
- store at 15 to 30°C (59 to 86°F)
- additional liquids by mouth are recommended while using this product
- **Tamper Evident: DO NOT USE IF TOP OR BOTTOM FLAP OF CARTON IS TORN OR MISSING.**

Inactive ingredients

benzalkonium chloride, disodium EDTA, purified water

Questions or comments?

866-323-0107

Distributed by: Marc Glassman, Inc.
West 130th Street, Cleveland, Ohio 44130

PRINCIPAL DISPLAY PANEL - 133 mL Bottle Carton

NDC 68998-233-12

Compare to the active ingredient of FLEET® Enema

Marc's®

Ready-to-Use

Enema

RECTAL SALINE SOLUTION

- **For relief of occasional constipation**
- **Comfortable, lubricated tip**
- **Works in minutes**

- **Not Made With Natural Rubber Latex**

**TWIN
PACK**

**2 Bottles
4.5 FL OZ (133 mL) Each**

CAUTION:
REMOVE BLUE
PROTECTIVE
SHIELD FROM
ENEMA TIP
BEFORE
INSERTING
PL6002B
V11

Tamper Resistant
Do not use if safety seal on the top flap
is broken or missing

Marc's

Ready-to-Use

Enema
Rectal Saline Solution

Positioning:

Left side position: Lie on left side with knees bent and arms resting comfortably.




Knee-chest position: Kneel, then lower head and chest forward until left side of face is resting on surface with left arm folded comfortably.



**HOW TO USE
THIS ENEMA**

- Remove protective shield from enema tip before inserting.

Hold bottle upright, grasping bottle cap with fingers. Grasp protective shield with other hand and pull gently to remove.



Administering enema:

- With steady pressure, gently insert enema tip into rectum with a slight side-to-side movement, with tip pointing toward navel. Insertion may be easier if person receiving enema bears down, as if having a bowel movement. Doing so helps relax muscles around anus.
- Do not force enema tip into rectum. Stop using if tip is hard to insert. Forcing the tip into the rectum can cause injury (especially if you have hemorrhoids). If enema tip causes rectal bleeding or pain, get immediate medical care.
- Squeeze bottle until nearly all liquid is gone. It is not necessary to empty bottle completely, as it contains more liquid than needed.
- Remove tip from rectum and stay in position until urge to evacuate is strong (usually 1 to 5 minutes). If no urge is felt after 5 minutes of using, try to empty bowel. Call a doctor promptly if no liquid comes out of the rectum after 30 minutes because dehydration could occur.

**Bottle, cap and shield are
NOT MADE WITH
NATURAL RUBBER LATEX**

Allergic reaction to materials containing Latex is a growing and serious medical problem.

Distributed by: Marc Glassman, Inc.

NDC 68998-233-12

Compare to the active ingredient
of FLEET® Enema

Marc's

Ready-to-Use

Enema

RECTAL SALINE SOLUTION

- For relief of occasional constipation
- Comfortable, lubricated tip
- Works in minutes
- Not Made With Natural Rubber Latex



**TWIN
PACK**

2 Bottles

4.5 FL OZ (133 ml) Each

CARTON SEALED FOR SAFETY. IF SAFETY SEAL ON THE TOP FLAP IS BROKEN OR MISSING, DO NOT USE.

Marc's

Ready-to-Use

Enema

RECTAL SALINE SOLUTION

COMPLETE ENEMA IN A DISPOSABLE SQUEEZE BOTTLE WITH SOFT, PRE-LUBRICATED TIP.

Protective Shield Prevents Contamination

Comfortable, Lubricated Tip For Ease Of Insertion

One-Way Safety Valve Controls Flow And Prevents Reflux



Marc's

Drug Facts

Active ingredients (in each 118 mL delivered dose)	Purpose
Dibasic Sodium Phosphate 7 g	Saline laxative
Monobasic Sodium Phosphate 19 g	Saline laxative

Use

- relieves occasional constipation
- this product generally produces bowel movement in 1 to 5 minutes

Warnings

For rectal use only.

Dosage warning: Using more than one enema in 24 hours can be harmful.

Do not use ■ on children under 2 years of age

Ask a doctor before use if you

- have already used a laxative for more than 3 days
- have kidney disease, have heart problems, or are dehydrated
- are 55 years of age or older
- are on a sodium-restricted diet
- have abdominal pain, nausea, or vomiting
- have a sudden change in bowel habits lasting more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

When using this product

- do not use more than directed. Serious side effects may occur from excess dosage
- do not use for more than 3 days, without asking a doctor


Stop use and ask a doctor if

- you have rectal bleeding
- you have no bowel movement within 30 minutes of enema use
- you have symptoms of dehydration (thirstiness, dizziness, vomiting, urinating less often than normal)

These symptoms may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Easy, Soft Squeeze Bottle



* This product is not manufactured or distributed by C. B. Fleet Company Inc., owner of the registered trademark Fleet®.



If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.

Directions (or as directed by a doctor)

Single daily dosage (per 24 hours)

Do not use if taking another sodium phosphate product.

Do not use more unless directed by a doctor. See Warnings.

Adults and children 12 years of age and older	1 bottle once daily
Children 2 to under 12 years of age	Use Pediatric Enema
Children under 2 years of age	DO NOT USE

Other information

- each 118 ml contains 4.3 g sodium
- store at 15 to 30°C (59 to 86°F)
- additional liquids by mouth are recommended while using this product
- **Tamper Evident: DO NOT USE IF TOP OR BOTTOM FLAP OF CARTON IS TORN OR MISSING.**

Inactive ingredients

benzalkonium chloride, disodium EDTA, purified water

Questions or comments? 866-323-0107

MARCS TWIN PACK

sodium phosphate, dibasic, unspecified form and sodium phosphate, monobasic, unspecified form enema

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68998-233
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM (UNII: GR686LBA74) (PHOSPHATE ION - UNII: NK08V8K8HR, SODIUM CATION - UNII: LYR4M0NH37)	SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM	19 g in 133 mL
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW) (PHOSPHATE ION - UNII: NK08V8K8HR, SODIUM CATION - UNII: LYR4M0NH37)	SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM	7 g in 133 mL

Inactive Ingredients

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
WATER (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68998-233-12	2 in 1 CARTON	02/06/2017	
1		133 mL in 1 BOTTLE, PLASTIC; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	02/06/2017	

Labeler - Marc Glassman Inc. (094487477)**Establishment**

Name	Address	ID/FEI	Business Operations
Natureplex LLC		062808196	MANUFACTURE(68998-233)

Revised: 7/2018

Marc Glassman Inc.