COLD FLU SEVERE- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated AmeriSource Bergen

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

GNP-1163-2022-0802

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-

800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	 take 2 caplets every 4 hours swallow whole - do not crush, chew, or dissolve do not take more than 10 caplets in 24 hours
children under 12 years	■ ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, D&C yellow #10 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

Compare to Tylenol® Cold + Flu Severe active ingredients†

Good Neighbor Pharmacy®

NDC 46122-418-62

For Adults

Cold + Flu Severe

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl, Guaifenesin

Pain Reliever/Fever Reducer, Cough Suppressant, Nasal Decongestant, Expectorant

Relief of:

- Head + Body Aches
- Fever + Sore Throat
- Cough
- Nasal Congestion

• Mucus + Chest Congestion

24 COOL TASTE CAPLETS

Actual Size



COLD FLU SEVERE

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-418
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg	

HYDROCHLORIDE

5 mg

Inactive Ingredients				
Ingredient Name	Strength			
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)				
CROSPOVIDONE (UNII: 2S7830E561)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
ALUMINUM OXIDE (UNII: LMI26O6933)				
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MALTODEXTRIN (UNII: 7CVR7L4A2D)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POLYVINYL ALCOHOL (UNII: 532B59J990)				
POVIDONE (UNII: FZ 989GH94E)				
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				

Product Characteristics				
Color	yellow	Score	no score	
Shape	OVAL	Size	19mm	
Flavor	MINT	Imprint Code	AAA;1136	
Contains				

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:46122- 418-62	2 in 1 CARTON	07/01/2018			
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/01/2018	

Revised: 8/2022 AmeriSource Bergen