#### PEPTO-BISMOL- bismuth subsalicylate tablet Procter & Gamble Manufactura S de RL de CV

-----

Pepto BISMOL ® Caplets

**Drug Facts** 

## **Active ingredient (in each caplet)**

Bismuth subsalicylate 262 mg

#### **Purpose**

Upset stomach reliever and antidiarrheal

#### Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - o gas
  - belching
  - fullness

## Warnings

## Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

# Allergy alert:

# Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

## Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

#### Ask a doctor before use if you have

- fever
- mucus in the stool

### Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

### Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- swallow with water, do not chew
- adults and children 12 years and over:
- 2 caplets (1 dose) every ½ hour or 4 caplets (2 doses) every hour as needed for diarrhea
- 2 caplets (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 caplets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

#### Other information

- each caplet contains: calcium 27 mg
- salicylate 99 mg
- low sodium
- avoid excessive heat (over 104°F or 40°C)

# Inactive ingredients

calcium carbonate, D&C red No. 27 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, polysorbate 80, povidone, silicon dioxide, sodium starch glycolate

## **Questions?**

1-800-717-3786

TAMPER EVIDENT: Do not use if inner printed seal on bottle is broken or missing.

**Made in Mexico** 

**DIST. BY PROCTER & GAMBLE,** 

**CINCINNATI OH 45202** 

### PRINCIPAL DISPLAY PANEL - 24 Caplet Carton

Pepto Bismol ®

Bismuth Subsalicylate

Upset Stomach Reliever/

Antidiarrheal

#### **5 SYMPTOM RELIEF**

NAUSEA HEARTBURN INDIGESTION UPSET STOMACH DIARRHEA

**24 CAPLETS** 



### **PEPTO-BISMOL**

bismuth subsalicylate tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58933-476
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII: 0414PZ 4LPZ)	BISMUTH SUBSALICYLATE	262 mg	

Inactive Ingredients			
Ingredient Name	Strength		
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)			
CALCIUM CARBONATE (UNII: H0G9379FGK)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MANNITOL (UNII: 30WL53L36A)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
POLYSORBATE 80 (UNII: 6OZP39ZG8H)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)			

Product Characteristics				
Color	pink	Score	no score	
Shape	OVAL (Caplet)	Size	16mm	
Flavor		Imprint Code	PeptoBismol	
Contains				

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-476- 01	1 in 1 POUCH	06/30/2004	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:58933-476- 02	1 in 1 CARTON	06/30/2004	
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:58933-476- 03	1 in 1 CARTON	06/30/2004	
3		40 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:58933-476- 12	17852 in 1 BAG; Type 0: Not a Combination Product	06/30/2004	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	06/30/2004	

Revised: 10/2023 Procter & Gamble Manufactura S de RL de CV