

**EQUATE NAPROXEN SODIUM- naproxen sodium capsule, liquid filled**  
**Wal-Mart Stores Inc**

-----  
**Wal-Mart Naproxen Sodium Capsules, 220 mg (NSAID) Drug Facts**

**Active ingredient (in each capsule)**

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen,

or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present in the painful area
  - any new symptoms appear
  - you have difficulty swallowing
  - it feels like the capsule is stuck in your throat

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:	<ul style="list-style-type: none"><li>• take 1 capsule every 8 to 12 hours while symptoms last</li><li>• for the first dose you may take 2 capsules within the first hour</li><li>• do not exceed 2 capsules in any 8- to 12-hour period</li><li>• do not exceed 3 capsules in a 24-hour period</li></ul>
children under 12 years:	<ul style="list-style-type: none"><li>• ask a doctor</li></ul>

### Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F) avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use

### Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

### Questions or comments?

1-888-287-1915

### Package/Label Principal Display Panel

EASY OPEN CAP

Compare to the Active Ingredient of Aleve® Liquid Gels

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Strength to Last 12 Hours

Actual Size

220 mg

80 LIQUID GELS\*\*

\*\*Liquid Filled Capsules

**EASY OPEN CAP**  
 NDC 49035-098-27  
**equate™**  
 Compare to the Active Ingredient of Aleve® Liquid Gels™

**Naproxen Sodium**  
 Capsules, 220 mg (NSAID)  
 Pain Reliever/Fever Reducer

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Strength to Last 12 Hours  
 220 mg 80 LIQUID GELS  
 Actual Size  
 \*\*\*Liquid Filled Capsules

**DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**  
**Active ingredient (in each capsule)**  
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\* .....Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Purposes**  
 ■ temporarily relieves minor aches and pains due to:  
 ■ minor pain of arthritis ■ backache ■ headache ■ the common cold  
 ■ menstrual cramps ■ toothache  
 ■ temporarily reduces fever

**Warnings**  
 Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling (an allergic reaction occurs, stop use and seek medical help right away.)  
 ■ asthma (wheezing) ■ shock ■ skin redness ■ rash  
 ■ stomach bleeding warning: This product contains an NSAID, which may cause severe stomach ulcers or bleeding problems. ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product. ■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if**  
 ■ you have a history of stomach problems, such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
 ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have asthma

**Ask a doctor or pharmacist before use if you are**  
 ■ under a doctor's care for any serious condition ■ taking any other drug

**When using this product**  
 ■ take with food or milk if stomach upset occurs  
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**  
 ■ you experience any of the following signs of stomach bleeding:  
 ■ feel faint ■ vomit blood ■ have bloody or black stools  
 ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear ■ you have difficulty swallowing ■ it feels like the capsule is stuck in your throat

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** ■ do not take more than directed ■ the smallest effective dose should be used ■ drink a full glass of water with each dose  
 ■ if taken with food, this product may take longer to work

Adults and children 12 years and older:  
 ■ take 1 capsule every 8 to 12 hours while symptoms last  
 ■ for the first dose you may take 2 capsules within the first hour  
 ■ do not exceed 2 capsules in any 8- to 12-hour period  
 ■ do not exceed 3 capsules in a 24-hour period

Children under 12 years:  
 ■ ask a doctor

**Other information** ■ each capsule contains: sodium 20 mg ■ store at 20-25°C (68-77°F) avoid high humidity and excessive heat above 40°C (104°F)  
 ■ read all directions and warnings before use

**Inactive ingredients** FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

**Questions or comments? 1-888-287-1915**

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.

©2014 by Watson Pharmaceuticals, Inc.  
 HSNAPD 4612715  
 \*This product is not manufactured or distributed by Bayer Healthcare or its subsidiaries.  
 Liquid Gels. 16452

09627 2E F2

# EQUATE NAPROXEN SODIUM

naproxen sodium capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49035-098
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

## Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
LACTIC ACID (UNII: 33X04XA5AT)	
MANNITOL (UNII: 3OWL53L36A)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONES (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

## Product Characteristics

<b>Color</b>	BLUE (Clear)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	742

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-098-27	80 in 1 BOTTLE		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021920	08/09/2013	

**Labeler** - Wal-Mart Stores Inc (051957769)

Revised: 6/2014

Wal-Mart Stores Inc