

**ACETAMINOPHEN EXTRA STRENGTH- acetaminophen tablet, coated**  
**Target Corporation**

-----  
**Target 44-519-Delisted**

***Active ingredient (in each gelcap)***

Acetaminophen 500 mg

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - headache
  - the common cold
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

## **Ask a doctor before use if you have**

liver disease.

## **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

## **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## ***Directions***

- **do not take more than directed**
- adults and children 12 years and over
  - take 2 gelcaps every 6 hours while symptoms last
  - do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
  - do not take for more than 10 days unless directed by a doctor
- children under 12 years: ask a doctor

## ***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid high humidity
- use by expiration date on package

## ***Inactive ingredients***

croscarmellose sodium, D&C red #33, FD&C blue #1, FD&C red #40, gelatin, hydroxypropyl cellulose, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, stearic acid, titanium dioxide

## ***Questions?***

**Call 1-800-910-6874**

***Principal Display Panel***

NDC 11673-519-15

Compare to active ingredient in  
Extra Strength Tylenol® Rapid Release Gels\*

extra strength

**acetaminophen**

**gelcaps, 500 mg**

pain reliever/fever reducer

**up&up™**

**50 GELCAPS**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*This product is not manufactured or distributed by Johnson  
& Johnson Corporation, owner of the registered trademark  
Extra Strength Tylenol® Rapid Release Gels.

50844 REV0322C51915

094 01 0548 R05 ID225450

Distributed by Target Corporation  
Minneapolis, MN 55403

TM & ©2022 Target Brands, Inc.

NDC 11673-519-15

Compare to active ingredient in  
Extra Strength Tylenol® Rapid Release Gels\*  
extra strength

# acetaminophen gelcaps, 500 mg

pain reliever/fever reducer



50 GELCAPS



acetaminophen gelcaps 500 mg acetaminophen gelcaps 500 mg

TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

## Drug Facts

### Active ingredient

(in each gelcap)  
Acetaminophen 500 mg.....Pain reliever/  
fever reducer

### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - the common cold
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever



\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Extra Strength Tylenol® Rapid Release Gels.  
50844 REV0322C51915  
094 01 0548 R05 ID225450  
Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2022 Target Brands, Inc.

no print / no varnish area  
lot no. & exp. date

PEEL HERE FOR MORE DRUG FACTS

## Drug Facts (continued)

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
 ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

### Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease.**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

### Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
  - new symptoms occur
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
- These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

STOP PEELING

## Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- do not take more than directed
- adults and children 12 years and over
- take 2 gelcaps every 6 hours while symptoms last
- do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor
- children under 12 years: ask a doctor

### Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid high humidity
- use by expiration date on package

### Inactive ingredients

crosscarmellose sodium, D&C red #33, FD&C blue #1, FD&C red #40, gelatin, hydroxypropyl cellulose, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, stearic acid, titanium dioxide

**Questions? Call 1-800-910-6874**

Target 44-519

## ACETAMINOPHEN EXTRA STRENGTH

acetaminophen tablet, coated

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:11673-519

Route of Administration

ORAL

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	red, blue	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	L;5
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-519-15	50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/10/2004	01/30/2025
2	NDC:11673-519-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/10/2004	01/30/2025
3	NDC:11673-519-20	225 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/10/2004	02/22/2023

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	05/10/2004	01/30/2025

**Labeler** - Target Corporation (006961700)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	manufacture(11673-519) , pack(11673-519)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11673-519)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11673-519)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(11673-519)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11673-519)

Revised: 5/2024

Target Corporation