

**MINICA-S- glycerin enema**  
**Sato Pharmaceutical Co., Ltd.**

-----  
**Minica-S**

**Active ingredient** Glycerin 80%

**Purpose** Laxative

**Uses** For relief of occasional constipation (irregularly). This product generally produces bowel movement in 1/4 to 1 hour.

**Warnings**

For rectal use only

**Do not use**

■ for a period of longer than 1 week unless directed by a doctor. Frequent or prolonged use of this preparation may result in dependence on laxatives.

**Ask a doctor before use if you have**

- abdominal pain, nausea, or vomiting.
- noticed a sudden change in bowel habits that persists over a period of 2 weeks.

**When using this product**

- may cause rectal discomfort or burning sensation.

**Stop use and ask a doctor if**

■ you have rectal bleeding or failure to have a bowel movement after use of a laxative. This may indicate a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

■ Adults and children 6 years and over: Use 1 rectal enema containing 8 mL of Glycerin daily or as directed by a doctor.

Gently insert stem with tip pointing toward navel. Squeeze unit until nearly all liquid is expelled. Remove tip from rectum.

■ Children under 6 years: Ask a doctor.

**Other information**

- A small amount of liquid will remain in unit.
- Store and use at room temperature.

**Inactive ingredients** Methylparaben, purified water.

**MINICA-S** DISPOSABLE ENEMA EXP. DATE 使用期限

\* Do not purchase if printed seal is broken or missing from top or bottom of carton. LOT  
 \* もし外面の天地の封印シールがはがれていたり、なくなっている場合には、購入しないでください。製造記号

US-P

NDC 49873-401-01

DISPOSABLE ENEMA  
 (使い捨て浣腸剤)

LAXATIVE / 便秘に  
**MINICA-S**  
 ミニカS

Manufactured by  
**SATO PHARMACEUTICAL CO., LTD.**  
 1-6-27 MOTOAKASAKA MINATO-KU TOKYO, JAPAN  
 Reports of serious side effects associated with use of this product can be sent to:  
 SATO PHARMACEUTICAL, INC 20501 Earl Street, Suite 3, Torrance, CA 90503

**2 ENEMAS (8mL x 2)**  
 浣腸剤2本(0.27 FL. OZ. x 2)



**MINICA-S**  
 DISPOSABLE ENEMA  
 8mL x 2 ENEMAS  
**OTBS**

<b>医薬品表示</b> 有効成分・分量 ..... 配合目的 グリセリン 80% ..... 100mL 効能 便秘(習慣性を除く)の緩和。通常15分〜1時間後に排便を促します。 使用上の注意 直腸にのみ使用してください。 使用しないください。 ■ 医師の指示なくして、1週間以上続けて使用しないでください。腹痛・長期常用すると慣れが生じ、薬害に陥りやすくなります。 使用前に蓋部又は薬液部にご相談ください。 ■ 薬液は、必ず、凍結がある場合、 ■ 排便習慣に突然の変遷をきたし、それが2週間以上続いている場合。 本品使用の際、時に腸の不快感、灼熱感を伴う場合があります。	<b>医薬品表示 (続き)</b> 使用を中止し、医師にご相談ください。 ■ 直腸からの出血又は赤痢を使用しても効果がない場合、腫瘍な状態を示している場合があります。 ■ 小児の手の届かないところに保管してください。誤って飲み込んでしまった場合は、直ちに専門家又はポイズンコントロールセンターにご相談ください。 用法・用量 ■ 大人及び6歳以上の小児(1日1本(グリセリン8mL))または医師の指示をのりにご使用ください。 ■ パイプの先端をしっかりと肛門に挿入し、漏洩が完全に止まるまで絞ってください。 ■ 6歳未満の小児に医師にご相談ください。 他の情報 ■ 容器内に少量の薬液が残ることがあります。■ 室温で保存し、使用してください。 その他の成分 メチルパラベン、精製水
---	---

<b>Drug Facts (continued)</b> Stop use and ask a doctor if ■ you have rectal bleeding or failure to have a bowel movement after use of a laxative. This may indicate a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Directions ■ Adults and children 6 years and over: Use 1 rectal enema containing 8mL of Glycerin daily or as directed by a doctor. Gently insert stem with tip pointing toward rectum. Squeeze until nearly all liquid is expelled. Remove tip from rectum. ■ Children under 6 years: Ask a doctor. Other information ■ A small amount of liquid will remain in unit. ■ Store and use at room temperature. Inactive ingredients Methylparaben, purified water.	<b>Drug Facts</b> Active ingredient Glycerin 80%.....Laxative Uses For relief of occasional constipation (regularity). This product generally produces bowel movement in 1/4 to 1 hour. Warnings For rectal use only Do not use ■ for a period of longer than 1 week unless directed by a doctor. Frequent or prolonged use of this preparation may result in dependence on laxatives. Ask a doctor before use if you have ■ abdominal pain, nausea, or vomiting. ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks. When using this product ■ may cause rectal discomfort or burning sensation.
---	---

**MINICA-S**  
 glycerin enema

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49873-401
<b>Route of Administration</b>	RECTAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>GLYCERIN</b> (UNII: PDC6A3C00X) (GLYCERIN - UNII: PDC6A3C00X)	GLYCERIN	80 mL in 100 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>METHYLPARABEN</b> (UNII: A218C7H19T)	
<b>WATER</b> (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49873-401-01	2 in 1 CARTON	05/01/1994	
1		8 mL in 1 APPLICATOR; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	05/01/1994	

**Labeler** - Sato Pharmaceutical Co., Ltd. (690575642)

## Establishment

Name	Address	ID/FEI	Business Operations
Sato Pharmaceutical Co., Ltd.		715699133	manufacture(49873-401) , label(49873-401) , pack(49873-401)

Revised: 12/2023

Sato Pharmaceutical Co., Ltd.