

**HYDROXYPROGESTERONE CAPROATE - hydroxyprogesterone caproate injection**  
**AuroMedics Pharma LLC**


-----  
**Hydroxyprogesterone Caproate Injection, USP**  
**Rx only**

**DESCRIPTION**

Hydroxyprogesterone Caproate Injection, USP is a sterile, long-acting preparation of the caproate ester of the naturally- occurring progestational hormone, hydroxyprogesterone, in an oil solution for intramuscular use.

The chemical name for hydroxyprogesterone caproate is pregn-4-ene-3,20-dione, 17[(1-oxohexyl)oxy]. It has a molecular formula of  $C_{27}H_{40}O_4$  and a molecular weight of 428.60. Hydroxyprogesterone caproate exists as white or creamy white crystalline powder.

The structural formula is:



Each 5 mL multiple-dose vial contains hydroxyprogesterone caproate, 250 mg/mL, in castor oil (28.6% v/v) and benzyl benzoate (46 % v/v) with the preservative benzyl alcohol (2% v/v).

**CLINICAL PHARMACOLOGY**

Hydroxyprogesterone is a potent, long-acting, progestational steroid ester which transforms proliferative endothelium into secretory endothelium, induces mammary gland duct development, and inhibits the production and/or release of gonadotropic hormone; it also shows slight estrogenic, androgenic, or corticoid effects as well, but should not be relied upon for these effects.

In advanced adenocarcinoma of the uterine corpus, Hydroxyprogesterone Caproate Injection, USP in a dosage of 1,000 mg or more, one or more times each week, often induces regressive changes.

*Absorption:* Peak serum levels of hydroxyprogesterone caproate appeared after 3 to 7 days in non-pregnant female subjects following a single intramuscular injection of 1,000

mg hydroxyprogesterone caproate. The pharmacokinetics of the 250 mg dose of hydroxyprogesterone caproate has not been evaluated in a study.

*Metabolism:* The conjugated metabolites include sulfated and glucuronidated products. *In vitro* data indicate that the metabolism of hydroxyprogesterone caproate is predominantly mediated by CYP3A4 and CYP3A5.

*Excretion:* Both conjugated metabolites and free steroids are excreted in the urine and feces, with the conjugated metabolites being prominent.

### **Hepatic Impairment**

The effect of hepatic impairment on the pharmacokinetics of hydroxyprogesterone caproate has not been evaluated.

Hydroxyprogesterone caproate is extensively metabolized and hepatic impairment may reduce the elimination of hydroxyprogesterone caproate.

### **Renal Impairment**

The effect of renal impairment on the pharmacokinetics of hydroxyprogesterone caproate has not been evaluated.

## **INDICATIONS AND USAGE**

Hydroxyprogesterone Caproate Injection, USP is indicated in non-pregnant women: for the treatment of advanced adenocarcinoma of the uterine corpus (Stage III or IV); in the management of amenorrhea (primary and secondary) and abnormal uterine bleeding due to hormonal imbalance in the absence of organic pathology, such as submucous fibroids or uterine cancer; as a test for endogenous estrogen production and for the production of secretory endometrium and desquamation.

## **CONTRAINDICATIONS**

Hydroxyprogesterone caproate is contraindicated in patients with known or suspected carcinoma of the breast, other hormone-sensitive cancer, or history of these conditions; undiagnosed abnormal vaginal bleeding; liver dysfunction or disease; missed abortion, and in those with a history of hypersensitivity to the drug. Hydroxyprogesterone caproate is also contraindicated as a diagnostic test for pregnancy and in patients with current or history of thrombotic or thromboembolic disorders.

## **WARNINGS**

### **Thrombotic and Thromboembolic Events**

Discontinue the medication pending examination if there is a sudden partial or complete loss of vision or if there is a sudden onset of proptosis, diplopia, or migraine. Medication should be stopped if examination reveals papilledema or retinal vascular lesions.

### **Allergic reactions**

Hypersensitivity reaction to progestins have been reported. Hydroxyprogesterone

caproate is contraindicated in women with a history of hypersensitivity to a progestin.

### **Glucose tolerance**

Progestins may decrease glucose tolerance and the blood glucose concentration should be monitored in diabetic users.

### **PRECAUTIONS**

Detectable amounts of progestins have been identified in the milk of mothers receiving the drug. Many studies have found no adverse effects of progestins on breastfeeding performance, or on the health, growth, or development of the infant.

Because progestational drugs may cause some degree of fluid retention, conditions that might be influenced by this effect, such as asthma, migraine, epilepsy, or cardiac or renal dysfunction require careful observation.

The pretreatment physical examination should include examination of the breasts and pelvic organs and a Papanicolaou smear.

In relation to irregular bleeding which does not respond predictably to the hormone therapy, nonfunctional causes should be borne in mind and adequate diagnostic measures instituted.

Any influence of prolonged sex hormone medication on pituitary, ovarian, adrenal, hepatic, or uterine function awaits further study.

The pathologist should be advised of progesterone therapy when relevant specimens are submitted.

Patients who have a history of psychic depression should be carefully observed and the drug discontinued if the depression recurs to a serious degree.

### **Information for The Patient**

Counsel patients that hydroxyprogesterone caproate injections may cause pain, soreness, swelling, itching or bruising. Inform the patient to contact her physician if she notices increased discomfort over time, oozing of blood or fluid, or inflammatory reactions at the injection site.

### **ADVERSE REACTIONS**

A. Serious arterial thrombotic and venous thromboembolic events, including cases of pulmonary emboli (some fatal), deep vein thrombosis, myocardial infarction, and strokes, have been reported in women using progestins.

B. neuroocular lesions (e.g., retinal thrombosis and optic neuritis); nausea; vomiting; gastrointestinal symptoms (such as abdominal cramps or bloating); edema; breakthrough bleeding, spotting, or withdrawal bleeding; breast tenderness; changes in body weight (increase or decrease); headache; increase in cervical mucus; allergic rash; abscess; pain at the injection site; migraine headaches.

C. chloasma or melasma, cholestatic jaundice, rise in blood pressure, mental depression, and amenorrhea during or after treatment.

D. posttreatment anovulation, cystitis, hirsutism, loss of scalp hair, changes in libido, changes in appetite, dizziness, fatigue, backache, itching, or amenorrhea.

E. The following laboratory tests may be affected by progestins: hepatic function (increased sulfobromophthalein retention and other tests); coagulation tests (increased in prothrombin and Factors VII, VIII, IX, and X); thyroid function tests (increase in PBI and butanol extractable protein-bound iodine, decrease in T3 uptake values).

A few instances of coughing, dyspnea, constriction of the chest, and/or allergic-like reactions have occurred following hydroxyprogesterone caproate therapy; the likelihood of these occurring may be increased at higher dosage levels.

## DOSAGE AND ADMINISTRATION

Suggested dosages are presented in the therapy guide. Because of the low viscosity of the vehicle, Hydroxyprogesterone Caproate Injection, USP may be administered with a small gauge needle. Care should be taken to inject the preparation deeply into the upper outer quadrant of the gluteal muscle following the usual precautions for intramuscular injection. Since the 250 mg potency provides a high concentration in a small volume, particular care should be observed to administer the full dose.

Note: Use of a wet needle or syringe may cause the solution to become cloudy; however, this does not affect the potency of the material.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Protect from light. Store vial in its box. Store upright.

Discard any unused product within 5 weeks after first use.

## THERAPY GUIDE

CYCLIC THERAPY SCHEDULE: (28-day cycle; repeated every 4 weeks);

Day 1 of each cycle: 20 mg Estradiol Valerate Injection, USP

2 weeks after Day 1: 250 mg Hydroxyprogesterone Caproate Injection, USP and 5 mg Estradiol Valerate Injection, USP

4 weeks after Day 1. This is Day 1 of next cycle

## SUGGESTED CYCLIC REGIMEN

Indications	Dosage	Started	Repeated	Stopped	Comments
Amenorrhea (primary and secondary): Abnormal uterine bleeding due to hormonal imbalance in the absence of organic pathology, such as	375 mg	Any time	-	-	Genital malignancy should be excluded before hormone therapy is started. Hydroxyprogesterone caproate is used as a "Medical D and C" to eliminate any proliferated endometrium from

submucous fibroids or uterine cancer.					previous estrogenic action by conversion to secretory endometrium and desquamation. To determine onset of normal cyclic functions, patient should be observed for 2 or 3 cycles after cessation of therapy.
	Cyclic Therapy Schedule	After 4 days of desquamation or, if there is no bleeding, 21 days after Hydroxyprogesterone Caproate Injection, USP alone	Every 4 weeks	After 4 cycles	
Production of secretory endometrium and desquamation	Patients not on estrogen therapy: Cyclic Therapy Schedule	Any time	Every 4 weeks	When cyclic therapy is no longer required	If estrogen deficiency has been prolonged, menstruation may not occur until estrogen has been given for several months.
	Patients currently on estrogen therapy: 375 mg Hydroxy progesterone Caproate Injection, USP	Any time	-	-	
	Cyclic Therapy Schedule	After 4 days of desquamation or, if there is no bleeding, 21 days after Hydroxyprogesterone Caproate Injection, USP alone	Every 4 weeks	When cyclic therapy is no longer required	

### SUGGESTED NON-CYCLIC REGIMEN

Indications	Dosage	Started	Repeated	Stopped	Comments
Adenocarcinoma of uterine corpus in advanced stage (Stage III or IV)	1,000 mg or more	At once	1 or more times each week (1 to 7 g per week)	When relapse occurs, or after 12 weeks with no objective response	Should not be used in early stage (Stage I or II) in place of established anticancer therapy. May be used in advanced stage concomitantly with other anticancer therapy (surgery, α

					radiation, or chemotherapy, or combination of these). Treatment results reported to date have been better in histologically well-differentiated forms of endometrial adenocarcinoma. (The drug has not been adequately studied in non-endometrioid adenocarcinoma of the uterine corpus).
Test for endogenous estrogen production ("Medical D and C")	250 mg	Any time	For confirmation, 4 weeks after 1 <sup>st</sup> injection	After 2 <sup>nd</sup> injection	Non-pregnant patient with responsive endometrium; bleeding 7 to 14 days after injection indicates endogenous estrogen.

## HOW SUPPLIED

Hydroxyprogesterone Caproate Injection, USP is available in vials providing hydroxyprogesterone caproate with a potency of 250 mg per mL. The product is formulated in castor oil and 46% benzyl benzoate and containing 2% (v/v) benzyl alcohol as a preservative.

Hydroxyprogesterone Caproate Injection, USP is a sterile, clear pale yellow to yellow color oily solution and is supplied as follows:

5 mL Multiple-Dose Vial  
Packaged Individually

NDC 55150-311-01

5 mL Multiple-Dose Vials  
in a carton of 5

NDC 55150-311-05

## Storage

Hydroxyprogesterone Caproate Injection, USP should be stored at controlled room temperature 20° to 25° C (68° to 77° F). Storage at low temperatures may result in the separation of some crystalline material which redissolves readily on heating in boiling water.

Protect from light. Store vial in its box. Store upright.

Discard any unused product within 5 weeks after first use.

The vial stopper is not made with natural rubber latex.

Distributed by:

**AuroMedics Pharma LLC**  
279 Princeton-Hightstown Road  
E.Windsor, NJ 08520

Manufactured by:  
**Eugia Pharma Specialities Limited**  
Hyderabad - 500032  
India

Revised: June 2022

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 1,250 mg per 5 mL (250 mg / mL) - Container Label**

**Rx only                      NDC 55150-311-01**  
**Hydroxyprogesterone**  
**Caproate Injection, USP**  
**1,250 mg per 5 mL**  
**(250 mg / mL)**  
**For Intramuscular Use Only**  
**5 mL Multiple Dose Vial**  
**AUROMEDICS**



Rx only      NDC 55150-311-01

**Hydroxyprogesterone**  
**Caproate Injection, USP**

**1,250 mg per 5 mL**  
**(250 mg / mL)**

**For Intramuscular Use Only**  
**5 mL Multiple Dose Vial**  
**AUROMEDICS**

Sterile

Dosage and Administration: See Insert for Dosage Information.

Store at 20° to 25°C (68° to 77°F). See USP Controlled Room Temperature.

Code: TS/DRUGS/24/2015

Mfd. in India for: **AuroMedics Pharma LLC**, E. Windsor, NJ 08520



P1424003


**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 1,250 mg per 5 mL (250 mg / mL) - Container-Carton (1 vial)**

**Rx only                      NDC 55150-311-01**  
**Hydroxyprogesterone**  
**Caproate Injection, USP**  
**1,250 mg per 5 mL**  
**(250 mg / mL)**  
**For Intramuscular Use Only**  
**5 mL Multiple Dose Vial**  
**AUROMEDICS**



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 1,250 mg per 5 mL (250 mg / mL) - Container-Carton (5 vials)**

**Rx only                      NDC 55150-311-05**  
**Hydroxyprogesterone**  
**Caproate Injection, USP**  
**1,250 mg per 5 mL**  
**(250 mg / mL)**  
**For Intramuscular Use Only**  
**5 x 5 mL Multiple Dose Vials**  
**AUROMEDICS**



## HYDROXYPROGESTERONE CAPROATE

hydroxyprogesterone caproate injection

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:55150-311
<b>Route of Administration</b>	INTRAMUSCULAR		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>HYDROXYPROGESTERONE CAPROATE</b> (UNII: 276F2042F5) (HYDROXYPROGESTERONE - UNII:21807M87J2)	HYDROXYPROGESTERONE CAPROATE	1250 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>CASTOR OIL</b> (UNII: D5340Y2I9G)	
<b>BENZYL BENZOATE</b> (UNII: N863NB338G)	
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55150-311-01	1 in 1 CARTON	05/09/2019	
1		5 mL in 1 VIAL; Type 0: Not a Combination Product		
2	NDC:55150-311-05	5 in 1 CARTON	05/09/2019	
2		5 mL in 1 VIAL; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211142	05/09/2019	

**Labeler** - AuroMedics Pharma LLC (968961354)**Establishment**

Name	Address	ID/FEI	Business Operations
EUGIA Pharma Specialities Limited		872201704	ANALYSIS(55150-311) , MANUFACTURE(55150-311)

Revised: 6/2022

AuroMedics Pharma LLC