

## **TRIAMCINOLONE ACETONIDE- triamcinolone acetonide cream**

### **Bryant Ranch Prepack**

-----  
**Triamcinolone Acetonide Cream USP, 0.025%, 0.1%, 0.5%**


**For Dermatologic Use Only**

**Not for Ophthalmic Use**

**Rx Only**

### **DESCRIPTION**

The topical corticosteroids constitute a class of primarily synthetic steroids used as anti-inflammatory and anti-pruritic agents. Triamcinolone acetonide is designated chemically as pregna-1,4-diene-3,20-dione,9-fluoro-11,21-dihydroxy-16,17-[(1-methylethylidene) bis (oxy)]-, (11 $\beta$ ,16 $\alpha$ )-. C<sub>24</sub>H<sub>31</sub>FO<sub>6</sub>, and M.W. of 434.51; CAS Reg. No. 76-25-5.



Each gram of 0.025%, 0.1% and 0.5% Triamcinolone Acetonide Cream USP contains 0.25 mg, 1 mg, or 5 mg triamcinolone acetonide respectively, in a washable cream base of cetyl alcohol, cetyl esters wax, glycerin, glyceryl monostearate, isopropyl palmitate, polysorbate-60, propylene glycol, purified water, sorbic acid, and sorbitan monostearate.

### **CLINICAL PHARMACOLOGY**

Topical corticosteroids share anti-inflammatory, anti-pruritic and vasoconstrictive actions. The mechanism of anti-inflammatory activity of the topical corticosteroids is unclear. Various laboratory methods, including vasoconstrictor assays, are used to compare and predict potencies and/or clinical efficacies of the topical corticosteroids. There is some evidence to suggest that a recognizable correlation exists between vasoconstrictor potency and therapeutic efficacy in man.

### **Pharmacokinetics -**

The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle, the integrity of the epidermal barrier, and the use of occlusive dressings. Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin increase percutaneous absorption. Occlusive dressings substantially increase the percutaneous absorption of topical corticosteroids. Thus, occlusive dressings may be a valuable therapeutic adjunct for treatment of resistant dermatoses (see **DOSAGE AND ADMINISTRATION**). Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to systemically administered corticosteroids. Corticosteroids are bound to plasma proteins in varying degrees. Corticosteroids are metabolized primarily in the liver and are then excreted by the kidneys. Some of the topical corticosteroids and their metabolites are also excreted into the bile.

## **INDICATIONS AND USAGE**

Topical corticosteroids are indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses.

## **CONTRAINDICATIONS**

Topical corticosteroids are contraindicated in those patients with a history of hypersensitivity to any of the components of the preparations.

## **PRECAUTIONS**

### **General -**

Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia, and glucosuria in some patients.

Conditions which augment systemic absorption include the application of the more potent steroids, use over large surface areas, prolonged use, and the addition of occlusive dressings. Therefore, patients receiving a large dose of a potent topical steroid applied to a large surface area or under an occlusive dressing should be evaluated periodically for evidence of HPA axis suppression by using the urinary free cortisol and ACTH stimulation tests. If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent steroid.

Recovery of HPA axis function is generally prompt and complete upon discontinuation of the drug. Infrequently, signs and symptoms of steroid withdrawal may occur, requiring supplemental systemic corticosteroids. Children may absorb proportionally larger amounts of topical corticosteroids and thus be more susceptible to systemic toxicity (see **PRECAUTIONS-Pediatric Use**).

If irritation develops, topical corticosteroids should be discontinued and appropriate therapy instituted. In the presence of dermatological infections, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, the corticosteroid should be discontinued until the infection

has been adequately controlled.

### **Information for Patients**

Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. Patients should be advised not to use this medication for any disorder other than for which it was prescribed.
3. The treated skin area should not be bandaged or otherwise covered or wrapped as to be occlusive unless directed by the physician.
4. Patients should report any signs of local adverse reactions especially under occlusive dressing.
5. Parents of pediatric patients should be advised not to use tight-fitting diapers or plastic pants on a child being treated in the diaper area, as these garments may constitute occlusive dressings.

### **Laboratory Tests**

The following tests may be helpful in evaluating the HPA axis suppression:

urinary free cortisol test

ACTH stimulation test

### **Carcinogenesis, Mutagenesis, Impairment of Fertility -**

Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of topical corticosteroids. Studies to determine mutagenicity with prednisolone and hydrocortisone have had negative results

### **Pregnancy: Teratogenic Effects: Pregnancy Category C -**

Corticosteroids are generally teratogenic in laboratory animals when administered systemically at relatively low dosage levels. The more potent corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. There are no adequate and well-controlled studies in pregnant women on the teratogenic effects from topically applied corticosteroids. Therefore, topical corticosteroids should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Drugs of this class should not be used extensively on pregnant patients, in large amounts, or for prolonged periods of time.

### **Nursing Mothers -**

It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids are secreted into breast milk in quantities not likely to have a deleterious effect on the infant. Nevertheless, caution should be exercised when topical corticosteroids are administered to a nursing woman.

### **Pediatric Use -**

Pediatric patients may demonstrate greater susceptibility to topical corticosteroid-induced HPA axis suppression and Cushing's syndrome than mature patients because of a larger skin surface area to body weight ratio.

Hypothalamic-pituitary-adrenal (HPA) axis suppression, Cushing's syndrome, and intracranial hypertension have been reported in children receiving topical corticosteroids. Manifestations of adrenal suppression in children include linear growth retardation, delayed weight gain, low plasma cortisol levels, and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Administration of topical corticosteroids to children should be limited to the least amount compatible with an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of children.

## **ADVERSE REACTIONS**

The following local adverse reactions are reported infrequently with topical corticosteroids, but may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae and miliaria.

## **OVERDOSAGE**

Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects (see **PRECAUTIONS**).

## **DOSAGE AND ADMINISTRATION**

Topical corticosteroids are generally applied to the affected area as a thin film from two to four times daily depending on the severity of the condition.

Occlusive dressing may be used for the management of psoriasis or recalcitrant conditions. If an infection develops, the use of occlusive dressing should be discontinued and appropriate antimicrobial therapy instituted.

## **HOW SUPPLIED**

Triamcinolone Acetonide Cream USP, 0.025% is available as follows:

- NDC 72162-2334-2: 15 g in a TUBE
- NDC 72162-2334-4: 80 g in a TUBE

Repackaged/Relabeled by:  
Bryant Ranch Prepack, Inc.  
Burbank, CA 91504

## STORAGE

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

Manufactured By Padagis®

Yeruham, Israel

Distributed By Padagis®

Allegan, MI 49010 • [www.padagis.com](http://www.padagis.com)

Rev 01-23

2N700 RC PH1

## Triamcinolone Acetonide Cream USP, 0.025%



GTIN  
Lot  
Exp  
SN

Each gram contains: 0.25 mg triamcinolone acetonide in a washable cream base of cetyl alcohol, cetyl esters wax, glycerin, glyceryl monostearate, isopropyl palmitate, polysorbate-60, propylene glycol, purified water, sorbic acid, and sorbitan monostearate.

For external use only. Not for ophthalmic use. Keep out of reach of children.

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

Usual Dosage: Apply 2 to 4 times a day; see insert information at [www.dailymed.nlm.nih.gov](http://www.dailymed.nlm.nih.gov)

NDC 72162-2334-2

Triamcinolone Acetonide Cream USP,

0.025%



Relabeled by:  
Bryant Ranch Prepack, Inc.  
Burbank, CA 91504 USA

Rx only  
NET WT 15 g  
Manufactured by:  
Padagis



## Extended Label

**Directions for puncturing tube seal:** Remove cap. Turn cap upside down and place puncture tip onto tube seal; push down until seal is punctured. Screw cap back on to close.

## TRIAMCINOLONE ACETONIDE

triamcinolone acetonide cream

### Product Information

|                         |                         |                    |                               |
|-------------------------|-------------------------|--------------------|-------------------------------|
| Product Type            | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:72162-2334(NDC:45802-063) |
| Route of Administration | TOPICAL                 |                    |                               |

| Active Ingredient/Active Moiety                                                               |                  |                                                   |                         |                    |
|-----------------------------------------------------------------------------------------------|------------------|---------------------------------------------------|-------------------------|--------------------|
| Ingredient Name                                                                               |                  |                                                   | Basis of Strength       | Strength           |
| <b>TRIAMCINOLONE ACETONIDE</b> (UNII: F446C597KA) (TRIAMCINOLONE ACETONIDE - UNII:F446C597KA) |                  |                                                   | TRIAMCINOLONE ACETONIDE | 0.25 mg in 1 g     |
|                                                                                               |                  |                                                   |                         |                    |
| Inactive Ingredients                                                                          |                  |                                                   |                         |                    |
| Ingredient Name                                                                               |                  |                                                   |                         | Strength           |
| <b>WATER</b> (UNII: 059QF0KO0R)                                                               |                  |                                                   |                         |                    |
| <b>CETYL ALCOHOL</b> (UNII: 936JST6JCN)                                                       |                  |                                                   |                         |                    |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)                                                    |                  |                                                   |                         |                    |
| <b>GLYCERYL MONOSTEARATE</b> (UNII: 230OU9XXE4)                                               |                  |                                                   |                         |                    |
| <b>POLYSORBATE 60</b> (UNII: CAL22UVI4M)                                                      |                  |                                                   |                         |                    |
| <b>SORBITAN MONOSTEARATE</b> (UNII: NVZ4I0H58X)                                               |                  |                                                   |                         |                    |
| <b>SORBIC ACID</b> (UNII: X045WJ989B)                                                         |                  |                                                   |                         |                    |
| <b>ISOPROPYL PALMITATE</b> (UNII: 8CRQ2TH63M)                                                 |                  |                                                   |                         |                    |
| <b>CETYL ESTERS WAX</b> (UNII: D072FFP9GU)                                                    |                  |                                                   |                         |                    |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)                                                            |                  |                                                   |                         |                    |
|                                                                                               |                  |                                                   |                         |                    |
| Packaging                                                                                     |                  |                                                   |                         |                    |
| #                                                                                             | Item Code        | Package Description                               | Marketing Start Date    | Marketing End Date |
| 1                                                                                             | NDC:72162-2334-2 | 1 in 1 CARTON                                     | 06/18/2024              |                    |
| 1                                                                                             |                  | 15 g in 1 TUBE; Type 0: Not a Combination Product |                         |                    |
| 2                                                                                             | NDC:72162-2334-4 | 1 in 1 CARTON                                     | 06/18/2024              |                    |
| 2                                                                                             |                  | 80 g in 1 TUBE; Type 0: Not a Combination Product |                         |                    |
|                                                                                               |                  |                                                   |                         |                    |
| Marketing Information                                                                         |                  |                                                   |                         |                    |
| Marketing Category                                                                            |                  | Application Number or Monograph Citation          | Marketing Start Date    | Marketing End Date |
| ANDA                                                                                          |                  | ANDA086413                                        | 10/09/2006              |                    |

**Labeler** - Bryant Ranch Prepack (171714327)

**Registrant** - Bryant Ranch Prepack (171714327)

## Establishment

| Name                 | Address | ID/FEI    | Business Operations                      |
|----------------------|---------|-----------|------------------------------------------|
| Bryant Ranch Prepack |         | 171714327 | REPACK(72162-2334) , RELABEL(72162-2334) |